

Epidemiology of ASF in wild boar




Prague, Czech Republic 25-27 October 2017

0



European Commission

Tracing the origin



Georgia June 2007




N Europe

© MAPS.COM

1. LITHUANIA
2. LUXEMBOURG
3. LIECHTENSTEIN
4. CZECH REPUBLIC
5. BOSNIA & HERZEGOVINA
6. SERBIA
7. MONTENEGRO
8. KOSOVO



**ASF: 0-70 km/year
since 2007**



Few certainties

Wild boar CAN ACT AS the true epidemiological reservoir of the virus;

The virus is maintained by the wild boars independently from the infection in domestic pigs and ticks

Infected Wild boar contaminate the environment making more likely secondary outbreaks in domestic pigs (non commercial and commercial farms)

How the virus spreads

Direct e contacts (nose to nose)

Contaminated environment (infected material)

Feeding infected wild boar carcasses

Virus prevalence in infected wild boar population: **1-4,5%**

Sero-prevalence in hunted WB: **0,5-2%**

Incubation **3-5 days**



Lethality **90-95%**

70-80% found dead wild boar are virus

≈ 50 km/year is the average speed, but the virus lasts also in old infected areas

The virus **spreads** through the **geographical continuity of the wild boar population** RATHER THAN of wild boar migration


Monthly incidence of ASF in domestic pigs and wild boar






European
Commission

NO. OF CASES OF ASF IN WILD BOAR IN POLAND (2014-2016)





Higher prevalence in summer: new born animals, insectes?


Lower prevalence in winter: virus survives in carcasses

Increasing prevalence: rutting period ?





Consumers of wild boar remains



+ 19 wild boar approaches without contact

Role of insects and carcasses no ticks

Maggots could increase contacts between wild boar and infected carcasses but they have been never positive to the virus (only DNA presence but no virus): enhanced summer transmission

Scavenging insects: long attraction for wild boar, increased probability of direct contact with infected carcasses

Carcasses: virus maintenance in the environment; direct transmission to the susceptible animals

Risk of spread after introduction of the virus

Delayed diagnosis

Wild boar population size and density

Forest connectivity


Inappropriate hunting methodologies

Lack of biosecurity measures applied during hunting


Infected wild boar carcasses available for healthy wild boars

Poaching

Geographical continuity



Winter feeding increases densities




Hunting and wild boar movement

Drive hunting with dogs: increase of range size during the hunting season

Season	100% MCP				95% kernel				50% kernel			
	Median	Q ₃ -Q ₁	Mean	SE	Median	Q ₃ -Q ₁	Mean	SE	Median	Q ₃ -Q ₁	Mean	SE
Pre-hunting	80	104	88	25	66	156	98	39	4	14	10	3
Hunting	428	1360	825	358	221	696	457	192	23	68	45	16
Post-hunting	195	544	358	151	189	488	284	99	20	88	45	20

Home range displacements during the hunting season (up to 15 km)



Eur J Wildl Res (2010) 56:307–318
DOI 10.1007/s10344-009-0314-z


ORIGINAL PAPER

Do intensive drive hunts affect wild boar (*Sus scrofa*) spatial behaviour in Italy? Some evidences and management implications

Laura Scillitani • Andrea Monaco • Silvano Toso



Driven hunt with dogs – effective method to reduce the population density



DIRTY AREA




How many
wild boars?

Density dependent spread


The number of NEW INFECTED wild boar is proportional to the wild boar population size

The duration of the epidemic is proportional to the wild boar population size


Poland: tendency to spread within areas with wild boar density > 1 individual/km²



2014 – 30 cases






2015 – 53 cases



2016 – 28 cases

Metssea asustustihedus jahipiirkonniti
(isendit 1000 ha jahimaa kohta)




Density of wild boars (individuals per 10 km² of hunting ground) in hunting districts by hunters estimations (census) in spring 2016.

SAK leiud metssigadel

- 11. - 17. juuni 2016
- Varasemad

- ▭ Maakonna piir
- ▭ Jahipiirkonna piir



Can we define the threshold density?

The threshold density (n_t) is that wild boar density at which an infectious wild boar does not encounter any susceptible wild boar in due time to spread the infection


Duration of infectiousness

Density/availability of susceptible hosts


If the wild boar population size is decreased till a certain density, the infection fade out through a density dependent mechanism

NO WILD BOARS = NO DISEASE


CLASSICAL SWINE FEVER in WILD BOAR



Density dependence of ASF



Apparently: not a density dependent spread



ASF in wild boar

A density dependent transmission during summer-autumn (new born and adult animals)....insects?

Virus survival during winter with few (or many) **infected carcasses** according to the local ecological situation

A mixed transmission: density dependent and frequency dependent => **NO THRESHOLD**


ASF in wild boar

The question is:


Which is the wild boar density that prevent the contact between a susceptible wild boar with an infected carcass?

An ASF virus will overwinter in a infected carcass.....3-4 months...and the virus will appear again during the late spring in alive susceptible individuals

CSF: a density dependent disease



ASF is not a truly density dependent infection. The final tail of the infection is determined by carcasses



Practically

ASF in wild boar eradication is PROBABILISTIC EVENT (stochastic) NOT a DETERMINISTIC one;

Eradication probability increases when: **wild boar population** size is **reduced** (as much as possible); **carcasses** are safely **disposed** (as much as possible); **hunting** is carried out under **bio-security**

ASF: the virus and the environment

Since the infection is not entirely transmitted through density dependent mechanism we have to shift to

The reduction of the environmental contamination of the virus

The problem then is not purely addressed in the mechanistic reduction of the wild boar density but in reducing the viral load of the environment

Standing Group of Experts on African swine fever in the Baltic and Eastern Europe region under the GF- TADs umbrella

SGE ASF3: Moscow, Russia, 15-16 March 2016

Wild boar population reduction should be considered, in combination with other control measures, within the framework of a wild boar management strategy **aimed at reducing ASF virus contamination of the environment.**

EU strategy

(see EFSA, 2015)

- ◆ Reduce the wild boar population size through targeted hunting of adult females
- ◆ Detection of – at least - 50% infected carcasses and their safe disposal
- ◆ Ban of winter/sustaining artificial feeding

Strategy applied - for at least - 100 km in front of the detected case

It is a medium term strategy that accepts the presence of the virus for a certain number of years

Thanks for the attention

Questions, comments?

BT SF



This presentation is delivered under contract with the Consumers, Health, Agriculture and Food Executive Agency (<http://ec.europa.eu/chafea>). The content of this presentation is the sole responsibility of Opera, the Istituto Zooprofilattico Sperimentale Lombardia e Emilia Romagna and the State Food and Veterinary Service of Latvia and it can in no way be taken to reflect the views of the Consumers, Health, Agriculture and Food Executive Agency or any other body of the European Union. The Consumers, Health, Agriculture and Food Executive Agency or any other body of the European Union will not be responsible under any circumstances for the contents of communication items prepared by the contractors.

OPERA

Viale Parioli 96 - 00197 Roma - Italy
Tel +39 06 96042652 / +39 06 8080111
Fax +39 06 89280678
info@opera-italy.it; www.btsftraining.com;
www.opera-italy.it

© Copyright holder: European Commission (2018)

Better Training for Safer Food BTSF

• *European Commission
Consumers, Health and Food Executive Agency
DRB A3/042
L-2920 Luxembourg*