

Avian Influenza H5N8 Situation in Austria

SCOPAFF

17./18. January 2017

Dr. Johann Damoser

HPAI poultry


- One case in a turkey farm in November 2016
- All restrictions already lifted (23.12.2016)



HPAI wild birds




Highly pathogenic avian influenza in wild birds
Positive samples



HPAI wild birds

Highly pathogenic avian influenza in wild birds Negative samples



HPAI wild birds

	negative	not analyzable	Ongoing investigation	positive	In total
No:	128	4	6	35	173

Concerned species:

curlew (*Brachvogel*), grey heron (*Graureiher*),
 great crested grebe (*Haubentaucher*), black-headed gull
 (*Lachmöwe*), common buzzard (*Mäusebussard*), tufted duck
 (*Reiherente*), mute swan (*Höckerschwan*), herring gull
 (*Silbermöve*), common gull (*Sturmmöwe*), pochard (*Tafelente*),
 water strider (*Waldwasserläufer*)

Risk areas - 1

1. Vorarlberg



2. Oberösterreich/Salzburg/Steiermark

Risk area - 2


- **Since 10 January 2017 the whole territory of Austria**
- According to Article 8 of the Austrian ordinance on HPAI, the following measures are applied within the risk area (in accordance with Directive 2005/94/EC):
 - All poultry has to be kept within a stable or at least in a closed equipment in order to avoid direct contact between wild birds and their faeces and poultry
 - No surface water for watering the poultry
 - Cleansing and disinfection with reasonable care
 - Notification of any decreased uptake of food and/or egg production
 - Notification of high mortality rate

Risk area - 2

- where indoor-keeping is not possible, (esp. in zoos), the CA can grant derogations by official notice
 - separation of poultry from water fowl (no direct or indirect contact)
 - Feeding only inside
 - Clinical investigation of those animals at least once a month
 - Serological investigation at the NRL every 3 months

Surveillance zone

- Due to an outbreak in the Slovak Republik



THANK YOU FOR YOUR ATTENTION