# Practical experiences: Use of local anaesthesia at castration Production of entire males

**SEGES Innovation** 






STØTTET AF

**Svine**afgiftsfonden



#### Must join course held by the veterinarian





- 2 hours theory
  - Mostly pictures and videos
  - No preparation or homework
- Practical training by the vet
  - Local anaesthesia minimum 10 pigs
- Lifelong certificate



We use both local anaesthesia and analgesia





#### **LOCAL ANAESTHESIA**





#### **ANALGESIA**













#### Inspiration for work routines using local anaesthesia

One litter at the time



- More litters
- Local anaesthesia before other routines
- Castration after other routines









#### Information at website in several languages

LOCAL ANAESTHESIA BEFORE CASTRATION

The pig must lie still







#### REMEMBER

- Castration bench
- Needle 25g (0.5 x 16 mm)
- Automatic injector or disposable syringe
- · Procamidor Vet. or Pronestesic
- Nitrile gloves
- Container for discarded needles and scalpels
- · Container for cut off testicles

## **LIST OF MATERIALS AND TOOLS**REMEMBER

- Castration bench
- Needle 25g (0.5 x 16 mm)
- Automatic injector or disposable syringe
- Procamidor Vet. or Pronestesic
- Nitrile gloves
- Container for discarded needles and scalpels
- Container for cut off testicles

#### **SELF INJECTION**

#### - NOW WHAT?


- Rinse with water in case of skin or eye contact
- Seek medical help if skin irritation occurs or if you self-inject
- Show the package leaflet to your GP







#### Information also made as short videos





#### Future - loose housing of lactating sows - focus piglet survival

- Today an important risk and frequent cause of piglet deaths
  - Hunger
  - Crushing/overlaying by the sow
- Very important and will become even more so when sows are loose
  - Need to ensure piglet access to udder
    - So piglets cannot be separated from sows for long
  - Important piglets recover quickly
    - Piglets must not be affected by anasthesia for long
      - · Therefore local anasthesia and analgesia
- Temporary confinement
  - We recommend sows to be confined until after castration to reduce risk of crushing
- Zero-confinement




#### **Production of male pigs – welfare**

- Male pigs do perform more aggression and sexual behaviours
  - Can cause scratches and leg-injuries
  - Mainly occurs in the final part of the growing period
  - Mounting behaviour can lead to leg injuries and increase in mortality
  - Important to be aware of indicators and reduce competition
  - Experience from three Danish herds
    - Split and mixed pens worked equally well



## Penile injuries!





## Penile injuries

Sex	Castrate	Male pigs	Im muno - <u>cast</u> r ate
No penile	218	197	196
No injuries (%)	87	89	92
Not developed (%)	88	3	2



Behaviour Aggression Pushing

Riding



#### Results productivity: castrates, male pigs and immunocastrates

DanBred DLY	Castrates	Male pigs	Immuno- castrates
g/day	1196 <sup>ab</sup>	1212a	1180b
FEsv/day	3.30a	2.92 <sup>b</sup>	3.07°
FEsv/kg	2.76a	2.41 <sup>b</sup>	2.61°



#### Results productivity: castrates, male pigs and immunocastrates

DanBred DLY	Castrates	Male pigs	Immuno- castrates
g/dag	1196 <sup>ab</sup>	1212a	1180 <sup>b</sup>
FEsv/dag	3.30a	2.92 <sup>b</sup>	3.07°
FEsv/kg	2.76a	2.41b	2.61°
Lean meat %	60.1a	63.5 <sup>b</sup>	62.6°
Slaughter loss	1.33 <sup>a</sup>	<u>1</u> .36 <sup>b</sup>	1.36 <sup>b</sup>



# Economy – basis DKK/pig

Insertion & carcass weight (30-89 kg)
Price reduction male pigs DKK 0.31/kg
Rejected DKK 4.6/kg (April 2021)
Feed price DKK/1.57/FEsv (5 year)
Pork price (5 year)

**Castration costs:** 

Analgesia DKK 0.72 Local anaesthesia DKK 2.93 Salary DKK 2.50 Lower mortality DKK 2.00 Total DKK 8.15

Improvac costs: Salary: DKK 1.50
Vaccine: DKK 24.00



#### **Economy**

	Castrates	Male pigs	Improvac
Castration DKK/pig	-8.15	-	-
FEsv/kg	0	-0.35	-0.15
Gain g/day	0	+16	-16
Lean meat%	0	+3.4	+2.5
Economy, DKK/pig	-	59.1	8.2
MAX rejected	-	14.4%	1.6%



#### Castration

- Local anaesthesia and analgesia MUST be used
  - Implement good work routines
- Do not castrate before the anaesthesia has had time to work - this also makes castration easier!
- Change scalpels and needles for each litter
- Use nitril gloves

## Producing male pigs

- Sorting boar semen does not work
- Sorting fathers (D boars) reduces boar taint but not enough!
- Immunocastration
  - Reduces boar taint markedly
    - But does not remove boar taint
  - Reduces aggressive behaviour
  - Economy is equal to producing castrates
  - BUT OUR MAIN MARKETS WILL NOT BUY THEM.

