

EFSA outputs on Avian Influenza

Frank Verdonck

PAFF meeting, 25 Oct 2017, Brussels





RISK ASSESSMENT ON

- AI introduction into the EU and into poultry holdings
- AI transmission and spread
- Mutation from LPAI to HPAI
- AI surveillance
- Biosecurity


EFSA scientific opinion

EFSA Journal 2017;15(10):4991 d oi:10.2903/j.efsa.2017.4991



AI INTRODUCTION

- Migratory water birds represent the most likely pathway of AIV introduction into the EU
- Mainly via the north eastern and eastern migratory routes




Clade	NE route	E route	S route	NW route
2.3.4.4	Benchmark	Slightly lower	Much lower	Much lower
2.2.1.2	Much lower	Much lower	Lower	Extremely low
2.3.2.1c	Similar	Similar	Lower	Extremely low



AI INTRODUCTION


 According to a mathematical model, AIV amplification and spread takes place when wild bird populations of sufficient size within the EU become infected





AI INTRODUCTION

The AIV prevalence in water birds as well as the size and composition of the wild bird reservoir are determining the probability of a holding to become infected






AI TRANSMISSION AND SPREAD

- The **transmission rate** between animals within a flock is assessed to be **higher for HPAI** viruses than LPAI viruses.
- Spread of HPAI viruses between farms is highly likely in the absence of control measures.
- In most cases, **LPAIV remain restricted** to a single farm, although horizontal spread has been observed in several occasions.



MUTATION FROM LPAI TO HPAI

No specific factors related to host species, environmental conditions or viral lineage were identified and likewise no molecular markers that would be useful predictors of increased risk of a specific LPAIV to mutate to an HPAI phenotype were recognized.





AI SURVEILLANCE OF POULTRY

- In gallinaceous poultry, passive surveillance through notification of <u>suspicious clinical signs/mortality</u> is the most effective method for early detection of HPAI outbreaks.
- For effective surveillance in **anseriform poultry passive surveillance** through notification of <u>suspicious clinical</u> <u>signs/mortality</u> needs to be accompagnied by <u>serological</u> <u>surveillance</u> and/or a <u>virological surveillance program</u> of birds found dead (bucket sampling).



AI SURVEILLANCE OF POULTRY

- The serological surveillance is unfit for early warning of LPAI outbreaks at the individual holding level. Serosurveillance could be effective in detecting clusters of LPAIV-infected holdings.
- Risk-based surveillance is useful as it targets flocks where AI introduction is considered to be higher, although there is limited quantitative (EUwide) evidence to weight the risk factors.




AI SURVEILLANCE OF WILD BIRDS

- Passive surveillance is an appropriate method for HPAI surveillance in wild birds if the HPAIV infections are associated with mortality.
- Active wild bird surveillance has a very low efficiency in detecting HPAI.



AI SURVEILLANCE OF WILD BIRDS

Targeted active wild bird surveillance combined with enhanced passive surveillance at a few priority regions in the EU may detect, if infection prevalence and sample sizes are sufficient, the presence of circulating AIV when these do not cause massive mortality among these birds.





BIOSECURITY

- The risk of AIV introduction and spread will remain high in production processes when movement of animals, restricting access throughout the whole production cycle and/or contact with wild birds is not reduced.
- If poultry cannot be confined during high-risk periods, it is recommended to prevent direct contact between wild birds and poultry by reducing the size of the outdoor area and/or by using netting. Feed and water should be provided under a roof or a horizontal fabric.
- Online biosecurity questionnaires could be used by farmers to check their current biosecurity level and subsequently to improve it based on the received feedback.



MONITORING AI SITUATION

- 2016-2017 outbreaks in the EU
- HPAI situation in other continents covering both human and animal health aspects!







European Union Reference Laboratory for Avian Influenza


collaboration between EU institutions and all affected MSs


EFSA scientific report

EFSA Journal 2017;15(10):5018 d oi:10.2903/j.efsa.2017. 5018



2016-2017 EPIDEMIC IN THE EU





■ The 2016/2017 HPAI epidemic was the **largest ever recorded in the EU** in terms of high number of outbreaks, wide geographic distribution and high number of dead wild birds.



2016-2017 EPIDEMIC IN THE EU

- Despite of a high number of human exposures to infected poultry during the ongoing outbreaks, no human cases have been identified in Europe.
- Challenges remain to identify all exposed people remain



CHARACTERISATION OF HPAI-AFFECTED HOLDINGS

- To further evolve from a descriptive towards a (quantitative) analytical analysis (risk factor analysis), there is a need to:
 - harmonise reporting amongst MSs (e.g. thresholds)
 - have poultry population data

Number of susceptible birds per affected holding	Commercial	Non-commercial
0-50	6/505 (1%)	318/455 (70%)
51-200	4/505 (1%)	102/455 (22%)
201-1,000	25/505 (5%)	22/455 (5%)
1,001-10,000	201/505 (40%)	12/455 (3%)
>10,000	269/505 (53%)	1/455 (0%)
Total	505	455



APPLIED PREVENTION AND CONTROL MEASURES

Case reports were submitted by 13 MSs to share information on applied prevention and control measures

Annex L — Applied prevention and control measures on avian influenza Italy

> Dorotea Tiziano, Mulatti Paolo, Bonfanti Lebana, Marangon Stefano Istituto Zooprofilatico Sperimentale delle Venezie

- Stringency and implementation of biosecurity is not clear across the EU.
- Communication among MSs is paramount in order to increase the level of preparedness, and to promptly apply control measures before the disease spread to non-affected MS, preventing and/or limiting the spread of the disease.



MONITORING AI IN OTHER CONTINENTS

- The current epidemiology of HPAIV H5N6 in Asia, with widespread occurrence in migratory birds of the order Anseriformes, and detection in apparently healthy northern pintails, indicates a risk of long-distance spread of this virus
- The HPAI situation in Africa of the subtypes H5N1 and H5N8 is evolving rapidly and requires close monitoring.
- Human infections due to transmission of H5N1, H5N6, H7N9 and H9N2 have only been observed in areas where these viruses circulate in the wild bird and/or poultry populations, mainly in Asia and Egypt.



ACKNOWLEDGEMENTS

- **WGs:** Andrew Breed, Adam Brouwer, Matthieu Guillemain, Timm Harder, Isabella Monne, Helen Roberts, Jan Arend Stegeman, Brown I, Mulatti P, Smietanka K, Staubach C, Willeberg P
- AHAW Panel: Simon More, Dominique Bicout, Anette Bøtner, Andrew Butterworth, Paolo Calistri, Klaus Depner, Sandra Edwards, Bruno Garin-Bastuji, Margaret Good, Christian Gortázar Schmidt, Virginie Michel, Miguel Angel Miranda, Søren Saxmose Nielsen, Mohan Raj, Liisa Sihvonen, Hans Spoolder, Hans-Hermann Thulke, Antonio Velarde, Preben Willeberg, Christoph Winckler,
- **Hearing experts:** Patrick Daniel, Teun Fabri, Nicolas Gaidet, Barbara Grabkowsky, Vittorio Guberti, Marie Guyot, Ursula Hoefle, Adeline Huneau Salaun, Thijs Kuiken, Hartmut Meyer, Virginie Michel, Ioana Neghirla, Éric Niqueux, Yali Si, Brigita Slavec, Krzysztof Smietanka, Christoph Staubach, Anna Luca Vescei, Josanne Verhagen, Sophie Von Dobschuetz
- **ECDC:** Adlhoch C, Kaja Kaasik Aaslav, Epidemic Intelligence team at ECDC and Pasi Penttinen
- **EFSA:** Francesca Baldinelli, Federica Barrucci, Chiara Fabris, Laura Martino, Olaf Mosbach-Schulz, Frank Verdonck, Joana Morgado, Zancanaro G
- AI consortium











ACKNOWLEDGEMENTS

Members States: Austria (Andrea Hoeflechner-Poeltl, Eveline Wodak), Belgium (Philippe Houdart, Katie Vermeersch), Bulgaria (Aleksandra Miteva, Anna Zdravkova), Croatia (Zlatko Krovina, Tihana Miškić) Cyprus (Giorgos Krasias), Czech Republic (Marie Vágnerová), Denmark (Thorkild Bastholm, Torben Grubbe), Estonia (Kärt Jaarma, Helen Prommik), Finland (Tiia Tuupanen), France (Loïc Evain, Isabelle Guerry, Alexandra Troyano-Groux), Germany (Franz Conraths, Andrea Coßmann), Greece (Sokratis Perdikaris), Hungary (Zsófia Szepesiné Kókány, Zsolt Terjék, Gabor Wyszoczky), Ireland (Stephanie Ronan, Eoin Ryan), Italy (Anna Sorgente), Latvia (Rudīte Vārna), Lithuania (Paulius Busauskas), Luxembourg (Roger Gindt), Malta (Joseph Caruana), the Netherlands (Nina Berendsen, Dennis Bol, Geert Eleveld, Marcel Spierenburg), Poland (Edyta Swieton), Portugal (Yolanda Vaz), Romania (Ioana Neghirla, Alexandru Supeanu, Claudiu Stroe), Slovakia (Vilem Kopriva, Barbora Pavlikova), Slovenia (Aleksandra Hari), Spain (Ana Fernandez Martín, Inés Moreno), Sweden (Annica Wallén Norell), United Kingdom (Adam Brouwer, Helen Roberts); Members States representatives that wrote the case reports on the AI secondary outbreaks and on the AI applied prevention and control measures, as reported in the Annex: Austria (Andrea Hoeflechner-Poeltl, Eveline Wodak), Belgium (Philippe Houdart, Bénédicte Lambrecht, Marjorie Piret, Mieke Steensels), Bulgaria (Aleksandra Miteva, Anna Zdravkova), Czech Republic (Milada Dubská, Petr Šatrán, Marie Vágnerová), Denmark (Pernille Dahl Nielsen, Stig Mellergaard), France (Mohamed Boukottaya, Anne Bronner, Alexandre Fediaevsky, Claire Guinat, Adeline Huneau-Salaün, Mathilde Paul), Greece (Sokratis Perdikaris), Hungary (Zsófia Szepesiné Kókány, Gerda Pállai, Anna Luca Vecsei, Gabor Wyszoczky), Ireland (Eoin Ryan), Italy (Tiziano Dorotea, Lebana Bonfanti, Stefano Marangon, Paolo Mulatti), the Netherlands (Marcel Spierenburg), Romania (Nicolae Drăgan, Ioana Neghirlă, Alexandru Supeanu, Claudiu Stroe), United Kingdom (Adam Brouwer, Ian Brown, Helen Roberts)