

Highly pathogenic avian influenza


Current Situation in Germany

SCoPAFF-meeting July 2023

bmel.de 💆 💿

Evolution in wild birds

Still high level of HPAI cases in wild birds, mainly affected species: colony breeders/seabirds








Situation in **poultry** and **captive birds**

Period of time	Number of outbreaks
Active outbreaks on 07/07/2023 (24/05/2023)	11 (26)
New from 24/05/2023 (~last PAFF) until 11/07/2023	2 Zoo: 0 Poultry: 2 NON-Poultry: 0
New season from 15/07/2022 until 11/07/2023	223 (212 lifted)



Evolution: **Poultry** and **captive birds**



New outbreak: Geese for fattening, measures are in place, epidemiological investigation is ongoing

Active outbreaks on 11/07/2023

DNK is part of the surveillance zone





Summary

- Number of cases in wild birds still high
- Infection pressure in the wild bird population is increasing because the colonies are now dissolving and moving further away from the colonies and the risk of spread at that time is high
- Number of HPAI outbreaks in **poultry** and **captive birds** nonetheless **low**
- Biosecurity measures in place
- Still essential: awareness



Thank you for your attention!

Contact:

Federal Ministry of Food and Agriculture Directorate 3 Division 323 – Animal Diseases - EU-Trade, International Affairs, Crises Centre 323@bmel.bund.de

Dr. Christiane Soltau 323@bmel.bund.de www.bmel.de



