

Corrigendum to the « Report on the monitoring and testing of ruminants for the presence of transmissible spongiform encephalopathy (TSE) in 2002

Table 46: Distribution of genotypes in TSE positive animals per Member State

	Known genotypes		Distribution of known genotypes					
	Number	% of TSE positives	NSP1	NSP2	NSP3		NSP4	NSP5
					ARQ/ARQ	Others		
Belgique/België	25	100%	0%	0%	8%	8%	12%	72%
Deutschland	0	0%	0%	0%	0%	0%	0%	0%
Ellas	57	84%	0%	2%	70%	23%	0%	5%
España	19	90%	0%	5%	84%	0%	0%	11%
France	61	25%	0%	7%	51%	2%	5%	36%
Ireland	23	35%	0%	0%	74%	17%	0%	9%
Italia	95	100%	0%	0%	85%	15%	0%	0%
Nederland	68	97%	0%	0%	7%	1%	3%	88%
United Kingdom	356	79%	0%	0%	24%	4%	5%	67%
Total	704	67%	0%	1%	39%	7%	4%	49%

Table 49: Relative susceptibility of genotypes to a TSE infection, based on monitoring results in France, Spain and the United Kingdom (NSP5=100%)

			NSP1	NSP2	NSP3		NSP4	NSP5
					ARQ/ARQ	others		
Frequency in random samples	FR	23%	39%	24%	3%	7%	5%	
	ES	6%	35%	49%	6%	1%	4%	
	UK	20%	42%	14%	12%	6%	6%	
Frequency in TSE cases	FR	0%	7%	51%	2%	5%	36%	
	ES	0%	5%	84%	0%	0%	11%	
	UK	0%	0%	24%	4%	5%	67%	
Relative susceptibility	FR	0%	3%	32%	7%	11%	100%	
	ES	0%	6%	66%	0%	0%	100%	
	UK	0%	0%	16%	3%	8%	100%	