



EUROPEAN COMMISSION  
HEALTH AND CONSUMERS DIRECTORATE-GENERAL

Director General

SANCO/10866/2013

*Programmes for the eradication, control and monitoring of certain  
animal diseases and zoonoses*

**The programme for  
the control of certain zoonotic salmonella in breeding,  
laying and broiler flocks of Gallus gallus and in flocks of  
turkeys (Meleagris gallopavo)**

**Romania**

**Approved\* for 2013 by Commission Decision 2012/761/EU**

\* in accordance with Council Decision 2009/470/EC

# Romanian National Control Programme for *Salmonella* in laying hens (*Gallus gallus*) 2013

## Part A

### a. Aim of the programme

To reduce or maintain the low prevalence of *Salmonellas* with public health significance in laying hens flocks (*Gallus gallus*) in Romanian holdings producing eggs intended for human consumption at least to the target levels set out in Regulation (EC) No 517/2011.

An annual minimum percentage of reduction of positive flocks of adult laying hens equal to at least:

- (i) 10 % if the prevalence in the preceding year was less than 10 %;
- (ii) 20 % if the prevalence in the preceding year was between 10 and 19 %;
- (iii) 30 % if the prevalence in the preceding year was between 20 and 39 %;
- (iv) 40 % if the prevalence in the preceding year was 40 % or more;

### b. Sampling programme

The National Control Programme for *Salmonella* in laying hens encompasses the following serovars of zoonotic *Salmonella*: *Salmonella enteritidis*, including the monophasic ST and *Salmonella typhimurium*.

The sampling programme will be in accordance to Regulation **2160/2003 EC** and Regulation **517/2011 EC**. We have also taken into account the Regulation 1177/2006 EC provisions.

The programme was elaborated in compliance with the requirements of Commission Regulation (EU) No 517/2011 of 25 May 2011 implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards a Union target for the reduction of the prevalence of certain *Salmonella* serotypes in laying hens of *Gallus gallus* and amending Regulation (EC) No 2160/2003 and Commission Regulation (EU) No 200/2010.

**c. Demonstrate the evidence that it complies with the specific requirements laid down in Parts D of Annex II to Regulation (EC) no 2160/2003**

### ***Specific requirements concerning flocks of laying hens***

1. Eggs shall not be used for direct human consumption as table eggs unless they originate from a commercial flock of laying hens subject to *Salmonella* national control programme established and is not under official restriction.

2. Eggs originating from flocks with unknown health status, that are suspected of being infected or that are infected with *Salmonella* serotypes for which a target for reduction has been set or which were identified as the source of infection in a specific human food-borne outbreak, may be used for human consumption only if they are treated in a manner that guarantees the destruction of all *Salmonella* serotypes with public health significance in accordance with Community legislation on food hygiene.

Eggs originating from flocks with unknown health status, that are suspected of being infected or that are infected with *Salmonella* serotypes for which a target for reduction has been set or which were identified as the source of infection in a specific human food-borne outbreak, shall be:

(a) considered as Class B eggs as defined in Article 2(4) of Commission Regulation (EC) No **557/2007** laying down detailed rules for implementing Council Regulation (EC) No **1028/2006** on marketing standards for eggs (1);

(b) marked with the indication referred to in Article 10 of Commission Regulation (EC) **No 557/2007** which clearly distinguishes them from Class A eggs prior to being placed on the market;

(c) prohibited access to packaging centres unless the competent authority is satisfied with the measures to prevent possible cross-contamination of eggs from other flocks.

3. When birds from infected flocks are slaughtered or destroyed, steps are taken to reduce the risk of spreading zoonoses as soon as possible. Slaughtering shall be carried out in accordance with Community legislation on food hygiene. Products derived from such birds may be placed on the market for human consumption in accordance with Community legislation on food hygiene. If they are not destined for human consumption, this products must be used or disposed of in accordance with Regulation (EC) **No 1774/2002**.

4. In order to exclude false-positive initial results, the competent authority may lift the restrictions laid down in point 2 of this Part:

(a) when the flock of layers is not the source of infection for humans by the consumption of eggs or egg products as a result of the epidemiological investigation of food-borne outbreaks in accordance with Article 8 of Directive **2003/99/EC**; and

(b) where the flock is subjected to a *Salmonella* national control programme and *Salmonella* serotypes which a target for reduction has been set, is not confirmed by the following sampling protocol carried out by the competent authority:

(i) the technical specifications referred to in Article 5 of Commission Decision **2004/665/EC** (seven samples); however, a sub-sample of 25 grams must be collected of each faecal material and dust sample for analysis; all samples must be analysed separately;

or

(ii) bacteriological investigation of the caeca and oviducts of 300 birds;

or

(iii) bacteriological investigation of the shell and the content of 4 000 eggs of each flock in pools of maxi-mum 40 eggs.

In addition to the sampling in point (b), the competent authority shall verify the absence of the use of antimicrobials, potentially affecting the result of the analyses of the sampling.

## 1. General

### 1.1 Summary referring to the occurrence of the salmonellosis in Romania

The programme for the control of Salmonella Enteritidis and Salmonella Typhimurium in laying hens flocks of Gallus gallus has been in operation in Romania since 2008. As a result, the number of Salmonella Enteritidis and Salmonella Typhimurium infected flocks of Gallus gallus in Romania is currently very low under 2%. During 2011, a totally of 411 laying hens flocks were tested for Salmonella and there were only 8 flocks positive for Salmonella Typhimurium and Salmonella Enteritidis . The prevalence for the target serotypes in broiler flock in 2011 was 1.9%, which is low and below the Community target.

The incidence of various serotypes of Salmonella in poultry farms, in 2011 in Romania

| NO. | SEROTYPE                                   | NO. OF STRAINS in Gallus gallus |
|-----|--|---------------------------------|
| 1.  | <b>S.Agona</b>                             | 7                               |
| 2.  | <b>S.Albany</b>                            | 8                               |
| 3.  | <b>S.Amsterdam</b>                         | 5                               |
| 4.  | <b>S.Enteritidis</b>                       | 114                             |
| 5.  | <b>S.Gallinarum</b>                        | 6                               |
| 6.  | <b>S.Gallinarum biovarianta Pullorum</b>   | 4                               |
| 7.  | <b>S.Gallinarum biovarianta Gallinarum</b> | 7                               |
| 8.  | <b>S.Glostrup</b>                          | 6                               |
| 9.  | <b>S.Hadar</b>                             | 15                              |
| 10. | <b>S.Havana</b>                            | 1                               |

|              |                      |            |
|--------------|----------------------|------------|
| 11.          | <b>S.Indiana</b>     | 2          |
| 12.          | <b>S.Infantis</b>    | 12         |
| 13.          | <b>S.Irumu</b>       | 3          |
| 14.          | <b>S.Kottbus</b>     | 1          |
| 15.          | <b>S.Livingstone</b> | 6          |
| 16.          | <b>S.Mbandaka</b>    | 17         |
| 17.          | <b>S.Molade</b>      | 1          |
| 18.          | <b>S.Montevideo</b>  | 7          |
| 19.          | <b>S.Namibia</b>     | 1          |
| 20.          | <b>S.Rissen</b>      | 1          |
| 21.          | <b>S.Senftenberg</b> | 9          |
| 22.          | <b>S.Taksony</b>     | 4          |
| 23.          | <b>S.Tennessee</b>   | 10         |
| 24.          | <b>S.Thompson</b>    | 48         |
| 25.          | <b>S.Typhimurium</b> | 1          |
| 26.          | <b>S.Uganda</b>      | 2          |
| 27.          | <b>S.Vilvoorde</b>   | 1          |
| <b>TOTAL</b> |                      | <b>303</b> |

## 1.2 The structure and organization of the relevant Competent Authorities

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in laying hens (*Gallus gallus*) flocks is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**

The departments responsible for implementing the National Control Programme of Salmonella in laying hens are:

- for implementing at national level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary and Food Chain Safety General Directorate** ;
- in respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the verification of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**
- Concerning the slaughter of flocks, the **Hygiene and Veterinary Epidemiology Directorate** is responsible for public health protection,

through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.

- In respect of EC Regulation No 183/2005 on feed hygiene at the central level is **The Directorate for Technical Coordination of Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**
- at county level for implementing the Regulation (EC) No 2160/2003,882/2004 and 183/2005 there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the national reference laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

#### **Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România  
Tel: 0374.322.013 / 0374.322.000 Fax: 0214.113.394 E-mail: [office@idah.ro](mailto:office@idah.ro)

- **H.I.P.V.H.** (Hygiene Institute and Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED**

#### **Hygiene Institute and Veterinary Public Health**

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201  
Tel: 021.252.46.51 Fax: 021.252.00.61 E-mail: [iispv@iispv.ro](mailto:iispv@iispv.ro)

### **1.3 Approved laboratories where samples collected within the programme are analyzed**

**Diagnostics:** The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL is show below:

| <b>Nr.</b> | <b>NAME</b> | <b>ADRESS</b> | <b>TELEPHON/ E-MAIL</b> |
|------------|-------------|---------------|-------------------------|
|------------|-------------|---------------|-------------------------|

| <b>Crt.</b> |                                  |  |   |
|-------------|----------------------------------|--|---|
| <b>1</b>    | <b>I.D.S.A</b>                   | București, Str. Dr.Staicovici, nr.63, sector 5, Cod 050557         | 0374/322.013<br><a href="mailto:office@idah.ro">office@idah.ro</a>  |
| <b>2</b>    | <b>L.S.V.S.A. Alba</b>           | Alba Iulia, Str. Lalelelor nr.7A, Cod 510217, Jud. Alba            | 0258/835.915<br><a href="mailto:office-alba@ansvsa.ro">office-alba@ansvsa.ro</a>  |
| <b>3</b>    | <b>L.S.V.S.A. Bacău</b>          | Bacau, Str Bucovinei, Nr 21, Jud. Bacău                            | 0234/586.233, int.111<br><a href="mailto:office-bacau@ansvsa.ro">office-bacau@ansvsa.ro</a>   |
| <b>4</b>    | <b>L.S.V.S.A.Bistrița-Năsăud</b> | Bistrita, Str. Tarpiului 26, Cod 420062, Jud. Bistrița Năsăud      | 0263/206.027; 0263/224.974<br><a href="mailto:office-bistrita-nasaud@ansvsa.ro">office-bistrita-nasaud@ansvsa.ro</a>  |
| <b>5</b>    | <b>L.S.V.S.A. Botoșani</b>       | Botosani , Str Tudor Vladimirescu Nr.17, Jud. Botoșani             | 0231/512.766<br><a href="mailto:office-botosani@ansvsa.ro">office-botosani@ansvsa.ro</a>  |
| <b>6</b>    | <b>L.S.V.S.A. Brașov</b>         | Brasov, Str. Calea Feldioarei Nr. 20 A, Cod 500450, Jud.Brașov     | 0268/440.257<br><a href="mailto:office-brasov@ansvsa.ro">office-brasov@ansvsa.ro</a><br><a href="mailto:dsvbv@rdslink.ro">dsvbv@rdslink.ro</a>                    |
| <b>7</b>    | <b>L.S.V.S.A. Brăila</b>         | Brăila, str. Calea Galați, nr.344, Cod 810385, Jud. Brăila         | 0239/610.689<br><a href="mailto:dsv@braila.rdsnet.ro">dsv@braila.rdsnet.ro</a><br><a href="mailto:office-braila@ansvsa.ro">office-braila@ansvsa.ro</a>            |
| <b>8</b>    | <b>L.S.V.S.A. Buzău</b>          | Buzau, Str. Horticolei, nr. 58 bis, Jud. Buzau                     | 0238-725001, 0238-725002<br><a href="mailto:office-buzau@ansv.ro">office-buzau@ansv.ro</a>  |
| <b>9</b>    | <b>L.S.V.S.A. Călărași</b>       | Călărași, Str. Prelungirea Dobrogei , nr. 4, Jud.Călărași          | 0242/313.676,<br><a href="mailto:office-calarasi@ansvsa.ro">office-calarasi@ansvsa.ro</a> ;<br><a href="mailto:office_dsv@satline.ro">office_dsv@satline.ro</a> ; |
| <b>10</b>   | <b>L.S.V.S.A Constanța</b>       | Constanta, Sos. Mangaliei Nr. 78, Jud. Constanța                   | 0241-682-417,<br>email: <a href="mailto:dsvct@rdscet.ro">dsvct@rdscet.ro</a>  |
| <b>11</b>   | <b>L.S.V.S.A. Cluj</b>           | Cluj-Napoca, Piata Marasti Nr. 1, Jud. Cluj                        | 0264-445729, 0264-448177<br><a href="mailto:office-cluj@ansvsa.ro">office-cluj@ansvsa.ro</a><br><a href="mailto:dsvcj@rdslink.ro">dsvcj@rdslink.ro</a>            |
| <b>12</b>   | <b>L.S.V.S.A. Dâmbovița</b>      | Târgoviște, Str. I.C. Brătianu, nr. 35, Cod 130055, Jud. Dâmbovița | 0372/737818<br><a href="mailto:office-dambovita@ansvsa.ro">office-dambovita@ansvsa.ro</a>   |
| <b>13</b>   | <b>L.S.V.S.A. Giurgiu</b>        | Giurgiu, Str. Bucuresti nr.72, Jud. Giurgiu                        | 0246/ 230.491<br><a href="mailto:office-giurgiu@ansvsa.ro">office-giurgiu@ansvsa.ro</a>   |
| <b>14</b>   | <b>L.S.V.S.A. Gorj</b>           | Târgul-Jiu, Str.Ecaterina Teodoroiu Nr.523, Jud. Gorj              | 0253/226.033<br><a href="mailto:dsv@intergorj.ro">dsv@intergorj.ro</a><br><a href="mailto:office-gorj@ansvsa.ro">office-gorj@ansvsa.ro</a>                        |
| <b>15</b>   | <b>L.S.V.S.A Hunedoara</b>       | Deva, Str. 22 Decembrie Nr. 226, Jud. Hunedoara                    | 0254-221145, 0254-230527<br><a href="mailto:office-hunedoara@ansvsa.ro">office-hunedoara@ansvsa.ro</a>  |
| <b>16</b>   | <b>L.S.V.S.A. Ialomița</b>       | Slobozia, str. Lacului, nr. 12 , Jud. Ialomița                     | 0243/ 232.069<br><a href="mailto:dsvsa-ialomita@ansvsa.ro">dsvsa-ialomita@ansvsa.ro</a>   |
| <b>17</b>   | <b>L.S.V.S.A. Iași</b>           | Iași, Aleea Mihail Sadoveanu 10, Jud. Iași                         | 0232-267501<br><a href="mailto:office-iasi@ansvsa.ro">office-iasi@ansvsa.ro</a>   |
| <b>18</b>   | <b>L.S.V.S.A. Maramureș</b>      | Baia Mare, Str. Vasile Alecsandri, Nr. 66, Jud. Maramureș          | 0262/ 224.031<br><a href="mailto:office-maramures@ansvsa.ro">office-maramures@ansvsa.ro</a>   |
| <b>19</b>   | <b>L.S.V.S.A. Mureș</b>          | Târgu Mures, str. Podeni nr.10 cod 540253, Jud. Mureș              | 0265/ 314.975<br><a href="mailto:office-mures@ansvsa.ro">office-mures@ansvsa.ro</a>   |
| <b>20</b>   | <b>L.S.V.S.A. Neamț</b>          | Piatra Neamț, Aleea Tiparului, nr. 12, cod 610263, Jud. Neamț      | 0233/ 223.259<br><a href="mailto:office-neamt@ansvsa.ro">office-neamt@ansvsa.ro</a>   |
| <b>21</b>   | <b>L.S.V.S.A. Prahova</b>        | Ploiesti, Str. Corlatesti, nr. 11, Jud. Prahova,                   | 0244 / 57.17.51<br><a href="mailto:office-prahova@ansvsa.ro">office-prahova@ansvsa.ro</a>   |
|             | <b>L.S.V.S.A. Satu Mare</b>      | Satu Mare, Str. Lăcrimioarei, Nr. 37, cod poștal                   | 0261/ 715.956   |

|    |                           |  |   |
|----|---------------------------|--|---|
| 22 |                           | 440067, Jud.Satu Mare  | <a href="mailto:office-satu-mare@ansvsa.ro">office-satu-mare@ansvsa.ro</a>                    |
| 23 | <b>L.S.V.S.A. Sibiu</b>   | Sibiu, str. Calea Şurii Mari, nr. 21, cod 550089, Jud. Sibiu | 0269/223.069<br><a href="mailto:office-sibiu@ansvsa.ro">office-sibiu@ansvsa.ro</a>            |
| 24 | <b>L.S.V.S.A. Suceava</b> | Suceava, Str. Scurta Nr. 2, Jud Suceava                      | Tel: 0230-522848,<br><a href="mailto:dsvsv@suceava.rdsnet.ro">dsvsv@suceava.rdsnet.ro</a>     |
| 25 | <b>L.S.V.S.A. Timiș</b>   | Timisoara, Str. Surorile Martir Caceu Nr.4, Jud. Timiș       | 0256-204911<br><a href="mailto:office-timis@ansv.ro">office-timis@ansv.ro</a>                 |
| 26 | <b>L.S.V.S.A. Vaslui</b>  | Bârlad, str. Trestiana, nr. 2, cod 731030, Jud.Vaslui        | 0235-421121, 0235-421413<br><a href="mailto:office-vaslui@ansv.ro">office-vaslui@ansv.ro</a>  |
| 27 | <b>L.S.V.S.A. Vrancea</b> | Focsani,B-dul Brailei, nr. 121 bis, cod 620122, Jud.Vrancea  | 0237-215561,0237-232727<br><a href="mailto:office-vrancea@ansv.ro">office-vrancea@ansv.ro</a> |

All Laboratories have to use the methods of the diagnostic presented at the point 1.4.

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the serotype according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

#### **1.4 Methods used in the examination of the samples in the framework of the programme**

Samples harvest by operators and samples harvest as official controls are prepared and tested in accordance with the requirements of the Commission Regulation (EC) No 517/2011, using the method recommended by the Community Reference Laboratory for *Salmonella* in Bilthoven, Netherlands. The method is described in the current version of ISO 6579:/A1:2007): 'Detection of *Salmonella* spp. in animal faeces and in samples of the primary production stage'. A semi-solid medium (modified semi-solid Rappaport-Vassiladis medium, MSRV) is used as the single selective enrichment medium. At least one isolate from each positive sample shall be serotyped according to the Kaufmann-White scheme. In general the sensitivity of a panel of 10 antimicrobials will be determined.

#### **1.5 Official controls**

One sample will be taken under the control of the Competent Authority for Regulation 2160/2003 from one layer flock on each holding with more than 1000 birds during the period of production of eggs for human consumption as specified in 2.1 of Annex to Commission Regulation (EC) No 517/2011.

Official samples will include a sample of dust (or when not available an additional sample of faecal material) in compliance with 2.1 and 2.2 of Annex to Commission Regulation (EC) No 517/2011.



The use of antimicrobials (as defined in Regulation (EC) No 1177/2006) will be checked when the official sample is taken. If the flock is under antimicrobial medication for animal health or animal welfare reasons, the flock will be sampled again after the withdrawal period of the product specified in the Marketing Authorization. Flock owners are required to keep records of antimicrobial used and to make these records available to the competent authorities.

The records of samples harvest by the operator will be made available for inspection to the Competent Authority or its agent and will provide details of the identity of the flock sampled, date of sample, slaughter date, type of sample, laboratory carrying out the examination, and the result. The number of flocks on the holding and the number of birds present will be recorded.

Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs requires poultry abattoirs to undertake microbiological testing for *Salmonella* on 5 samples a week (each sample is 3 neck skins). Establishments producing minced meat, meat preparations and mechanically separated meat must also undertake weekly testing for *Salmonella*.

### **Official controls at other stages of the food chain.**

Under the terms of the EC Feed Hygiene **Regulation 183/2005** provisions, feed businesses operators must be approved or registered by the Local Authority. Approvals/registrations are issued for the producers of compound feeds, feed materials, feed additives and premixtures. Approval requires a prior-inspection visit by the Local Authority to ensure that the establishments are in conformity with the required standards. The registration is followed by placing of premises on the list of feed business operators. The competent authority performs checks according to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order.

### **1.6 Measures taken by the Competent Authorities with regard to animals or products in which the presence of *Salmonella* spp. have been detected**

When a layer flock of *Gallus gallus* is suspected of being infected with *Salmonella* Enteritidis or *Salmonella* Typhimurium the flock will be investigated. The flock is suspected of being infected when *S. Enteritidis* or *S. Typhimurium* is isolated from a sample of faeces, boot swabs, tissue or organs from the birds in the flock, or from dust in the environment. This applies to layer flocks from day old through to end of production.

The Competent Authority will confirm the result of the sample by taking repeat samples in the flock consisting of two pairs of boot swabs per house (or equivalent faeces samples) plus one dust sample of at least 100 g (or equivalent faeces or boot swabs if dust is not available) as defined in **Commission Regulation (EC) No 517/2011 in Annex 2.1**. For the purposes of establishing the progress towards the target if *S. Enteritidis* or *S. Typhimurium* is isolated from any such sample the flock is classed as positive.

Isolates of *Salmonella Enteritidis* and *Salmonella Typhimurium* will be examined to determine if they are vaccine strains according to the manufacturer's protocol. If vaccine strains are confirmed in samples the flock will not be classed as positive for the purposes of establishing the progress towards the target.

Official sampling will be carried out under the control of the Competent Authority in all replacement flocks where the previous flock in a building was positive for *S. Enteritidis* or *S. Typhimurium*. This sampling will take place according to **Regulation (EC) No 517/2011 Annex 2.1b**.

Official samples will also be taken under the control of the Competent Authority when there is an epidemiological link to a case of human disease in a foodborne outbreak in accordance with Article 8 of **Directive (EC) No 2003/99**.

- Official samples will be taken under the control of the Competent Authority in all other flocks on a holding when *S. Enteritidis* or *S. Typhimurium* is suspected in a flock on the same holding.
- Operators will be invited to contact their veterinary for advice on how to reduce or eliminate the *Salmonella*. Advice will include recommendations on management, cleaning and disinfection, pest control, biosecurity, monitoring, and the potential use of vaccines.

**Arrangements for implementing the specific requirements concerning flocks of laying hens set out in Regulation 2160/2003 Annex II D.**

1) Starting **from 01 January 2010** when a layer flock of *Gallus gallus* is suspected of being infected with *Salmonella Enteritidis* or *Salmonella Typhimurium* the flock is placed under official control by the Competent Authority. If the flock is in the laying phase no further eggs from the flock may be placed on the market for human consumption unless they have been heat treated in such a way as to eliminate *Salmonella*.

2) The Competent Authority will confirm the result of the sample by taking repeat samples in the flock consisting of two pairs of boot swabs per house (or equivalent faeces samples) plus one dust sample of at least 100 g or equivalent faeces or boot swabs if dust is not available). All other flocks on the premises will be investigated in the same way if *S. Enteritidis* or *S. Typhimurium* is detected in the official sample of the flock suspected to be infected with *Salmonella*.

3) If no *Salmonella Enteritidis* or *Salmonella Typhimurium* are isolated from the repeat official samples (see 2) above), the official control measures are withdrawn.

4) If *S. Enteritidis* or *S. Typhimurium* is isolated from the official faeces/boot swab samples no further eggs may be placed directly on the market for human consumption. This restriction will remain in place for the duration of production of that flock. If *S. Enteritidis* or *S. Typhimurium* is isolated from official faeces or boot swabs in any other flock on the holding, eggs from the flock may not be placed directly on the market for human consumption, and this restriction will remain in place while the flock remains in production.

5) If *S. Enteritidis* or *S. Typhimurium* is isolated only from a dust sample, the flock will remain under official control. Eggs may continue to be sent direct for human consumption. Official samples of boot swabs (or equivalent faeces) and dust will be taken at 2 week intervals for analysis. Eggs may continue to be sent for human consumption if *S. Enteritidis* or *S. Typhimurium* are not isolated from the faeces samples or boot swabs.

6) A notice will be served requiring the owner to clean and disinfect the buildings where the infected flock was kept. The owner or person responsible for the flock is required to clean and disinfect the building where the infected birds were kept, and provide evidence to the Competent Authority that the cleaning and disinfection has been satisfactory by taking appropriate samples and having them analysed for the presence of *Salmonella*. Re-stocking may not take place until the cleaning and disinfection has been carried out and representative samples taken from the house after cleaning and disinfection have been shown to be negative for *S. Enteritidis* and *S. Typhimurium*).

7) When *S. Enteritidis* or *S. Typhimurium* is isolated from a sample taken from a flock before it comes into lay, the flock will be placed under official control. An official sample of the rearing flock suspected of being infected with *Salmonella* will be taken to confirm the infection.

The official sample taken will be as detailed in SAMPLING PROTOCOL and consist of either faeces or boot swabs. If *Salmonella Enteritidis* or *S. Typhimurium* is isolated from the official samples the flock may not be used to produce eggs direct for human consumption unless the eggs are treated in such a way as to eliminate *Salmonella*.

8) If the operator/owner of the laying flock disputes the results of the official test he/she may arrange to have samples taken of either

a) caecae and oviducts from 300 birds in the flock selected under supervision of the Competent Authority, or

b) 4000 eggs

9) and have them examined at his/her own expense at the national reference laboratory for the presence of *S. Enteritidis* or *S. Typhimurium*. Examination of eggs shall include both shell and contents. If *S. Enteritidis* or *S. Typhimurium* are not confirmed in these samples official controls on the flock will be removed and eggs may be placed on the market for direct human consumption. The Operator of the flock will be required to continue to comply with the sampling of the flock as detailed in Regulation (EC) No 517/2011 Annex 2.1. If one or more of the samples

taken from the birds is positive for *S. Enteritidis* or *S. Typhimurium* the flock will remain under official control and eggs may only be placed on the market for human consumption after treatment to eliminate *Salmonella*.

10) Antimicrobial treatment may not be used for the control of *Salmonella* in the national control programme except within the limits set by Commission Regulation (EC) No.1177/2006 on the use of specific control methods in the framework of the national programmes for the control of *Salmonella*. Vaccines to aid in the control of *Salmonella* may be used within the limits set by Commission Regulation (EC) No.1177/2006

11) Operators will be required to indicate at the time of sampling if the flock is being treated, or had received treatment during the last 2 week period, with antimicrobials for animal health or welfare reasons. If the flock is being treated with antimicrobials, or has been treated with antimicrobials during the last two week period for animal health or welfare reasons the flock will be re-sampled no sooner than at the end of the withdrawal period for the antimicrobial being used in the treatment.

### **1.7 Relevant national legislation.**

- N.S.V.F.S.A President **Order no. 34** to approve the sanitary veterinary norm regarding to monitoring zoonoses and zoonotic agents

- Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order

- N.S.V.F.S.A President **Order no. 205** for the approval of national reference laboratories and duties

- N.S.V.F.S.A President **Order no. 160** for the approval of the sanitary veterinary norm regarding the community reference laboratory for the zoonoses epidemiology and *Salmonella* and the national reference laboratory for *Salmonella*

### **1.8 Financial assistance provided to food and feed businesses operators in the context of the National Control Programme**

In laying hens flocks of *Gallus gallus* financial assistance is provided in the context of the control programme. This is set up by the **Government Decision no. 1214/2009** regarding the methodology for for determining and paying compensation to be paid to owners of slaughtered animals, killed or otherwise affected by the rapid liquidation of outbreaks of animal diseases.

The amount to compensate the animal owners in the context SNCP is determined in accordance with the provision of Government Decision No. 1214/2009 and is established by an evaluation committee consisting of:

- a) the representative of the county sanitary veterinary and food safety directorate (local CA)
- b) local representative of the Ministry of Agriculture and Rural Development with responsibilities in determining the genetic value of the animal
- c) the representative of the decision local unit from the local disease control center, designated by the Prefect,
- d) the local mayor or the person designated by him.

Convening of the evaluation committee is made by the Prefect at the written request of the county sanitary veterinary and food safety directorate (local CA).

On the proposal of the local county sanitary veterinary and food safety directorate the compensation committee considers and approves the replacement value of the animals according to the genetic value of animal, zootechnical value, sex, age, weight, physiological status, category production, at market price at the time when the liquidation of animals from the outbreak disease took place, and the average unit value based on the total amount of compensation for animals or products concerned according to Art. 4 of Regulation (EC) no. 349/2005 of the Commission, with the following amendments.

Romania applied to the Commission for co-financing for certain aspects of the control programme within the terms of Council Decision **2009/470/EC** on expenditure in the veterinary field.

## **2. Food and Feed businesses covered by the programme.**

The programme for the control of *Salmonella* in laying hens is part of the controls along the whole food chain.

### **2.1 The structure of the production of the given species and products thereof.**

The structure of the laying hens flocks and holding in Romania at the end of 2010 is presented below:

| <b>Region<br/>(County+NUTS<br/>CODE)</b> | <b>Type of flocks(b)</b> | <b>Total number of<br/>flocks ©</b> | <b>Total number of<br/>animals</b> | <b>Total number of<br/>holdings</b> |
|--|--------------------------|-------------------------------------|------------------------------------|-------------------------------------|
| <b>Total</b>                             | <b>Laying hens</b>       | <b>574</b>                          | <b>10936375</b>                    | <b>183</b>                          |

**In Romania in 2011 there were 183 holdings with 574 laying hens flocks in total.**

### **2.2 The structure of the production of food**

At the end of the laying flock production period the birds are slaughtered and may go for human consumption. In accordance with Regulation 853/2004 provisions, poultry meat intended for human consumption must be obtained in approved slaughterhouses. There are **40** approved slaughterhouses **in Romania**. Eggs sold at retail level within Romania are required to be marked with (stamped) with a code identifying the establishment (production site), country of origin and method of production (i.e organic, free range, barn or cage).

Poultry feed is supplied to farms by a small number of manufacturers. The major manufacturers of poultry feed operate to assurance schemes, apply HACCP principles and monitor for Salmonella.

### **2.3 Relevant guides for good animal husbandry practices or other guidelines**

Some of the relevant guides for good animal husbandry practices are presented at the below link:

[http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella\\_12404ro.pdf](http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella_12404ro.pdf)

[http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010\\_13247ro.pdf](http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010_13247ro.pdf)

[http://www.ansvsa.ro/documente/admin/community\\_guide\\_layers\\_hygiene\\_practice\\_pullet\\_egg\\_ro\\_17813ro.pdf](http://www.ansvsa.ro/documente/admin/community_guide_layers_hygiene_practice_pullet_egg_ro_17813ro.pdf)

Other relevant guides for good animal husbandry practices and other guidelines are under constructions.

All farm registered must have their individual plan for good farmer practices, which are subject to approval by the official veterinarian responsible for the control of the holding concerned.

### **2.4 Routine veterinary supervision of farms**

The owner is responsible for the health and welfare of the poultry on the holding, and for ensuring that a veterinarian is consulted on disease and welfare issues as appropriate. It is mandatory for each holding to have a contract with a private veterinarian who is responsible for veterinary care.

A veterinarian on behalf of the Competent Authority carries out inspections on farms for animal welfare reasons, to take samples for residues to administer and enforce the legislation on marketing of eggs and to check medicine records.

Also a veterinarian on behalf of the Competent Authority visit the farms and take official samples in the framework of Salmonella NCP according with the legislation in force.

It is mandatory for each county sanitary veterinary and food safety directorate (local CA) to report to the NSVFSA every month the number of samples and results of these tests for each flock. Also the Salmonella NRL has the

obligation to notify immediately NSVFSA and CSVFSD each positive sample for the relevant Salmonella.

## **2.5 Registration of farms**

All commercial poultry holdings in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centers / holdings of origin and means of transport for health and animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products.

All commercial holdings are entered into our national data base.

All poultry breeding flocks of more than 250 birds are included in the Salmonella National Control Programme.

## **2.6 Record-keeping at farms**

a. All laying hens flock operators are required to keep records of veterinary medicines use, including vaccines, which must be available for inspection.

b. Records relating to movement of flocks on to/off the holding must be kept.

c. Records giving details of sampling for *Salmonella* and results will be kept either at the holding or be readily available.

## **2.7 Documents to accompany animals when dispatched**

Operators wishing to export birds or hatching eggs to another EU Member State (or certain third countries) must comply with **EU Directive 2009/158/EC** and ensure that the consignment is accompanied by a completed and signed Intra-trade Animal Health Certificate (ITAHC) for poultry breeding and production. This must be completed and signed by the Official Veterinarian as well as the operator to confirm compliance with the relevant articles of Directive.

The ITAHC will also require the reference number of the operator's poultry health certificate. The ITAHC is amended to include the results of the last test for Salmonella as is required in the Commission Regulation (EC) 2160/2003 Art 9.1 prior to any dispatching of the live animals, or hatching eggs, from the food business operators place of origin. The date and the result of testing are included in the relevant health certificate provided in the Community legislation.

## **2.8 Other relevant measures to ensure the traceability of animals.**

It is require to the operators of poultry flocks to keep records of poultry or hatching eggs entering or leaving the premises. The records must contain

information on the number, date, and origin or destination. These records must be available to the Competent Authority for inspection.

All official veterinary health certificates issued for the intra community trade of poultry and hatching eggs are recorded on the Trade Control and Expert System (TRACES). This system allows tracking of exports of live animals and hatching eggs accompanied by veterinary health certification. TRACES generate ITAHCs issued for intra-Community movements. TRACES are an internet-based service which is owned and maintained by the Commission.

## **PART B**

### **1. Identification of the programme**

|                             |  |
|-----------------------------|--|
| Member State:               | <b>Romania</b>   |
| Disease:                    | <b>Salmonella Enteritidis and Typhimurium in Laying hens (GALLUS gallus)</b>   |
| Year of implementation:     | <b>2013</b>  |
| Reference of this document: | <b>National Sanitary Veterinary and Food Safety Authority<br/>No. /</b>  |
| Contact :                   | <b>Dr. Niculae LAZĂR,<br/>Director,<br/>phone: +4 021 315 78 75;<br/>fax: +4021 31249 67,<br/>e-mail: <a href="mailto:lazar.niculae@ansvsa.ro">lazar.niculae@ansvsa.ro</a></b> |



2. **Historical data on the epidemiological evolution of the disease**

*Salmonella* has been recognized as an important zoonotic pathogen for many years. *Salmonella Enteritidis* and *Salmonella Typhimurium* have accounted for the majority of cases of human salmonellosis and have consistently been the most commonly implicated pathogens in general outbreaks of food-borne disease.

A National Control Programme of *Salmonella* was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 517/2011. The national control programme for *Salmonella* in laying hens flock was coming into effect in January 2008.

In line with Regulation EC No 1003/2005 since 01 January 2007 the programme includes the control of *Salmonella Enteritidis*, *Salmonella Typhimurium*, *S. Hadar*, *S. Infantis* and *S. Virchow*. As a result of the control programme the number of *Salmonella Enteritidis* and *Salmonella Typhimurium* infected breeding flocks of *Gallus gallus* in Roumania is currently low. Of the other three *Salmonella* serovars, *Salmonella Hadar*, *Salmonella Infantis* and *Salmonella Virchow*, the occurrence is likewise at low levels. Breeding flocks which are confirmed to be infected with zoonotic *Salmonella* are compulsorily slaughtered.

The success of the control programme in breeding flocks means that the day old chicks of layers which are placed on farms should be free of *S. Enteritidis* and *S. Typhimurium*.

## The incidence of various serotypes of Salmonella in poultry, in 2010

| NO.          | SEROTYPE                                  | NO. OF STRAINS in Gallus gallus |
|--------------|---|---------------------------------|
| 1.           | <b>S.Agona</b>                            | 7                               |
| 2.           | <b>S.Albany</b>                           | 8                               |
| 3.           | <b>S.Amsterdam</b>                        | 5                               |
| 4.           | <b>S.Enteritidis</b>                      | 114                             |
| 5.           | <b>S.Gallinarum</b>                       | 6                               |
| 6.           | <b>S.Gallinarum biovariata Pullorum</b>   | 4                               |
| 7.           | <b>S.Gallinarum biovariata Gallinarum</b> | 7                               |
| 8.           | <b>S.Glostrup</b>                         | 6                               |
| 9.           | <b>S.Hadar</b>                            | 15                              |
| 10.          | <b>S.Havana</b>                           | 1                               |
| 11.          | <b>S.Indiana</b>                          | 2                               |
| 12.          | <b>S.Infantis</b>                         | 12                              |
| 13.          | <b>S.Irumu</b>                            | 3                               |
| 14.          | <b>S.Kottbus</b>                          | 1                               |
| 15.          | <b>S.Livingstone</b>                      | 6                               |
| 16.          | <b>S.Mbandaka</b>                         | 17                              |
| 17.          | <b>S.Molade</b>                           | 1                               |
| 18.          | <b>S.Montevideo</b>                       | 7                               |
| 19.          | <b>S.Namibia</b>                          | 1                               |
| 20.          | <b>S.Rissen</b>                           | 1                               |
| 21.          | <b>S.Senftenberg</b>                      | 9                               |
| 22.          | <b>S.Taksony</b>                          | 4                               |
| 23.          | <b>S.Tennessee</b>                        | 10                              |
| 24.          | <b>S.Thompson</b>                         | 48                              |
| 25.          | <b>S.Typhimurium</b>                      | 1                               |
| 26.          | <b>S.Uganda</b>                           | 2                               |
| 27.          | <b>S.Vilvoorde</b>                        | 1                               |
| <b>TOTAL</b> |   | <b>303</b>                      |

It is a statutory requirement for all laboratories which isolate *Salmonella* from a flock of chickens or its environment to report the

finding and supply the isolate to the National Reference Laboratory (NRL) for *Salmonella*. The isolates are serotyped, phage-typed, where appropriate, and tested for antimicrobial sensitivity by the NRL. This information is recorded and analyzed. The number of reports received depends on the level and sensitivity of monitoring which is undertaken by the producers. The reports provide useful

information on the serovars which are most common in poultry, and indicate trends.

### 3. Description of the submitted programme

#### **Objectives**

The target for Romania in 2009 can not be set as we did not participate to the baseline study regarding laying hens. The main objective of our programme for the reduction of Salmonella enteritidis and Salmonella typhimurium in adult laying hens of Gallus gallus shall be a reduction of the maximum percentage of positive adult laying flocks according to Regulation (EC) No 2160/2003 and Regulation (EC) 517/2011.

The subsidiary objectives of the programme include further reduction of the incidence of infected table eggs and eventually the reduction of the occurrence of human salmonellosis and a consequent reduction in suffering, mortality and health service costs. This programme is in accordance to Council Directive 1999/74/EC, Commission Decision 2008/425/EC, Council Decision 92/65/EEC, Council Decision 2009/470/EEC, Regulation 2160/2003/EC, Regulation 517/2011/EC and Regulation 1177/2006/EC.

#### **Target animal population**

The National Control Programme for Salmonella in laying flocks will be held in all holdings of laying hens consisting of at least 350 poultry of Gallus gallus which produce eggs for human consumption. Laying hens holdings which have between 350 and 1000 of birds will not be the subject of official testing, but will perform tests on the initiative of operators (self-control).

Small flocks that are reared to supply eggs for private domestic use, or small quantities of primary product supplied directly by the producer to the final consumer, will be exempt, as permitted in Regulation (EC) No 2160/2003 Article 1.3.

The number of holdings and flocks in laying hens sector at the end of 2011 the are shown **in the table below**.

**Table . Number of holdings and flocks in the laying hens sector**

| Region<br>(County+NUTS<br>CODE) | Type of flocks(b) | Total number of<br>flocks © | Total number of<br>animals |
|---------------------------------|-------------------|-----------------------------|----------------------------|
| RO121-AB                        | laying hens       | 9                           | 145104                     |
| RO421-AR                        | laying hens       | 14                          | 102000                     |
| RO311-AG                        | laying hens       | 12                          | 298944                     |
| RO211-BC                        | laying hens       | 3                           | 16500                      |
| RO111-BH                        | laying hens       | 7                           | 49575                      |

|              |                    |            |                 |
|--------------|--------------------|------------|-----------------|
| RO112-BN     | laying hens        | 4          | 65539           |
| RO212-BT     | laying hens        | 1          | 18240           |
| RO221-BR     | laying hens        | 34         | 996765          |
| RO122-BV     | laying hens        | 10         | 158736          |
| RO321-BUC    | laying hens        | 0          | 0               |
| RO222-BZ     | laying hens        | 0          | 0               |
| RO312-CL     | laying hens        | 5          | 55082           |
| RO422-CS     | laying hens        | 7          | 329404          |
| RO113-CJ     | laying hens        | 25         | 125300          |
| RO223-CT     | laying hens        | 14         | 422480          |
| RO123-CV     | laying hens        | 0          | 0               |
| RO313-DB     | laying hens        | 38         | 1000000         |
| RO411-DJ     | laying hens        | 4          | 11862           |
| RO224-GL     | laying hens        | 19         | 783000          |
| RO314-GR     | laying hens        | 83         | 1411578         |
| RO412-GJ     | laying hens        | 2          | 135456          |
| RO124-HR     | laying hens        | 4          | 106000          |
| RO423-HD     | laying hens        | 18         | 181050          |
| RO315-IL     | laying hens        | 16         | 422308          |
| RO213-IS     | laying hens        | 3          | 33000           |
| RO322-IF     | laying hens        | 0          | 0               |
| RO114-MM     | laying hens        | 26         | 222610          |
| RO413-MH     | laying hens        | 0          | 0               |
| RO125-MS     | laying hens        | 71         | 1500000         |
| RO214-NT     | laying hens        | 15         | 186849          |
| RO414-OT     | laying hens        | 11         | 237540          |
| RO316-PH     | laying hens        | 0          | 0               |
| RO116-SJ     | laying hens        | 10         | 133900          |
| RO115-SM     | laying hens        | 18         | 286000          |
| RO126-SB     | laying hens        | 14         | 118934          |
| RO215-SV     | laying hens        | 3          | 3600            |
| RO317-TR     | laying hens        | 16         | 428552          |
| RO424-TM     | laying hens        | 19         | 200939          |
| RO225-TL     | laying hens        | 0          | 0               |
| RO415-VL     | laying hens        | 0          | 0               |
| RO226-VN     | laying hens        | 3          | 64178           |
| RO216-VS     | laying hens        | 36         | 685350          |
| IDAH         | laying hens        | 0          | 0               |
| <b>Total</b> | <b>laying hens</b> | <b>574</b> | <b>10936375</b> |

### Sampling programmes

The National Salmonella Control Programme encompasses the following serovars of zoonotic Salmonella: Salmonella enteritidis and Salmonella typhimurium .

The sampling programme will be in accordance to Regulation 2160/2003 EC and Regulation 517/2011 EC. We have also taken into account Regulation 1177/2006 EC.

#### **4.1. Summary of measures under the programme year: 2013**

- Control
- Testing
- Slaughter of positive animals
- Killing of positive animals
- Vaccination
- Treatment of animal products
- Disposal of products
- Monitoring or surveillance
- Other measures

#### **4.2. Designation of the central authority charged with supervising and coordinating the departments responsible for implementing the programme:**

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in laying hens flocks of *Gallus gallus* is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**



The departments responsible for implementing the National Control Programme of Salmonella in laying hens are:

- for implementing at central level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary General Directorate** ;
- In respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the checking of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**.
- Concerning the slaughter of flocks, the **Hygiene and Veterinary Epidemiology Directorate** is responsible for public health protection, through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.
- In respect of EC Regulation No **183/2005** on feed hygiene at the central level is **The Directorate for Technical Coordination of**

**Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**

- At county level for implementing the Regulation (EC) No **2160/2003**, **882/2004** and **183/2005** there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the National Reference Laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

**NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

**Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România  
Tel: 0374.322.013 / 0374.322.000 Fax: 0214.113.394 E-mail: [office@idah.ro](mailto:office@idah.ro)

- **H.I.P.V.H.** (Hygiene Institute for Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

**NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED**

**Hygiene Institute for Sanitary Veterinary Public Health**

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201  
Tel: 021.252.46.51 Fax: 021.252.00.61 E-mail: [iispv@iispv.ro](mailto:iispv@iispv.ro)

The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL at February 2010 is show in the link below:

[http://www.ansvsa.ro/documente/admin/lista%20laboratoarelor%202011%20-%2028.01.2011\\_11043ro.pdf](http://www.ansvsa.ro/documente/admin/lista%20laboratoarelor%202011%20-%2028.01.2011_11043ro.pdf)

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the serotype according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

### **1. N.S.V.F.S.A.**

- proposal of the plan of disease control.
- elaborate and submit to the EC the National Control Programme
- evaluate, coordinate the implementation of the National Control Programme in Romania and propose the update of this programme if is necessary (depending of the national situation)
- adoption of measures based on the disease situation in Romania
- submission of reports to the E.C.
- training the specialists from C.S.V.F.S.D

### **2. S.V.F.S.D.**(Sanitary Veterinary and Food Safety Direction).

- Coordination of the programme at the county level
- Official sampling is performed by the official veterinarian
- Propose the compensation costs to the Commission which will evaluate compensation documents.

**3. I.D.A.H.** Institute of Diagnosis and Animal Health is the National Reference Laboratory concerning animal health and was also designated as NRL for Salmonella in animals.

Responsibilities and tasks of the National Reference Laboratory for Salmonella (**I.D.A.H.**), pursuant to Directive 2003/99/EC and Regulation (EC) No 2160/2003, according to provision of N.S.V.F.S.A.. President Order no.160/2006 and 205/2007 are:

#### **1. General duties**

- (a) To collaborate with the Community reference laboratory in their area of competence.
- (b) To coordinate, as appropriate, the activities of laboratories responsible for the analysis of samples in accordance with, in particular, Articles 4, 5 and 7 of Directive 2003/99/EC.
- (c) To coordinate the activities of laboratories responsible for the analysis of samples in accordance with Article 12(1) of Regulation (EC) No 2160/2003/EC.
- (d) Where appropriate, to organise comparative tests between the laboratories referred to under (b) and (c) and to assure an appropriate follow-up of such comparative testing.

- (e) To ensure the dissemination to the competent authority and to the laboratories referred to under (b) and (c), of the information that the Community reference laboratory supplies.
- (f) To provide scientific and technical assistance to the National Sanitary Veterinary and Food Safety Authority in their area of competence.
- (g) Characterize the pathogen isolates, genetic typing of this agents
- (h) Keep in maxim security conditions, the isolates
- (i) Give to the C.E., CRL, OMS, and national reference Laboratory of other member states, with the accord of N.S.V.F.S.A. all the information required.
- (j) train of the specialists from the government and private laboratory.

## **2. Specific functions and duties**

- (a) To participate, as appropriate in the monitoring and control programme for Salmonella and related anti-microbial resistance pursuant to Directive 2003/99/EC and in the analysis and testing of *Salmonella* pursuant to Regulation (EC) No 2160/2003.
- (b) To inform, as appropriate, the Community reference laboratory on aspects related to Salmonella vaccine strains and other specific control methods.
- (d) To gather data and information on the activities developed and methods used in relevant laboratories and to inform the Community reference laboratory thereof.
- (e) To monitor the epidemiological evolution of salmonella in Roumania.

## **5. S.V.F.S.L.**

There are 41 county Sanitary Veterinary and Food Safety Laboratories. Only 27 of them apply quality assurance systems that conform to the requirements of the current EN/ISO standard and are designated by National Reference Laboratory for Salmonella in animals to perform bacteriological examinations in the framework of the programme under the supervision of IDAH-NRL for Salmonella.

## **6. H.I.P.S.V.H.**

Hygiene Institute for Public Sanitary Veterinary Health is the National Reference Laboratory concerning the expertise for food products of animal origin and feed and it is also designed as the National Reference Laboratory for Salmonella in food and feed –public veterinary health.



### **4.3. Description and delimitation of the geographical and administrative areas in which the programme is to be implemented:**

The National Control Programme will be implemented throughout Romania, covering all the national territory and will cover all laying hens flocks of *Gallus gallus* consisting of at least 350 poultry which produce eggs for human consumption .

The administrative boundaries are the boundaries of the country. Roumania is administrative divided in 42 counties. There are 42 County Sanitary Veterinary and Food Safety Directorates and 41 County Sanitary veterinary and food Safety Laboratories.

### **4.4 Measures implemented under the programme**

#### **Bio-Security Measures**

Bio-security is a combination of practices, which are intended to prevent the spread of disease-causing organisms within the poultry farm. Where these are performed in parallel with the sanitation and disinfection procedures, bio-security measures could eradicate or, at least, reduce the level of pathogens to values, at which no hazard of infection would be likely.

The bio-security measures in commercial poultry farms are in accordance to the N.S.V.F.S.A. President Order 147/2006 regarding the bio-security measures in commercial poultry farms and the movements of live poultry, products and poultry by-products.

#### **Bio-security measures on holdings:**

- Health status of poultry
- On entering to all houses on the farm must be located disinfection barrier
- Control of movement of people
- Transport hygiene
- Feed hygiene
- Water hygiene
- Rodent, insect and bird control
- Cleaning and disinfecting of buildings
- Recording of all events and operations

According to the provision of Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program approved every year by N.S.V.F.S.A. President Order, after each cleaning and disinfections the owner is oblige to take samples to verify the efficient of the disinfection. The official

veterinarian take sample to verify the efficiency of the disinfection only in case of a positive flock.

Hygiene measures on poultry farms are also assessed during visits for the collection of official samples and during general visits to premises for other purpose.

#### **4.4.1. Measures and terms of legislation as regards the registration of holdings:**

All commercial poultry flocks in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centers / holdings of origin and means of transport for health and animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products. Poultry holdings shall be registered and sanitary veterinary approved in order to be able to operate.

The legal framework for carrying out commercial activity with poultry and poultry products consists of:

- **Order** of the President of the National Sanitary Veterinary and Food Safety Authority no **144/2006** for the approval of the Sanitary veterinary norm on animal health conditions governing intra-Community trade in, and imports from third countries of, poultry and hatching eggs transposing Council Directive 90/539/EEC on animal health conditions governing intra-Community trade in, and imports from third countries of poultry and hatching eggs.

In order to be sanitary veterinary approved, poultry holdings shall satisfy, in accordance with the legislation in force, the following conditions:

- Appropriate facilities and operation;
- Application of the “Program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” approved by Order of the National Sanitary Veterinary and Food Safety Authority President .
- At least one inspection visit per year by the official veterinarian;
- Additional checks to verify the compliance of the establishment with the hygiene measures and the operation of the establishments.

Each poultry holding receives a distinct approval number, number that can be the same with the one given in compliance with the **Council Regulation no. 2782/75/EEC** on the production and marketing of eggs for hatching and of farmyard poultry chicks.

#### **4.4.2. Measures and terms of legislation as regards the identification of animals:** Not applicable.

#### 4.4.3. Measures and terms of legislation as regards the notification of the disease:

**Commission Decision 176/2005** of 1 March 2005 laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2006/924/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2010/160/EU** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2008/755/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2009/847/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

The internal notification is made under the provision of **N.S.V.F.S.A. President ORDER no. 79 /2008** to approve the sanitary and veterinary norm on domestic notification and official declaration of animal diseases.

#### 4.4.4. Measures and terms of legislation as regards the measures in case of a positive result:

Whenever a flock is found positive by own-check sampling in the framework of the SNCP in breeding hens this flock is considered a suspect flock and movement restriction and other relevant restrictive measures are mandatory imposed on this flock.

When a layer flock of *Gallus gallus* is suspected of being infected with *Salmonella* Enteritidis or *Salmonella* Typhimurium the flock will be investigated. The flock is suspected of being infected when *S. Enteritidis* or *S. Typhimurium* is isolated from a sample of faeces, boot swabs, tissue or organs from the birds in the flock, or from dust in the environment. This applies to layer flocks from day old through to end of production.

The Competent Authority will confirm the result of the sample by taking repeat samples in the flock consisting of two pairs of boot swabs per house (or equivalent faeces samples) plus one dust sample of at least 100 g (or equivalent faeces or boot swabs if dust is not available) as defined in **Commission Regulation (EC) No 517/2011 in Annex 2.1**. For the purposes of establishing the progress towards the target if *S. Enteritidis* or *S. Typhimurium* is isolated from any such sample the flock is classed as positive.

Isolates of *Salmonella Enteritidis* and *Salmonella Typhimurium* will be examined to determine if they are vaccine strains according to the manufacturer's

protocol. If vaccine strains are confirmed in samples the flock will not be classed as positive for the purposes of establishing the progress towards the target.

Official sampling will be carried out under the control of the Competent Authority in all replacement flocks where the previous flock in a building was positive for *S. Enteritidis* or *S. Typhimurium*. This sampling will take place according to **Regulation (EC) No 517/2011 Annex 2.1b**.

Official samples will also be taken under the control of the Competent Authority when there is an epidemiological link to a case of human disease in a foodborne outbreak in accordance with Article 8 of **Directive (EC) No 2003/99**.

- Official samples will be taken under the control of the Competent Authority in all other flocks on a holding when *S. Enteritidis* or *S. Typhimurium* is suspected in a flock on the same holding.
- Operators will be invited to contact their veterinary for advice on how to reduce or eliminate the *Salmonella*. Advice will include recommendations on management, cleaning and disinfection, pest control, biosecurity, monitoring, and the potential use of vaccines.

### **Arrangements for implementing the specific requirements concerning flocks of laying hens set out in Regulation 2160/2003 Annex II D.**

1) Starting from **01 January 2010** when a layer flock of *Gallus gallus* is suspected of being infected with *Salmonella Enteritidis* or *Salmonella Typhimurium* the flock is placed under official control by the Competent Authority. If the flock is in the laying phase no further eggs from the flock may be placed on the market for human consumption unless they have been heat treated in such a way as to eliminate *Salmonella*.

2) The Competent Authority will confirm the result of the sample by taking repeat samples in the flock consisting of two pairs of boot swabs per house (or equivalent faeces samples) plus one dust sample of at least 100 g or equivalent faeces or boot swabs if dust is not available). All other flocks on the premises will be investigated in the same way if *S. Enteritidis* or *S. Typhimurium* is detected in the official sample of the flock suspected to be infected with *Salmonella*.

3) If no *Salmonella Enteritidis* or *Salmonella Typhimurium* are isolated from the repeat official samples (see 2) above), the official control measures are withdrawn.

4) If *S. Enteritidis* or *S. Typhimurium* is isolated from the official faeces/boot swab samples no further eggs may be placed directly on the market for human consumption. This restriction will remain in place for the duration of production of that flock. If *S. Enteritidis* or *S. Typhimurium* is isolated from official faeces or boot swabs in any other flock on the holding, eggs from the flock may not be placed directly on the market for human consumption, and this restriction will remain in place while the flock remains in production.

5) If *S. Enteritidis* or *S. Typhimurium* is isolated only from a dust sample, the flock will remain under official control. Eggs may continue to be sent direct for human consumption. Official samples of boot swabs (or equivalent faeces) and dust will be taken at 2 week intervals for analysis. Eggs may continue to be sent for human consumption if *S. Enteritidis* or *S. Typhimurium* are not isolated from the faeces samples or boot swabs.

Where the presence of *Salmonella* *Enteritidis* and *Salmonella* *Typhimurium* is not detected but antimicrobials or bacterial growth inhibitory effects are detected it shall be accounted for as an infected laying flock for the purpose of the Union target.

6) A notice will be served requiring the owner to clean and disinfect the buildings where the infected flock was kept. The owner or person responsible for the flock is required to clean and disinfect the building where the infected birds were kept, and provide evidence to the Competent Authority that the cleaning and disinfection has been satisfactory by taking appropriate samples and having them analysed for the presence of *Salmonella*. Re-stocking may not take place until the cleaning and disinfection has been carried out and representative samples taken from the house after cleaning and disinfection have been shown to be negative for *S. Enteritidis* and *S. Typhimurium*).

7) When *S. Enteritidis* or *S. Typhimurium* is isolated from a sample taken from a flock before it comes into lay, the flock will be placed under official control. An official sample of the rearing flock suspected of being infected with *Salmonella* will be taken to confirm the infection.

The official sample taken will be as detailed in SAMPLING PROTOCOL and consist of either faeces or boot swabs. If *Salmonella* *Enteritidis* or *S. Typhimurium* is isolated from the official samples the flock may not be used to produce eggs direct for human consumption unless the eggs are treated in such a way as to eliminate *Salmonella*.

8) If the operator/owner of the laying flock disputes the results of the official test he/she may arrange to have samples taken of either

a) caecae and oviducts from 300 birds in the flock selected under supervision of the Competent Authority, or

b) 4000 eggs

9) and have them examined at his/her own expense at the national reference laboratory for the presence of *S. Enteritidis* or *S. Typhimurium*. Examination of eggs shall include both shell and contents. If *S. Enteritidis* or *S. Typhimurium* are not confirmed in these samples official controls on the flock will be removed and eggs may be placed on the market for direct human consumption. The Operator of the flock will be required to continue to comply with the sampling of the flock as detailed in Regulation (EC) No 517/2011 Annex 2.1. If one or more of the samples taken from the birds is positive for *S. Enteritidis* or *S. Typhimurium* the flock will remain under official control and eggs may only be placed on the market for human consumption after treatment to eliminate *Salmonella*.

10) Antimicrobial treatment may not be used for the control of Salmonella in the national control programme except within the limits set by Commission Regulation (EC) No.1177/2006 on the use of specific control methods in the framework of the national programmes for the control of Salmonella. Vaccines to aid in the control of Salmonella may be used within the limits set by Commission Regulation (EC) No.1177/2006

11) Operators will be required to indicate at the time of sampling if the flock is being treated, or had received treatment during the last 2 week period, with antimicrobials for animal health or welfare reasons. If the flock is being treated with antimicrobials, or has been treated with antimicrobials during the last two week period for animal health or welfare reasons the flock will be re-sampled no sooner than at the end of the withdrawal period for the antimicrobial being used in the treatment.

In case of suspicion or conformation of **Salmonella enteritidis** or **Salmonella typhimurium** the NRL shall notify immediately the N.S.V.F.S. and local C.S.V.F.S.D..

### **Control of the use of feed antibiotics by official sampling**

According to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order, feeding stuffs intended for poultry nutrition are checked in view to avoid the contamination with Salmonella spp. Also, in conformity with the same legislation the feed stuffs are checked in view to detect the use of antibiotics.

Residues examination is performed according to the Romanian annual plan for examination for residues in live animals and animal origin products .

For broiler, hens, turkeys, other poultry a sample consists on one or more animals depending on the requirements of the analytical methods.

For each category of poultry considered, the minimum number of samples to be taken each year must be at least equal to one per 200 tones of annual production, with a minimum of 100 samples for each group of substances if the annual production of the category of birds considered is over 5 000 tones.

The following breakdown must be respected:

**Group A:** 50 % of the total samples. The equivalent of one fifth of these samples must be taken at farm level. Each sub-group of Group A must be checked each year using a minimum of 5 % of the total number of samples to

be collected for Group A. The balance will be allocated according to the experience and background information of Romania.

**Group B:** 50 % of the total samples, 30 % of samples must be checked for Group B 1 substances (antibiotics and sulfamides)

**4.4.5. Measures and terms of legislation as regards the different qualifications of animals and herds:** Not applicable.

**4.4.6. Control procedures and in particular rules on the movement of animals liable to be affected or contaminated by a given disease and the regular inspection of the holdings or areas concerned.**

According to the provisions of N.S.V.F.S.A. President Order 147/2006, Regulation 2160/2003/EC, the following measures are to be adopted in order to prevent the dissemination of *Salmonella enteritidis*, *Salmonella typhimurium*, into commercial holdings. Animals from infected flocks belonging to commercial holdings are to be kept isolated and special conditions apply for removal of these animals. No bird may leave the house concerned unless the competent authority has authorized the slaughter or/and destruction under supervision of slaughter in a slaughterhouse designated by the competent authority. All the birds in the house must be slaughtered in accordance with the provisions of the **REGULATION (EC) No. 853/2004 laying down specific hygiene rules for food of animal origin in order** to reduce as much as possible the risk of spreading salmonella.

**4.4.7. Measures and terms of legislation as regards the control (testing, vaccination ...) of the disease.**

The legal basis is the provisions of the **Regulation 2160/2003/EC**, **Regulation 517/2011/EC** and **Regulation 1177/2006 EC**.

The vaccination against Salmonella in laying hens it is not mandatory in Romania, but the CCA strongly recommend this. In case the owner of the flock wants to vaccinate against Salmonella he will send his vaccination programme to the local CA. If the vaccination programme is in accordance with the provision of Regulation 1177/2006 EC, the local CA approve it. Only the flocks which have an approved vaccination programme can vaccinate against Salmonella.

According to the provision of Regulation (EC) no. 1177/2006 CE in Romania the antimicrobials use for Salmonella control is forbidden.

A national *Salmonella* control programme was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 517/2011. The national control plan for *Salmonella* in laying flocks come into effect in January 2008.

**The NCP defines a flock as a single group or multiple groups of chickens which share the same production unit (i.e. using the same air-space or range area). Where housing systems are not typical, the situation is likely to be assessed on a case by case basis. Multiple groups of chickens which have ‘beak-to-beak’ contact (inside or outside the house) are likely to be treated as a single flock.**

All laying hens flocks will be included in the national control programme unless exempted in Regulation (EC) No 2160/2003 under Article 1.3, i.e. eggs produced for private domestic consumption, or where there is direct supply of small quantities of products to the final consumer or to local retail establishments directly supplying the primary products to the final consumer.

**Operators** will be required to implement the sampling programme in the Annex to EC Regulation 517/2011 (**self-control sampling**). For convenience the ‘Sampling protocol’ is repeated in **THE SAMPLING PROTOCOL**, showed below.

## **SAMPLING PROTOCOL.**

For each flock\*

### **Samples taken by operators (self control) during the rearing phase of layers.**

#### **A.Day old**

(a) One chick box liner, up to a maximum of 10, for every 500 chicks delivered from each hatchery. Samples taken on the day of arrival.

(b) The carcasses of all chicks, up to a maximum of 60, from each hatchery which are dead on arrival.

#### **2 weeks before entering laying phase**

A minimum of 2 pairs of boot swabs per house, or composite faeces sample taken according to the table below.

The number of sites from which separate faeces samples (minimum 1g each) are to be taken in order to make a composite sample shall be as follows:

| <b>Number of birds kept in a building</b> | <b>Number of faeces samples to be taken in the building</b> |
|---|---|
| 1-24                                      | Number equal to the number of birds up to a maximum of 20   |
| 25-29                                     | 20  |
| 30-39                                     | 25  |
| 40-49                                     | 30  |
| 50-59                                     | 35  |
| 60-89                                     | 40  |
| 90-199                                    | 50  |
| 200-499                                   | 55  |
| 500 or more                               | 60  |



Samples to be sent same day to Authorised Laboratory for testing for *Salmonella*. If samples are not dispatched on the day of collection to the laboratory they must be stored at 4° C but not frozen, and must be submitted within 48 hours of collection.

Boot swabs, faeces, and dust samples will be prepared according to the methods outlined in Section 3 of the Annex to Regulation (EC) No 517/2011.

### **Monitoring in Laying Flocks by the operator (self-control)**

Monitoring by operators shall take place according to Regulation (EC) No 517/2011 Annex Point 2 Monitoring in Laying Flocks every 15 weeks starting when the birds are 22 – 26 weeks of age.

### **Official sampling**

One sample will be taken under the control of the Competent Authority for Regulation 2160/2003 from one layer flock on each holding with more than 1000 birds during the period of production of eggs for human consumption as specified in 2.1 of Annex to Commission Regulation (EC) No 517/2011.

Official samples will include a sample of dust (or when not available an additional sample of faecal material) in compliance with 2.1 and 2.2 of Annex to Commission Regulation (EC) No 517/2011.

The use of antimicrobials (as defined in Regulation (EC) No 1177/2006) will be checked when the official sample is taken. If the flock is under antimicrobial medication for animal health or animal welfare reasons, the flock will be sampled again after the withdrawal period of the product specified in the Marketing Authorization. Flock owners are required to keep records of antimicrobial used and to make these records available to the competent authorities.

The records of samples harvest by the operator will be made available for inspection to the Competent Authority or its agent and will provide details of the identity of the flock sampled, date of sample, slaughter date, type of sample, laboratory carrying out the examination, and the result. The number of flocks on the holding and the number of birds present will be recorded.

When an official sample is taken it may replace the sample required to be taken by the operator.

## **4. Measures of the submitted programme**

### **Transport and preparation of the samples:**

Samples shall be sent to the approved laboratories within 24 hours after collection or in exceptional cases in maxim 48 hours after its had been harvest. The samples shall be kept refrigerated until its will be sent at the laboratory

The boot swabs, faeces, and dust samples shall be carefully unpacked to avoid dislodging adherent faecal material, pooled and placed in 225 ml buffered peptone water (BPW) which has been pre-warmed to room temperature. The sample shall be swirled to fully saturate it and culture shall be continued by using the detection method, described below.

#### **Detection method:**

The Salmonella spp. detection from faeces of animals made by the NRL for Salmonella in animals or C.S.V.F.S.L. is made according to PSO 001 - "Isolation and identification of the Salmonellosis" done by RENAR in 2006 (accreditation certificate no. 222 L/2006) and revised in May 2009. The procedure is in accordance with SR EN ISO 6579:2003 / A1: 2007 and with the standard procedure of CRL - Salmonella, Bilthoven, Netherlands.

In that detection method, a semi-solid medium (modified semi-solid Rappaport-Vassiladis medium, MSR/V) is used as the single selective enrichment medium.

#### **Serotyping**

At least one isolate from each positive sample shall be serotyped, following the Kaufmann-White scheme.

#### **Storage of strains:**

At least one isolated strain per house and per year shall be collected by the NRL and stored for future phage typing or anti-microbial susceptibility testing, using the normal methods for culture collection, which must ensure integrity of the strains for a minimum of two years.

#### **Use of vaccines**

Live Salmonella vaccines shall not be used in the framework of national control programme where the manufacturer does not provide an appropriate method to distinguish bacteriological wild-type strains of salmonella from vaccine strains.

Live salmonella vaccines shall not be used in the framework of national control programmes in laying hens during production unless the safety of the use has been demonstrated and they are authorised for such purpose in accordance with Directive 2001/82/EC.

#### **4.4.8. Measures and terms of legislation as regards the compensation for owners of slaughtered and killed animals:**

In laying flocks of *Gallus gallus* financial assistance is provided in the context of the control programme. This is set up by the **Gouvernement Decision no. 1214/2009** regarding the methodology for determining and paying compensation to be paid to owners of slaughtered animals, killed or otherwise affected by the rapid liquidation of outbreaks of animal diseases.

#### **4.4.9. Information and assessment on bio-security measures management and infrastructure in place in the flocks/holdings involved**

This information was given in the point 4.4.

## **5. General description of the benefits:**

### **Costs of the programme**

This programme is in accordance with Council Decision 2009/470/ EEC, Commission Decision 2008/425/EC and Council Decision 92/65/ EEC. The detailed financial costs for the national control programme of Salmonella in laying hens are laid down in point 8.

### **Benefits of the programme**

The overall aim of the Salmonella National Control Programme is to control the occurrence of Salmonella in the eggs sector on a very low level and thereby protect humans against infection with food-borne salmonellas. It has been known that poultry often harbour latent infections with Salmonella, which may pose a serious human health risk.

The anticipated benefits of this programme reducing the *Salmonella enteritidis* and *Salmonella typhimurium* are the minimising of human health problems and a consequent reduction in suffering, mortality and health service costs.

## **6. Data on the epidemiological evolution during the last year:**

## 6.1. Evolution of zoonotic salmonellosis

### 6.1.1 Data on evolution of zoonotic salmonellosis

Year: 2009

Situation on date: 31.12.2009

Animal species: laying hens (*Gallus gallus*)

Disease: zoonotic salmonellosis

| Region<br>(County+NUTS<br>CODE) | Type of<br>flocks(b) | Total<br>number<br>of<br>flocks<br>© | Total<br>number of<br>animals | Total<br>number of<br>flocks<br>under the<br>programme | Total<br>number of<br>animals<br>under the<br>programme | Number<br>of<br>flocks<br>checke<br>d(d) | Number of positive<br>flocks (e)                            |                           | Number of flocks<br>depopulated                                      |                           | Total number of animals<br>slaughtered or destroyed         |                       | Quantity of the<br>eggs destroyed<br>(number or kg)                  |                           | Quantity of eggs channelled to<br>egg products (number or kg) |                       |
|---------------------------------|----------------------|--------------------------------------|-------------------------------|--|---|--|---|---------------------------|--|---------------------------|---|-----------------------|--|---------------------------|---|-----------------------|
|                                 |                      |                                      |                               |  |   |  | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>seroty<br>pes(g) | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>serotypes(g) | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) | Serotypes<br>targeted in<br>the control<br>programe<br>(f)    | Other<br>serotypes(g) |
| RO121-AB                        | Laying<br>hens       | 5                                    | 138860                        | 5  | 138860  | 5  | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO421-AR                        | Laying<br>hens       | 6                                    | 52250                         | 6  | 52250   | 6  | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO311-AG                        | Laying<br>hens       | 11                                   | 183162                        | 11   | 183162  | 11                                       | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO211-BC                        | Laying<br>hens       | 15                                   | 103409                        | 15   | 103409  | 15                                       | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO111-BH                        | Laying<br>hens       | 5                                    | 90000                         | 2  | 60000   | 2  | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO112-BN                        | Laying<br>hens       | 4                                    | 90500                         | 4  | 90500   | 4  | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO212-BT                        | Laying<br>hens       | 1                                    | 21757                         | 1  | 21757   | 1  | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO221-BR                        | Laying<br>hens       | 36                                   | 1065613                       | 36   | 1065613   | 36                                       | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO122-BV                        | Laying<br>hens       | 6                                    | 87942                         | 6  | 87942   | 6  | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |
| RO321-BUC                       | Laying<br>hens       | 0                                    | 0                             | 0  | 0   | 0  | 0   | 0                         | 0  | 0                         | 0   | 0                     | 0  | 0                         | 0   | 0                     |

|          |             |    |         |    |         |    |             |   |   |   |   |       |   |   |   |   |
|----------|-------------|----|---------|----|---------|----|-------------|---|---|---|---|-------|---|---|---|---|
| RO222-BZ | Laying hens | 0  | 0       | 0  | 0       | 0  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO312-CL | Laying hens | 0  | 0       | 0  | 0       | 0  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO422-CS | Laying hens | 1  | 38901   | 1  | 38901   | 1  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO113-CJ | Laying hens | 27 | 119598  | 27 | 119598  | 27 | 1 suspicion | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO223-CT | Laying hens | 16 | 372079  | 16 | 372079  | 11 | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO123-CV | Laying hens | 10 | 184156  | 10 | 184156  | 10 | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO313-DB | Laying hens | 29 | 689216  | 29 | 689216  | 29 | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO411-DJ | Laying hens | 0  | 0       | 0  | 0       | 0  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO224-GL | Laying hens | 33 | 522418  | 20 | 507678  | 20 | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO314-GR | Laying hens | 36 | 638718  | 36 | 638718  | 36 | 0           | 4 | 0 | 4 | 0 | 49920 | 0 | 0 | 0 | 0 |
| RO412-GJ | Laying hens | 6  | 28190   | 6  | 28190   | 6  | 1           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO124-HR | Laying hens | 1  | 19601   | 1  | 19601   | 0  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO423-HD | Laying hens | 23 | 273863  | 23 | 278363  | 23 | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO315-IL | Laying hens | 9  | 224994  | 4  | 224994  | 4  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO213-IS | Laying hens | 3  | 35200   | 3  | 35200   | 2  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO322-IF | Laying hens | 0  | 0       | 0  | 0       | 0  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO114-MM | Laying hens | 19 | 202632  | 19 | 202632  | 19 | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO413-MH | Laying hens | 1  | 10000   | 1  | 10000   | 1  | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO125-MS | Laying hens | 29 | 191.970 | 29 | 191.970 | 29 | 0           | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |

|              |             |     |         |     |         |     |            |   |   |   |   |       |   |   |   |   |
|--------------|-------------|-----|---------|-----|---------|-----|------------|---|---|---|---|-------|---|---|---|---|
| RO214-NT     | Laying hens | 13  | 82795   | 13  | 82795   | 13  | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO414-OT     | Laying hens | 12  | 196301  | 12  | 196301  | 12  | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO316-PH     | Laying hens | 4   | 13429   | 4   | 13429   | 4   | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO116-SJ     | Laying hens | 4   | 110947  | 4   | 110947  | 4   | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO115-SM     | Laying hens | 13  | 125510  | 13  | 125510  | 13  | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO126-SB     | Laying hens | 21  | 164538  | 21  | 164538  | 21  | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO215-SV     | Laying hens | 2   | 4000    | 2   | 4000    | 2   | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO317-TR     | Laying hens | 3   | 180000  | 3   | 180000  | 3   | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO424-TM     | Laying hens | 21  | 182510  | 21  | 182510  | 21  | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO225-TL     | Laying hens | 0   | 0       | 0   | 0       | 0   | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO415-VL     | Laying hens | 3   | 49300   | 3   | 49300   | 3   | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO226-VN     | Laying hens | 5   | 131913  | 5   | 131913  | 5   | 0          | 1 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| RO216-VS     | Laying hens | 20  | 449729  | 20  | 449729  | 15  | 0          | 0 | 0 | 0 | 0 | 0     | 0 | 0 | 0 | 0 |
| <b>Total</b> | Laying hens | 453 | 7076001 | 432 | 7035761 | 420 | 1suspicion | 5 | 0 | 4 | 0 | 49920 | 0 | 0 | 0 | 0 |

Year: 2010

Situation on date: 31.12.2010

Animal species: laying hens (*Gallus gallus*)

Disease: zoonotic salmonellosis

| Region<br>(County+NU<br>TS CODE) | Type<br>of<br>flocks(<br>b) | Total<br>numb<br>er of<br>flocks<br>© | Total<br>number of<br>animals | Total<br>number of<br>flocks<br>under the<br>program<br>me | Total<br>number of<br>animals<br>under the<br>program<br>me | Numb<br>er of<br>flocks<br>check<br>ed(d) | Number of<br>positive flocks<br>(e)  |                           | Number of<br>flocks<br>depopulated   |                           | Total number of<br>animals<br>slaughtered or<br>destroyed                      |                           | Quantity of the<br>eggs destroyed<br>(number or kg)                     |                           | Quantity of eggs<br>channelled to egg<br>products (number<br>or kg)     |                           |
|----------------------------------|-----------------------------|---------------------------------------|-------------------------------|--|---|---|--|---------------------------|--|---------------------------|--|---------------------------|---|---------------------------|---|---------------------------|
|                                  |                             |                                       |                               |  |   |   | Seroty<br>pes<br>target<br>ed in<br>the<br>contro<br>l<br>progra<br>mme<br>(f) | Other<br>seroty<br>pes(g) | Seroty<br>pes<br>targete<br>d in<br>the<br>control<br>progra<br>mme<br>(f) | Other<br>seroty<br>pes(g) | Seroty<br>pes<br>target<br>ed in<br>the<br>contro<br>l<br>progra<br>mme<br>(f) | Other<br>seroty<br>pes(g) | Serotyp<br>es<br>targete<br>d in<br>the<br>control<br>progra<br>mme (f) | Other<br>seroty<br>pes(g) | Serotype<br>s<br>targete<br>d in<br>the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) |
| RO421-AR                         | Laying<br>Hens              | 7                                     | 57451                         | 7  | 57451   | 7   | 0  | 0                         | 0  | 0                         | 0  | 0                         | 0   | 0                         | 0   | 0                         |
| RO121-AB                         | Laying<br>Hens              | 9                                     | 144083                        | 9  | 144083  | 9   | 0  | 0                         | 0  | 0                         | 0  | 0                         | 0   | 0                         | 0   | 0                         |
| RO311-AG                         | Laying<br>Hens              | 7                                     | 110000                        | 7  | 110000  | 7   | 0  | 0                         | 0  | 0                         | 0  | 0                         | 0   | 0                         | 0   | 0                         |
| RO211-BC                         | Laying<br>Hens              | 3                                     | 12000                         | 3  | 12000   | 3   | 0  | 0                         | 0  | 0                         | 0  | 0                         | 0   | 0                         | 0   | 0                         |
| RO111-BH                         | Laying<br>Hens              | 21                                    | 104500                        | 21   | 104500  | 6   | 0  | 0                         | 0  | 0                         | 0  | 0                         | 0   | 0                         | 0   | 0                         |
| RO112-BN                         | Laying<br>Hens              | 5                                     | 94632                         | 5  | 94632   | 5   | 0  | 0                         | 0  | 0                         | 0  | 0                         | 0   | 0                         | 0   | 0                         |
| RO212-BT                         | Laying<br>Hens              | 1                                     | 15727                         | 1  | 15727   | 1   | 0  | 0                         | 0  | 0                         | 0  | 0                         | 0   | 0                         | 0   | 0                         |



|           |             |    |         |    |         |    |   |           |   |   |       |   |        |   |   |   |
|-----------|-------------|----|---------|----|---------|----|---|-----------|---|---|-------|---|--------|---|---|---|
| RO221-BR  | Laying Hens | 35 | 773321  | 35 | 773321  | 28 | 0 | <b>13</b> | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO122-BV  | Laying Hens | 8  | 96334   | 8  | 96334   | 8  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO321-BUC | Laying Hens | 0  | 0       | 0  | 0       | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO222-BZ  | Laying Hens | 0  | 0       | 0  | 0       | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO312-CL  | Laying Hens | 0  | 0       | 0  | 0       | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO422-CS  | Laying Hens | 1  | 193455  | 1  | 193455  | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO113-CJ  | Laying Hens | 27 | 119598  | 27 | 119598  | 27 | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO223-CT  | Laying Hens | 16 | 341298  | 16 | 341298  | 9  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO123-CV  | Laying Hens | 60 | 5951280 | 13 | 1289444 | 13 | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO313-DB  | Laying Hens | 29 | 699509  | 29 | 699509  | 29 | 0 | <b>2</b>  | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO411-DJ  | Laying Hens | 0  | 0       | 0  | 0       | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO224-GL  | Laying Hens | 22 | 592956  | 22 | 592956  | 22 | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO314-GR  | Laying Hens | 42 | 545367  | 34 | 483767  | 30 | 0 | <b>2</b>  | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO412-GJ  | Laying Hens | 6  | 133035  | 6  | 133035  | 6  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO124-HR  | Laying Hens | 3  | 111307  | 1  | 87747   | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO423-HD  | Laying Hens | 13 | 263000  | 13 | 263000  | 5  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO315-IL  | Laying Hens | 12 | 305590  | 12 | 305590  | 12 | 2 | 0         | 2 | 0 | 18782 | 0 | 175950 | 0 | 0 | 0 |
| RO213-IS  | Laying Hens | 4  | 34270   | 3  | 34270   | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO322-IF  | Laying Hens | 0  | 0       | 0  | 0       | 0  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO114-MM  | Laying Hens | 21 | 192609  | 21 | 192609  | 21 | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |
| RO413-MH  | Laying Hens | 1  | 9800    | 1  | 9800    | 1  | 0 | 0         | 0 | 0 | 0     | 0 | 0      | 0 | 0 | 0 |

|              |             |            |                 |            |                |            |          |           |          |          |              |          |               |          |          |          |
|--------------|-------------|------------|-----------------|------------|----------------|------------|----------|-----------|----------|----------|--------------|----------|---------------|----------|----------|----------|
| RO125-MS     | Laying Hens | 30         | 211000          | 30         | 211000         | 30         | 1        | 0         | 1        | 0        | 850          | 0        | 4100          | 0        | 0        | 0        |
| RO214-NT     | Laying Hens | 13         | 132323          | 13         | 132323         | 13         | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO414-OT     | Laying Hens | 11         | 258049          | 11         | 258049         | 11         | 0        | 1         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO316-PH     | Laying Hens | 5          | 13353           | 5          | 13353          | 5          | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO116-SJ     | Laying Hens | 9          | 120400          | 9          | 120400         | 9          | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO115-SM     | Laying Hens | 10         | 133000          | 10         | 133000         | 10         | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO126-SB     | Laying Hens | 19         | 181823          | 19         | 181823         | 18         | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO215-SV     | Laying Hens | 1          | 2000            | 1          | 2000           | 1          | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO317-TR     | Laying Hens | 7          | 398357          | 7          | 398357         | 7          | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO424-TM     | Laying Hens | 18         | 181243          | 18         | 181243         | 18         | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO225-TL     | Laying Hens | 0          | 0               | 0          | 0              | 0          | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO415-VL     | Laying Hens | 6          | 108000          | 6          | 108000         | 6          | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO226-VN     | Laying Hens | 5          | 64473           | 5          | 64473          | 5          | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| RO216-VS     | Laying Hens | 20         | 455935          | 20         | 455935         | 11         | 0        | 0         | 0        | 0        | 0            | 0        | 0             | 0        | 0        | 0        |
| <b>Total</b> | Laying Hens | <b>507</b> | <b>13161078</b> | <b>449</b> | <b>8414082</b> | <b>393</b> | <b>3</b> | <b>18</b> | <b>3</b> | <b>0</b> | <b>19632</b> | <b>0</b> | <b>180050</b> | <b>0</b> | <b>0</b> | <b>0</b> |

Year: 2011

Situation on date: 31.12.2011

Animal species: laying hens (*Gallus gallus*)

Disease: zoonotic salmonellosis

| Region<br>(County+NU<br>TS CODE) | Type<br>of<br>flocks(<br>b) | Total<br>numb<br>er of<br>flock<br>s © | Total<br>number<br>of<br>animals | Total<br>number<br>of flocks<br>under<br>the<br>program<br>me | Total<br>number<br>of<br>animals<br>under<br>the<br>program<br>me | Numb<br>er of<br>flocks<br>check<br>ed(d) | Number of<br>positive<br>flocks (e)  |                                       | Number of<br>flocks<br>depopulated   |                                       | Total number of<br>animals<br>slaughtered or<br>destroyed       |                               | Quantity of the<br>eggs destroyed<br>(number or kg)             |                               | Quantity of eggs<br>channelled to egg<br>products (number<br>or kg) |                           |
|----------------------------------|-----------------------------|--|----------------------------------|---|---|---|--|---------------------------------------|--|---------------------------------------|---|-------------------------------|---|-------------------------------|---|---------------------------|
|                                  |                             |  |                                  |   |   |   | Serot<br>ypes<br>target<br>ed in<br>the<br>contr<br>ol<br>progr<br>amme<br>(f) | Oth<br>er<br>ser<br>oty<br>pes<br>(g) | Serot<br>ypes<br>target<br>ed in<br>the<br>contr<br>ol<br>progr<br>amme<br>(f) | Oth<br>er<br>ser<br>oty<br>pes<br>(g) | Serotypes<br>targeted<br>in the<br>control<br>programm<br>e (f) | Other<br>serot<br>ypes(<br>g) | Serotypes<br>targeted<br>in the<br>control<br>programm<br>e (f) | Other<br>serot<br>ypes(<br>g) | Serotyp<br>es<br>targeted<br>in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) |
| RO121-AB                         | laying<br>hens              | 9                                      | 145104                           | 7   | 139642  | 6   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |
| RO421-AR                         | laying<br>hens              | 14                                     | 102000                           | 4   | 29000   | 4   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |
| RO311-AG                         | laying<br>hens              | 12                                     | 298944                           | 7   | 202724  | 7   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |
| RO211-BC                         | laying<br>hens              | 3                                      | 16500                            | 3   | 16500   | 3   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |
| RO111-BH                         | laying<br>hens              | 7                                      | 49575                            | 7   | 49575   | 7   | 1  | 0                                     | 1  | 0                                     | 820   | 0                             | 1500  | 0                             | 0   | 0                         |
| RO112-BN                         | laying<br>hens              | 4                                      | 65539                            | 4   | 65539   | 4   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |
| RO212-BT                         | laying<br>hens              | 1                                      | 18240                            | 1   | 18240   | 1   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |
| RO221-BR                         | laying<br>hens              | 34                                     | 996765                           | 34  | 996765  | 34  | 5  | 15                                    | 5  | 0                                     | 123295  | 0                             | 833399  | 0                             | 960385  | 0                         |
| RO122-BV                         | laying<br>hens              | 10                                     | 158736                           | 10  | 158736  | 9   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |
| RO321-BUC                        | laying<br>hens              | 0                                      | 0                                | 0   | 0   | 0   | 0  | 0                                     | 0  | 0                                     | 0   | 0                             | 0   | 0                             | 0   | 0                         |

|          |             |    |         |    |         |    |   |   |   |   |       |   |         |   |   |   |
|----------|-------------|----|---------|----|---------|----|---|---|---|---|-------|---|---------|---|---|---|
| RO222-BZ | laying hens | 0  | 0       | 0  | 0       | 0  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO312-CL | laying hens | 5  | 55082   | 5  | 55082   | 5  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO422-CS | laying hens | 7  | 329404  | 7  | 329404  | 6  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO113-CJ | laying hens | 25 | 125300  | 23 | 119470  | 7  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO223-CT | laying hens | 14 | 422480  | 14 | 422480  | 14 | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO123-CV | laying hens | 0  | 0       | 0  | 0       | 0  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO313-DB | laying hens | 38 | 1000000 | 38 | 1000000 | 38 | 0 | 4 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO411-DJ | laying hens | 4  | 11862   | 4  | 11862   | 4  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO224-GL | laying hens | 19 | 783000  | 19 | 783000  | 19 | 2 | 3 | 2 | 0 | 52596 | 0 | 1084200 | 0 | 0 | 0 |
| RO314-GR | laying hens | 83 | 1411578 | 36 | 1411578 | 36 | 0 | 5 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO412-GJ | laying hens | 2  | 135456  | 2  | 135456  | 2  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO124-HR | laying hens | 4  | 106000  | 4  | 106000  | 4  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO423-HD | laying hens | 18 | 181050  | 18 | 181050  | 17 | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO315-IL | laying hens | 16 | 422308  | 16 | 422308  | 16 | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO213-IS | laying hens | 3  | 33000   | 2  | 33000   | 3  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO322-IF | laying hens | 0  | 0       | 0  | 0       | 0  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO114-MM | laying hens | 26 | 222610  | 26 | 222610  | 18 | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO413-MH | laying hens | 0  | 0       | 0  | 0       | 0  | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO125-MS | laying hens | 71 | 1500000 | 26 | 549295  | 26 | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO214-NT | laying hens | 15 | 186849  | 13 | 162957  | 12 | 0 | 0 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |
| RO414-OT | laying hens | 11 | 237540  | 11 | 237540  | 11 | 0 | 3 | 0 | 0 | 0     | 0 | 0       | 0 | 0 | 0 |

|              |                    |            |                 |            |                |            |          |           |          |          |               |          |                |          |               |          |
|--------------|--------------------|------------|-----------------|------------|----------------|------------|----------|-----------|----------|----------|---------------|----------|----------------|----------|---------------|----------|
| RO316-PH     | laying hens        | 0          | 0               | 0          | 0              | 0          | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO116-SJ     | laying hens        | 10         | 133900          | 10         | 133900         | 10         | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO115-SM     | laying hens        | 18         | 286000          | 10         | 156000         | 10         | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO126-SB     | laying hens        | 14         | 118934          | 13         | 118934         | 13         | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO215-SV     | laying hens        | 3          | 3600            | 3          | 3000           | 2          | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO317-TR     | laying hens        | 16         | 428552          | 16         | 428552         | 16         | 0        | 1         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO424-TM     | laying hens        | 19         | 200939          | 19         | 200939         | 9          | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO225-TL     | laying hens        | 0          | 0               | 0          | 0              | 0          | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO415-VL     | laying hens        | 0          | 0               | 0          | 0              | 0          | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO226-VN     | laying hens        | 3          | 64178           | 3          | 64178          | 2          | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| RO216-VS     | laying hens        | 36         | 685350          | 36         | 685350         | 36         | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| IDAH         | laying hens        | 0          | 0               | 0          | 0              | 0          | 0        | 0         | 0        | 0        | 0             | 0        | 0              | 0        | 0             | 0        |
| <b>Total</b> | <b>laying hens</b> | <b>574</b> | <b>10936375</b> | <b>451</b> | <b>9650666</b> | <b>411</b> | <b>8</b> | <b>31</b> | <b>8</b> | <b>0</b> | <b>176711</b> | <b>0</b> | <b>1919099</b> | <b>0</b> | <b>960385</b> | <b>0</b> |

## 6.4 Data on vaccination programme

Data not available

## 7. Targets

### 7.1. Targets related to testing :to investigate the presence of Salmonella in laying flocks

**Disease (a): zoonotic Salmonella**

**Animal species:**

| Region (b)   | Type of the test (c) | Target population (d) | Type of sample (e)     | Objective (f) | Number of planned tests |
|--------------|----------------------|-----------------------|------------------------|---------------|-------------------------|
| All regions  | Bacteriological      | Laying hens           | Faeces boot/sock swabs | Surveillance  | 2000                    |
|              | Bacteriological      | Laying hens           | Faeces boot/sock swabs | Confirmation  | 350                     |
| <b>Total</b> | -                    | -                     | -                      | -             |                         |

(a) Disease and species if necessary

(b) Region as defined in the approved eradication programme of the Member State

(c) Description of the test

(d) Specification of the targeted species and the categories of targeted animals

(e) Description of the sample (faeces)

(f) Description of the objective ( e.g. qualification, surveillance, confirmation of suspected cases, monitoring of campaigns, seroconversion, control on deleted vaccines, testing of vaccine , control of vaccination

## 7.1.2. Targets on testing of laying flocks

Year: 2011

Animal species: Gallus gallus

Disease/infection(a): Zoonotic Salmonella

| Region      | Type of flock <sup>(b)</sup> | Total number of flocks <sup>(c)</sup> | Total number of animals | Total n° of flocks under the programme | Total n° of animals under the programme | Expected n° of flocks to be checked <sup>(d)</sup> | Number of flocks <sup>(d)</sup> expected to be positive <sup>(a)</sup> |      |      | Number of flocks expected to be depopulated |      | Total n° of animals expected to be slaughtered or destroyed <sup>(f)</sup> |      | Expected quantity of eggs to be destroyed (number or kg) <sup>(d)</sup> |    | Expected quantity of eggs channelled to egg products (number or kg.) <sup>(d)</sup> |      |
|-------------|------------------------------|---------------------------------------|-------------------------|--|---|--|--|------|------|---|------|--|------|---|----|---|------|
|             |                              |                                       |                         |  |   |  | (a1)   | (a2) | (a3) | a4  | (a3) | (a4)   | (a3) | (a4)  | a3 | (a4)  | (a3) |
| All regions | Laying hens                  | 600                                   | 10.000.000              | 600                                    | 10.000.000                              | 600  | 5  | 5    | 50   | 10  | 50   | 200000   | 0    | 2000000   | 0  | 1000000   | 0    |
| Total       | Laying hens                  | 600                                   | 10.000.000              | 600                                    | 10.000.000                              | 600  | 5  | 5    | 50   | 10  | 50   | 200000   | 0    | 2000000   | 0  | 1000000   | 0    |

- (a) For zoonotic salmonella indicate the serotypes covered by the control programmes: (a1) for Salmonella Enteritidis, (a2) for Salmonella Typhimurium, (a3) for other serotypes-specify as appropriate, (a4) for Salmonella Enteritidis or Salmonella Typhimurium.
- (b) For example, breeding flocks (rearing, adult flocks), production flocks, laying hen flocks, etc. Flocks equals herds or as appropriate.
- (c) Total number of flocks existing in the region including eligible flocks and non-eligible flocks for the programme.
- (d) Check means to perform a flock level test under the programme for the presence of Salmonella. In this column a flock should not be counted twice even if it has been checked more than once.
- (e) If a flock has been checked, in accordance with footnote (d), more than once, a positive sample should be taken into account only once.

## 7.2. Targets on vaccination

### 7.2.1. Targets on vaccination of laying hens flocks- Data not available

Disease <sup>(a)</sup>: Zoonotic Salmonella

Animal species: gallus gallus

| Region       | Total number of herds in vaccination or programme | Total number of animals in vaccination programme | Targets on vaccination programme     |                                       |   |  |
|--------------|---|--|--------------------------------------|---------------------------------------|---|--|
|              |   |  | No of herds in vaccination programme | No of herds expected to be vaccinated | No of animals expected to be vaccinated | No of doses of vaccine expected to be administered |
| All regions  | -   | -  | -                                    | -                                     | -                                       | -  |
| <b>Total</b> | -   | -  | -                                    | -                                     | -                                       | -  |

**The vaccination is not mandatory and the costs regarding purchase of vaccine doses and the vaccination are incurred by the business operators.**



## 8. Detailed analysis of the cost of the programme<sup>1</sup>

Costs mentioned below are estimated for a one-year period (1 of January 2013-31 of December 2013 )

| <b>Costs related to</b>          | <b>Specification</b>  | <b>Number of units</b> | <b>Unitary cost in €</b> | <b>Total amount in €</b> | <b>Community funding requested (yes/no)</b> |
|----------------------------------|---|------------------------|--------------------------|--------------------------|---|
| <b>1. Testing</b>                |   |                        |                          |                          |   |
| <b>1.1. Cost of the analysis</b> |   |                        |                          |                          |   |
|                                  | Test: bacteriological exam for Salmonella spp.detection   | 2000                   | 16.70                    | 33400                    | YES   |
|                                  | Test : Salmonella spp. serotyping   | 350                    | 64.97                    | 22739.5                  | YES   |
|                                  | Test : the detection of antimicrobials or bacterial growth inhibitory effect in tissues from birds from flocks tested for Salmonella with HPLC method | 300                    | 47.44                    | 14232                    | YES   |
| <b>1.2. Cost of sampling</b>     | Official sample harvest   | 2000                   | 0.5                      | 1000                     | YES   |
|                                  | Disposable sterile containers for sampling of faeces  | 2000                   | 0,5                      | 1000                     | NO  |
|                                  | One use gloves-pairs  | 2000                   | 0,1                      | 200                      | NO  |
|                                  | Boot swabs  | 1000                   | 2                        | 2000                     | NO  |
| <b>1.3. Other costs</b>          | Overcoats   | 2000                   | 1                        | 2000                     | NO  |
| <b>2. Vaccination</b>            |   |                        |                          |                          |   |
| <b>2.1. Purchase of vaccine</b>  |   | -                      | -                        | -                        | -   |
| <b>2.2. Distribution costs</b>   |   | -                      | -                        | -                        | -   |
| <b>2.3. Administering costs</b>  |   | -                      | -                        | -                        | -   |

|  |   |                |             |                            |            |
|--|---|----------------|-------------|----------------------------|------------|
| <b>2.4. Control costs</b>  |   | -              | -           | -                          | -          |
| <b>3. Slaughter and destruction</b>  |   |                |             |                            |            |
| <b>3.1. Compensation of animals</b>  | Compensation for the value of a commercial laying bird of <i>Gallus gallus</i> culled   | <b>200000</b>  | <b>4.4</b>  | <b>880000</b>              | <b>YES</b> |
|  | Compensation for table eggs of <i>Gallus gallus</i> destroyed   | <b>2000000</b> | <b>0.08</b> | <b>160000</b>              | <b>YES</b> |
| <b>3.2. Transport costs</b>  |   | -              | -           | -                          | -          |
| <b>3.3. Destruction costs</b>  |   | -              | -           | -                          | -          |
| <b>3.4. Loss in case of slaughtering</b>   |   | -              | -           | -                          | -          |
| <b>3.5 Costs from treatment of products (eggs, hatching eggs, heat treat of broilers, etc)</b> |   | -              | -           | -                          | -          |
| <b>4. Cleaning and disinfection</b>  |   |                |             |                            |            |
|  | Test: Bacteriological test to verify the efficiency of disinfection of poultry houses after depopulation of a Salmonella-positive flock | <b>500</b>     | <b>4.1</b>  | <b>2050</b>                | <b>YES</b> |
| <b>5. Salaries (staff contracted for the programme only)</b>                                   |   |                |             |                            |            |
|  |   |                |             |                            |            |
| <b>6. Consumables and specific equipment</b>   |   |                |             |                            |            |
| <b>7. Other costs</b>  |   |                |             |                            |            |
| <b>TOTAL</b>   |   |                |             | <b>1113422</b><br><b>€</b> | <b>YES</b> |



# Romanian National Control Programme for *Salmonella* in breeding hens (*Gallus gallus*)

2013

## Part A

### a. Aim of the programme

Starting from 1 January 2010, the Romanian target, as referred to in Article 4(1) of Regulation (EC) No 2160/2003, for the reduction of *Salmonella* spp. in breeding flocks of *Gallus gallus* (the Union target) shall be a reduction to 1 % or less of the maximum percentage of adult breeding flocks of *Gallus gallus* remaining positive for *Salmonella enteritidis*, *Salmonella infantis*, *Salmonella hadar*, *Salmonella typhimurium*, including also the monophasic strains of *S. Typhimurium* and *Salmonella virchow* (the relevant salmonella serotypes).

### b. Sampling programme

The National Control Programme for *Salmonella* in breeding hens encompasses the following serovars of zoonotic *Salmonella*: *Salmonella enteritidis*, *Salmonella typhimurium*, *Salmonella infantis*, *Salmonella hadar* and *Salmonella virchow*.

The sampling programme will be in accordance to Regulation **2160/2003 EC** and Regulation **200/2010 EC**. We have also taken into account the provisions of Regulation **1177/2006 EC** and Regulation (EC) **no 213/2009** amending Regulation (EC) **no 2160/2003** of the European Parliament and of the Council and Regulation (EC) no 1003/2005 as regards the control and testing of *Salmonella* in breeding flocks of *Gallus gallus* and turkeys.

The programme was elaborated in compliance with the requirements of Commission Regulation (EU) **No 200/2010** implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards a Union target for the reduction of the prevalence of *Salmonella* serotypes in adult breeding flocks of *Gallus gallus*.

**c. Demonstrate the evidence that it complies with the specific requirements laid down in Parts C of Annex II to Regulation (EC) no 2160/2003 (and Commission Regulation (EC) No 213/2009)**

## **Specific requirements concerning breeding flocks of *Gallus gallus***

1. Starting with 1 January 2010 the measures laid down below in points 3 to 5 must be taken whenever the analysis of samples carried out in accordance with point 2.2.2.1 and 2.2.2.2 of Annex of the Regulation 200/2010 indicates the presence of *Salmonella enteritidis* or *Salmonella typhimurium* in a breeding flock of *Gallus gallus* in the circumstances set out in point 2.

2. (a) If the competent authority has approved the method of analysis used for samples taken in accordance with part B of the Regulation 2160/2003/EC , it may require that the measures laid down in points 3 to 5 be taken when such analysis detects the presence of *Salmonella enteritidis* or *Salmonella typhimurium*.

(b) Otherwise, the measures laid down in points 3 to 5 must be taken whenever the competent authority confirms a suspicion of the presence of *Salmonella enteritidis* or *Salmonella typhimurium* arising from the analysis of samples carried out in accordance with part B of the Regulation 2160/2003/EC.

### **3. Non-incubated eggs from the flock must be destroyed.**

However, such eggs may be used for human consumption if they are treated in a manner that guarantees the elimination of *Salmonella enteritidis* and *Salmonella typhimurium* in accordance with Community legislation on food hygiene.

4. **All birds**, including day-old chicks, **in the flock must be slaughtered or destroyed so as to reduce as much as possible the risk of spreading Salmonella**. Slaughtering must be carried out in accordance with Community legislation on food hygiene. Products derived from such birds may be placed on the market for human consumption in accordance with Community legislation on food hygiene and, once applicable, part E of the Regulation 2160/2003. If not destined for human consumption, such products must be used or disposed of in accordance with Regulation (EC) No 1774/2002 of the European Parliament and of the Council, laying down health rules concerning animal by-products not intended for human consumption.

5. Where **eggs** for hatching from flocks in which *Salmonella enteritidis* or *Salmonella typhimurium* is present are still **present in a hatchery, they must be destroyed or treated** in accordance with Regulation (EC) No 1774/2002.

## 1. General

### 1.1 Summary referring to the occurrence of the salmonellosis in Romania

During 2011, a totally of 396 breeding flocks were tested for Salmonella infection from wich a no of 1535 were tested in official control and there were no flocks positive for Salmonella Typhimurium, including monophasic ST and Salmonella Enteritidis . The prevalence for the target serotypes in breeding flock in 2011 was 0%, which is low and below the Community target.

The incidence of various serotypes of Salmonella in breeding hens farms, in 2011 in Romania

| NO.          | SEROTYPE                          | NO. OF STRAINS in Gallus gallus |
|--------------|-----------------------------------|---------------------------------|
| 1            | S.Agona                           | 1                               |
| 2            | S.Amsterdam                       | 4                               |
| 3            | S.Enteritidis                     | 7                               |
| 4            | S.Glostrup                        | 1                               |
| 5            | S.Infantis                        | 3                               |
| 6            | S.Kentucky                        | 4                               |
| 7            | S.Mbandaka                        | 8                               |
| 8            | S.Senftenberg                     | 3                               |
| 9            | S.Tennessee                       | 2                               |
| 10           | S.Thompson                        | 7                               |
| 11           | S.Typhimurium (vaccineted strain) | 2                               |
| <b>TOTAL</b> |                                   | <b>42</b>                       |

### 1.2 The structure and organization of the relevant Competent Authorities

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in breeding hens (*Gallus gallus*) flocks is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**

București, Str. Dudului, nr. 37, sect. 6, cod poștal 060603; tel: 0374.150.200, fax: 3124967; e-mail: [office@ansv.ro](mailto:office@ansv.ro)

The departments responsible for implementing the National Control Programme of Salmonella in breeding hens are:

- for implementing at national level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary General Directorate** ;
- in respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the verification of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**
- Concerning the slaughter of flocks, the **Hygiene and Veterinary Epidemiology Directorate** is responsible for public health protection, through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.
- In respect of EC Regulation No 183/2005 on feed hygiene at the central level is **The Directorate for Technical Coordination of Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**
- at county level for implementing the Regulation (EC) No 2160/2003,882/2004 and 183/2005 there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the national reference laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

#### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

##### **Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România

Tel: 0374.322.013 / 0374.322.000

Fax: 0214.113.394 E-mail: [office@idah.ro](mailto:office@idah.ro)

- **H.I.P.V.H.** (Hygiene Institute and Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

#### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED**

##### **Hygiene Institute and Veterinary Public Health**

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201

Tel: 021.252.46.51 Fax: 021.252.00.61 E-mail: [iispv@iispv.ro](mailto:iispv@iispv.ro)

### 1.3 Approved laboratories where samples collected within the programme are analyzed

**Diagnostics:** The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL is show below:

| Nr. Crt. | NAME                             | ADRESS   | TELEPHON/ E-MAIL  |
|----------|----------------------------------|--|---|
| 1        | <b>I.D.S.A</b>                   | București, Str. Dr.Staicovici, nr.63, sector 5, Cod 050557         | 0374/322.013<br><a href="mailto:office@idah.ro">office@idah.ro</a>  |
| 2        | <b>L.S.V.S.A. Alba</b>           | Alba Iulia, Str. Lalelelor nr.7A, Cod 510217, Jud. Alba            | 0258/835.915<br><a href="mailto:office-alba@ansvsa.ro">office-alba@ansvsa.ro</a>  |
| 3        | <b>L.S.V.S.A. Bacău</b>          | Bacau, Str Bucovinei, Nr 21, Jud. Bacău                            | 0234/586.233, int.111<br><a href="mailto:office-bacau@ansvsa.ro">office-bacau@ansvsa.ro</a>   |
| 4        | <b>L.S.V.S.A.Bistrița-Năsăud</b> | Bistrita, Str. Tarpiului 26, Cod 420062, Jud. Bistrița Năsăud      | 0263/206.027; 0263/224.974<br><a href="mailto:office-bistrita-nasaud@ansvsa.ro">office-bistrita-nasaud@ansvsa.ro</a>  |
| 5        | <b>L.S.V.S.A. Botoșani</b>       | Botosani , Str Tudor Vladimirescu Nr.17, Jud. Botoșani             | 0231/512.766<br><a href="mailto:office-botosani@ansvsa.ro">office-botosani@ansvsa.ro</a>  |
| 6        | <b>L.S.V.S.A. Brașov</b>         | Brasov, Str. Calea Feldioarei Nr. 20 A, Cod 500450, Jud.Brașov     | 0268/440.257<br><a href="mailto:office-brasov@ansvsa.ro">office-brasov@ansvsa.ro</a><br><a href="mailto:dsvbv@rdslink.ro">dsvbv@rdslink.ro</a>                    |
| 7        | <b>L.S.V.S.A. Brăila</b>         | Brăila, str. Calea Galați, nr.344, Cod 810385, Jud. Brăila         | 0239/610.689<br>dsv@braila.rdsnet.ro<br><a href="mailto:office-braila@ansvsa.ro">office-braila@ansvsa.ro</a>  |
| 8        | <b>L.S.V.S.A. Buzău</b>          | Buzau, Str. Horticolei, nr. 58 bis, Jud. Buzau                     | 0238-725001<br>0238-725002<br><a href="mailto:office-buzau@ansv.ro">office-buzau@ansv.ro</a>  |
| 9        | <b>L.S.V.S.A. Călărași</b>       | Călărași, Str. Prelungirea Dobrogei , nr. 4, Jud.Călărași          | 0242/313.676,<br><a href="mailto:office-calarasi@ansvsa.ro">office-calarasi@ansvsa.ro</a> ;<br><a href="mailto:office_dsv@satline.ro">office_dsv@satline.ro</a> ; |
| 10       | <b>L.S.V.S.A Constanța</b>       | Constanta, Sos. Mangaliei Nr. 78, Jud. Constanța                   | 0241-682-417,<br>email: <a href="mailto:dsvct@rdsct.ro">dsvct@rdsct.ro</a>  |
| 11       | <b>L.S.V.S.A. Cluj</b>           | Cluj-Napoca, Piata Marasti Nr. 1, Jud. Cluj                        | 0264-445729, 0264-448177<br><a href="mailto:office-cluj@ansvsa.ro">office-cluj@ansvsa.ro</a><br><a href="mailto:dsvcj@rdslink.ro">dsvcj@rdslink.ro</a>            |
| 11       | <b>L.S.V.S.A. Dâmbovița</b>      | Târgoviște, Str. I.C. Brătianu, nr. 35, Cod 130055, Jud. Dâmbovița | 0372/737818<br><a href="mailto:office-dambovita@ansvsa.ro">office-dambovita@ansvsa.ro</a>   |
| 13       | <b>L.S.V.S.A. Giurgiu</b>        | Giurgiu, Str. Bucuresti nr.72, Jud. Giurgiu                        | 0246/ 230.491<br><a href="mailto:office-giurgiu@ansvsa.ro">office-giurgiu@ansvsa.ro</a>   |
| 14       | <b>L.S.V.S.A. Gorj</b>           | Târgul-Jiu, Str.Ecaterina Teodoroiu Nr.523, Jud. Gorj              | 0253/226.033<br><a href="mailto:dsv@intergorj.ro">dsv@intergorj.ro</a><br><a href="mailto:office-gorj@ansvsa.ro">office-gorj@ansvsa.ro</a>                        |



|    |                             |  |  |
|----|-----------------------------|--|--|
| 15 | <b>L.S.V.S.A Hunedoara</b>  | Deva, Str. 22 Decembrie Nr. 226, Jud. Hunedoara                        | 0254-221145, 0254-230527<br><a href="mailto:office-hunedoara@ansvsa.ro">office-hunedoara@ansvsa.ro</a> |
| 16 | <b>L.S.V.S.A. Ialomița</b>  | Slobozia, str. Lacului, nr. 12 , Jud. Ialomița                         | 0243/ 232.069<br><a href="mailto:dsvsa-ialomita@ansvsa.ro">dsvsa-ialomita@ansvsa.ro</a>                |
| 17 | <b>L.S.V.S.A. Iași</b>      | Iași, Aleea Mihail Sadoveanu 10, Jud. Iași                             | 0232-267501<br><a href="mailto:office-iasi@ansvsa.ro">office-iasi@ansvsa.ro</a>                        |
| 18 | <b>L.S.V.S.A. Maramureș</b> | Baia Mare, Str. Vasile Alecsandri, Nr. 66, Jud. Maramureș              | 0262/ 224.031<br><a href="mailto:office-maramures@ansvsa.ro">office-maramures@ansvsa.ro</a>            |
| 19 | <b>L.S.V.S.A. Mureș</b>     | Târgu Mures, str. Podeni nr.10 cod 540253, Jud. Mureș                  | 0265/ 314.975<br><a href="mailto:office-mures@ansvsa.ro">office-mures@ansvsa.ro</a>                    |
| 20 | <b>L.S.V.S.A. Neamț</b>     | Piatra Neamț, Aleea Tiparului, nr. 12, cod 610263, Jud. Neamț          | 0233/ 223.259<br><a href="mailto:office-neamt@ansvsa.ro">office-neamt@ansvsa.ro</a>                    |
| 21 | <b>L.S.V.S.A. Prahova</b>   | Ploiesti, Str. Corlatesti, nr. 11, Jud. Prahova,                       | 0244 / 57.17.51<br><a href="mailto:office-prahova@ansvsa.ro">office-prahova@ansvsa.ro</a>              |
| 22 | <b>L.S.V.S.A. Satu Mare</b> | Satu Mare, Str. Lăcrimioarei, Nr. 37, cod poștal 440067, Jud.Satu Mare | 0261/ 715.956<br><a href="mailto:office-satu-mare@ansvsa.ro">office-satu-mare@ansvsa.ro</a>            |
| 23 | <b>L.S.V.S.A. Sibiu</b>     | Sibiu, str. Calea Șurii Mari, nr. 21, cod 550089, Jud. Sibiu           | 0269/223.069<br><a href="mailto:office-sibiu@ansvsa.ro">office-sibiu@ansvsa.ro</a>                     |
| 24 | <b>L.S.V.S.A. Suceava</b>   | Suceava, Str. Scurta Nr. 2, Jud Suceava                                | Tel: 0230-522848,<br><a href="mailto:dsvsv@suceava.rdsnet.ro">dsvsv@suceava.rdsnet.ro</a>              |
| 25 | <b>L.S.V.S.A. Timiș</b>     | Timisoara, Str. Surorile Martir Caceu Nr.4, Jud. Timiș                 | 0256-204911<br><a href="mailto:office-timis@ansv.ro">office-timis@ansv.ro</a>                          |
| 26 | <b>L.S.V.S.A. Vaslui</b>    | Bârlad, str. Trestiana, nr. 2, cod 731030, Jud.Vaslui                  | 0235-421121,<br>0235-421413<br><a href="mailto:office-vaslui@ansv.ro">office-vaslui@ansv.ro</a>        |
| 27 | <b>L.S.V.S.A. Vrancea</b>   | Focsani,B-dul Brailei, nr. 121 bis, cod 620122, Jud.Vrancea            | 0237-215561,<br>0237-232727<br><a href="mailto:office-vrancea@ansv.ro">office-vrancea@ansv.ro</a>      |

All Laboratories have to use the methods of the diagnostic presented at the point 1.4.

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the serotype according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

#### **1.4 Methods used in the examination of the samples in the framework of the programme**

Samples harvest by operators and samples harvest as official controls are prepared and tested in accordance with the requirements of the Commission Regulation (EC) No 200/2010, using the method recommended by the Community Reference Laboratory for *Salmonella* in Bilthoven, Netherlands. The method is described in the current version of ISO 6579:/A1:2007): 'Detection of *Salmonella* spp. in animal faeces and in samples of the primary production stage'. A semi-solid medium (modified semi-solid Rappaport-Vassiladis medium, MSRV) is used as the single selective enrichment medium. At least one isolate from each positive

sample shall be serotyped according to the Kaufmann-White scheme. In general the sensitivity of a panel of 10 antimicrobials will be determined.

### **1.5 Official controls**

Three samples will be taken under the control of the Competent Authority for Regulation 2160/2003 from each breeding flock during production of eggs for hatching as specified in 2.1.2.2 of Annex to Commission Regulation (EC) No 200/2010.

The use of antimicrobials (as defined in Regulation (EC) No 1177/2006) will be checked when the official sample is taken. If the flock is under antimicrobial medication for animal health or animal welfare reasons, the flock will be sampled again after the withdrawal period of the product specified in the Marketing Authorization. Flock owners are required to keep records of antimicrobial used and to make these records available to the competent authorities.

The records of samples harvest by the operator will be made available for inspection to the Competent Authority or its agent and will provide details of the identity of the flock sampled, date of sample, slaughter date, type of sample, laboratory carrying out the examination, and the result. The number of flocks on the holding and the number of birds present will be recorded.

Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs requires poultry abattoirs to undertake microbiological testing for *Salmonella* on 5 samples a week (each sample is 3 neck skins). Establishments producing minced meat, meat preparations and mechanically separated meat must also undertake weekly testing for *Salmonella*.

#### **Official controls at other stages of the food chain.**

Under the terms of the EC Feed Hygiene **Regulation 183/2005** provisions, feed businesses operators must be approved or registered by the Local Authority. Approvals/registrations are issued for the producers of compound feeds, feed materials, feed additives and premixtures. Approval requires a prior-inspection visit by the Local Authority to ensure that the establishments are in conformity with the required standards. The registration is followed by placing of premises on the list of feed business operators. The competent authority performs checks according to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order.

## **1.6 Measures taken by the Competent Authorities with regard to animals or products in which the presence of *Salmonella* spp. have been detected**

When a breeding flock of *Gallus gallus* is suspected of being infected with *Salmonella* Enteritidis or *Salmonella* Typhimurium the flock is placed under official control by the Competent Authority. This applies to breeding flocks from day old through to end of production. If the flock is in the laying phase no further eggs may be sent for hatching and no birds or hatching eggs may leave the holding, except under licence issued by the Competent Authority. When infection with *Salmonella* Enteritidis or *Salmonella* Typhimurium has been confirmed, the owner is required to have the birds slaughtered in accordance with Community legislation on food hygiene. Infection with *S. Typhimurium* and *S. Enteritidis* will be confirmed by samples taken by the Competent Authority as detailed in with Annex 2.2.2.2 of Regulation (EC) 200/2010.

The owner or person responsible for the flock is required to clean and disinfect the building where the infected birds were kept, and provide evidence to the Competent Authority that the cleaning and disinfection has been satisfactory. Bacteriological test to verify the efficiency of disinfection of poultry houses after depopulation of a *Salmonella*-positive flock will be carried out by the Competent Authority. Re-stocking may not take place until the cleaning and disinfection has been carried out properly.

Hatching eggs present in the hatchery from the time the flock was suspected to be infected are removed and destroyed in accordance with Regulation (EC) No 1774/2002.

When a breeding flock is suspected of being infected with *S. Hadar*, *S. Infantis*, or *S. Virchow*, the operator/owner of the flock will be required to draw up a plan in consultation with his/her veterinarian and the Competent Authority for the monitoring and control of the infection. The plan should include an epidemiological investigation, a review of biosecurity measures, any additional monitoring procedures, and measures to be taken to reduce or eliminate the infection. As appropriate the flock may be placed under official control.

The operator/owner in consultation with his/her veterinarian may consider vaccination of the flock against *Salmonella* with a product which has a marketing authorization in line with the requirements of Commission Regulation (EC) No **1177/2006** implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards requirements for the use of specific control methods in the framework of the national programmes for the control of salmonella in poultry.

Vaccination may only be used as a preventative measure; it is not an alternative to the requirements in Annex II C of Commission Regulation (EC) No 2160/2003 for the use of specific control methods in the framework of the national programmes for the control of *Salmonella*. Antimicrobial treatment may not be used for the control of *Salmonella* in the national control programme except within

the limits set by Commission Regulation **(EC) No 1177/2006** implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards requirements for the use of specific control methods in the framework of the national programmes for the control of salmonella in poultry.

Isolates of *Salmonella Enteritidis* and *Salmonella Typhimurium* will be examined to determine if they are vaccine strains according to the manufacturer's protocol. If vaccine strains are confirmed in samples the flock will not be classed as positive for the purposes of establishing the progress towards the target.

In exceptional cases when the competent authority has reason to question the results of the testing (such as false positive or false negative results), it may decide to repeat the testing in accordance with 2.2.2.2 point (b) from the provision Annex of Regulation 200/2010.

Operators will be invited to contact their veterinary for advice on how to reduce or eliminate the *Salmonella*. Advice will include recommendations on management, cleaning and disinfection, pest control, biosecurity, monitoring, and the potential use of vaccines.

### **1.7 Relevant national legislation.**

- N.S.V.F.S.A President **Order no. 34** to approve the sanitary veterinary norm regarding to monitoring zoonoses and zoonotic agents

- Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order

- N.S.V.F.S.A President **Order no. 205** for the approval of national reference laboratories and duties

- N.S.V.F.S.A President **Order no. 160** for the approval of the sanitary veterinary norm regarding the community reference laboratory for the zoonoses epidemiology and Salmonella and the national reference laboratory for Salmonella

### **1.8 Financial assistance provided to food and feed businesses operators in the context of the National Control Programme**

In breeding hens flocks of *Gallus gallus* financial assistance is provided in the context of the control programme. This is set up by the **Government Decision no. 1214/2009** regarding the methodology for determining and paying compensation to be paid to owners of slaughtered animals, killed or otherwise affected by the rapid liquidation of outbreaks of animal diseases.

The amount to compensate the animal owners in the context SNCP is determined in accordance with the provision of Government Decision No. 1214/2009 and is established by an evaluation committee consisting of:

- a) the representative of the county sanitary veterinary and food safety directorate (local CA)
- b) local representative of the Ministry of Agriculture and Rural Development with responsibilities in determining the genetic value of the animal
- c) the representative of the decision local unit from the local disease control center, designated by the Prefect,
- d) the local mayor or the person designated by him.

Convening of the evaluation committee is made by the Prefect at the written request of the county sanitary veterinary and food safety directorate (local CA).

On the proposal of the local county sanitary veterinary and food safety directorate the compensation committee considers and approves the replacement value of the animals according to the genetic value of animal, zootechnical value, sex, age, weight, physiological status, category production, at market price at the time when the liquidation of animals from the outbreak disease took place, and the average unit value based on the total amount of compensation for animals or products concerned according to Art. 4 of Regulation (EC) no. 349/2005 of the Commission, with the following amendments.

Romania applied to the Commission for co-financing for certain aspects of the control programme within the terms of Council Decision **2009/470/EC** on expenditure in the veterinary field.

## **2. Food and Feed businesses covered by the programme.**

The programme for the control of *Salmonella* in breeding hens is part of the controls along the whole food chain.

### **2.1 The structure of the production of the given species and products thereof.**

The structure of the breeding hens flocks and holdings in Romania at the beginning of 2012 is presented below:

| Region (County+NUTS CODE) | Type of flocks       | Number of flocks | Number of animals | Number of holdings |
|---------------------------|----------------------|------------------|-------------------|--------------------|
| <b>Total</b>              | <b>Breeding hens</b> | <b>451</b>       | <b>2374102</b>    | <b>41</b>          |

In Romania in 2011 there were 41 holdings with a total number of 451 flocks.

### **2.2 The structure of the production of food**

At the end of the breeding flock production period the birds are slaughtered and may go for human consumption. In accordance with Regulation 853/2004

provisions, poultry meat intended for human consumption must be obtained in approved slaughterhouses. There are **40** approved slaughterhouses **in Romania**.

Poultry feed is supplied to farms by a small number of manufacturers. The major manufacturers of poultry feed operate to assurance schemes, apply HACCP principles and monitor for Salmonella.

### **2.3 Relevant guides for good animal husbandry practices or other guidelines**

Some of the relevant guides for good animal husbandry practices are presented at the below link:

[http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella\\_12404ro.pdf](http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella_12404ro.pdf)

[http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010\\_13247ro.pdf](http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010_13247ro.pdf)

[http://www.ansvsa.ro/documente/admin/community\\_guide\\_layers\\_hygiene\\_practice\\_pullet\\_egg\\_ro\\_17813ro.pdf](http://www.ansvsa.ro/documente/admin/community_guide_layers_hygiene_practice_pullet_egg_ro_17813ro.pdf)

Other relevant guides for good animal husbandry practices and other guidelines are under constructions.

All farm registered must have their individual plan for good farmer practices, which are subject to approval by the official veterinarian responsible for the control of the holding concerned

### **2.4 Routine veterinary supervision of farms**

The owner is responsible for the health and welfare of the poultry on the holding, and for ensuring that a veterinarian is consulted on disease and welfare issues as appropriate. It is mandatory for each holding to have a contract with a private veterinarian who is responsible for veterinary care.

A veterinarian on behalf of the Competent Authority carries out inspections on farms for animal welfare reasons, to take samples for residues to administer and enforce the legislation on marketing of eggs and to check medicine records. Also a veterinarian on behalf of the Competent Authority visit the farms and take official samples in the framework of Salmonella NCP according with the legislation in force.

It is mandatory for each county sanitary veterinary and food safety directorate (local CA) to report to the NSVFSA every month the number of samples and results of these tests for each flock. Also the Salmonella NRL has the obligation to notify immediately NSVFSA and CSVFSD each positive sample for the relevant Salmonella.

## **2.5 Registration of farms**

All commercial poultry holdings in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centers / holdings of origin and means of transport for health and animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products. . All commercial holdings are entered into our national data base.

All poultry breeding flocks of more than 250 birds are included in the Salmonella National Control Programme.

## **2.6 Record-keeping at farms**

a. All breeding hens flock operators are required to keep records of veterinary medicines use, including vaccines, which must be available for inspection.

b. Records relating to movement of flocks on to/off the holding must be kept.

c. Records giving details of sampling for *Salmonella* and results will be kept either at the holding or be readily available.

## **2.7 Documents to accompany animals when dispatched**

Operators wishing to export birds or hatching eggs to another EU Member State (or certain third countries) must comply with **EU Directive 2009/158/EC** and ensure that the consignment is accompanied by a completed and signed Intra-trade Animal Health Certificate (ITAHC) for poultry breeding and production. This must be completed and signed by the Official Veterinarian as well as the operator to confirm compliance with the relevant articles of Directive.

The ITAHC will also require the reference number of the operator's poultry health certificate. The ITAHC is amended to include the results of the last test for Salmonella as is required in the Commission Regulation (EC) 2160/2003 Art 9.1 prior to any dispatching of the live animals, or hatching eggs, from the food business operators place of origin. The date and the result of testing are included in the relevant health certificate provided in the Community legislation.

## **2.8 Other relevant measures to ensure the traceability of animals.**

It is require to the operators of poultry flocks to keep records of poultry or hatching eggs entering or leaving the premises. The records must contain

information on the number, date, and origin or destination. These records must be available to the Competent Authority for inspection.

All official veterinary health certificates issued for the intra community trade of poultry and hatching eggs are recorded on the Trade Control and Expert System (TRACES). This system allows tracking of exports of live animals and hatching eggs accompanied by veterinary health certification. TRACES generate ITAHCs issued for intra-Community movements. TRACES are an internet-based service which is owned and maintained by the Commission.

## **PART B**

### **1. Identification of the programme**

|                             |  |
|-----------------------------|--|
| Member State:               | <b>Romania</b>   |
| Disease:                    | <b>Salmonella Enteritidis, S. Typhimurium, S. infantis, S. Hadar, S. Virchow in Breeding hens (GALLUS gallus)</b>  |
| Year of implementation:     | <b>2013</b>  |
| Reference of this document: | <b>National Sanitary Veterinary and Food Safety Authority<br/>No. /</b>  |
| Contact :                   | <b>Dr. Nicolae LAZĂR,<br/>Director,<br/>phone: +4 021 315 78 75;<br/>fax: +4021 31249 67,<br/>e-mail: <a href="mailto:lazar.nicolae@ansvsa.ro">lazar.nicolae@ansvsa.ro</a></b> |

**Date sent to the Commission : 27.04.2012**



*Salmonella* has been recognized as an important zoonotic pathogen for many years. *Salmonella Enteritidis* and *Salmonella Typhimurium* have accounted for the majority of cases of human salmonellosis and have consistently been the most commonly implicated pathogens in general outbreaks of food-borne disease.

A National Control Programme of *Salmonella* was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 1003/2006. The national control programme for *Salmonella* in breeding flock was coming into effect in January 2007.

In line with Regulation EC No 1003/2005 since 01 January 2007 the programme includes the control of *Salmonella Enteritidis*, *Salmonella Typhimurium* S. Hadar, S. Infantis and S. Virchow. As a result of the control programme the number of *Salmonella Enteritidis* and *Salmonella Typhimurium* infected breeding flocks of *Gallus gallus* in Roumania is currently low. Of the other three *Salmonella* serovars, *Salmonella* Hadar, *Salmonella* Infantis and *Salmonella* Virchow, the occurrence is likewise at low levels. Breeding flocks which are confirmed to be infected with zoonotic *Salmonella* are compulsorily slaughtered.

All breeding flocks of 250 birds or more were included in the programme. Operators will be required to implement the sampling programme in Annex IIB of EC Regulation 2160/2003. Samples for the detection of *Salmonella* will be taken from *Gallus gallus* day-old chicks to be used for breeding, when the birds are approximately 4 weeks of age, and approximately 2 weeks before the birds come into lay. During the production phase of laying eggs for hatching the flocks will be sampled every two weeks by the operator on the holding to verify the achievement of the target in adult breeding flocks. Sampling to verify the achievement of the target was as detailed in the Annex to Commission Regulation (EC) No 1003/2005. Samples were submitted to a laboratory authorized by the Competent Authority and which applies quality assurance systems that conform to the requirements of the current EN/ISO standard.

During 2011, a totally of 396 breeding flocks were tested for *Salmonella* infection from wich a no of 1535 were tested in official control and there were no flocks positive for *Salmonella Typhimurium*, including monophasic ST and *Salmonella Enteritidis* . The prevalence for the target serotypes in breeding flock in 2011 was 0%, which is low and below the Community target.

## The incidence of various serotypes of Salmonella in poultry, in 2011

| NO.          | SEROTYPE                          | NO. OF STRAINS in Gallus gallus |
|--------------|-----------------------------------|---------------------------------|
| 1            | S.Agona                           | 1                               |
| 2            | S.Amsterdam                       | 4                               |
| 3            | S.Enteritidis                     | 7                               |
| 4            | S.Glostrup                        | 1                               |
| 5            | S.Infantis                        | 3                               |
| 6            | S.Kentucky                        | 4                               |
| 7            | S.Mbandaka                        | 8                               |
| 8            | S.Senftenberg                     | 3                               |
| 9            | S.Tennessee                       | 2                               |
| 10           | S.Thompson                        | 7                               |
| 11           | S.Typhimurium (vaccineted strain) | 2                               |
| <b>TOTAL</b> |                                   | <b>42</b>                       |

It is a statutory requirement for all laboratories which isolate *Salmonella* from a flock of chickens or its environment to report the finding and supply the isolate to the National Reference Laboratory (NRL) for *Salmonella*. The isolates are serotyped, phage-typed (Cantacuzino Institute), where appropriate, and tested for antimicrobial sensitivity by the NRL. This information is recorded and analyzed. The number of reports received depends on the level and sensitivity of monitoring which is undertaken by the producers. The reports provide useful information on the serovars which are most common in poultry, and indicate trends.

### **3. Description of the submitted programme**

#### **Objectives**

The main objective of our programme as referred to in Article 4(1) of Regulation (EC) No 2160/2003, for the reduction of *Salmonella* spp. in breeding flocks of *Gallus gallus* shall be a reduction to 1 % or less of the maximum percentage of adult breeding flocks of *Gallus gallus* remaining positive for *Salmonella enteritidis*, *Salmonella infantis*, *Salmonella hadar*, *Salmonella typhimurium* and *Salmonella virchow* (the relevant *Salmonella* serotypes), according to Regulation (EC) No 2160/2003 and Regulation (EC) 200/2010.

The subsidiary objectives of the programme include further reduction of the incidence of infected table eggs, the reduction of incidence of broiler meat and eventually the reduction of the occurrence of human salmonellosis and a

consequent reduction in suffering, mortality and health service costs. This programme is in accordance to Council Directive 1999/74/EC, Commission Decision 2008/425/EC, Council Decision 92/65/ EEC, Council Decision 2009/470/ EEC, Regulation 2160/2003/EC, Regulation 200/2010/EC and Regulation 1177/2006/ EC.

### Target animal population

The National Control Programme for Salmonella in breeding flocks will be held in all holdings of breeding hens consisting of at least 250 poultry of Gallus gallus .

The number of holdings and flocks in breeding hens sector in 2011 are shown **in the table below.**

| Region<br>(County+NUTS<br>CODE) | Type of<br>flocks | Number of<br>flocks | Number of animals |
|---------------------------------|-------------------|---------------------|-------------------|
| RO121-AB                        | Breeding<br>hens  | 18                  | 110600            |
| RO311-AG                        | Breeding<br>hens  | 24                  | 150734            |
| RO211-BC                        | Breeding<br>hens  | 36                  | 308872            |
| RO111-BH                        | Breeding<br>hens  | 10                  | 85350             |
| RO221-BR                        | Breeding<br>hens  | 7                   | 47626             |
| RO122-BV                        | Breeding<br>hens  | 46                  | 159641            |
| RO222-BZ                        | Breeding<br>hens  | 24                  | 143032            |
| RO312-CL                        | breeding hens     | 36                  | 201333            |
| RO223-CT                        | Breeding<br>hens  | 30                  | 216776            |
| RO123-CV                        | Breeding<br>hens  | 2                   | 7484              |
| RO313-DB                        | Breeding<br>hens  | 61                  | 250000            |
| RO224-GL                        | Breeding<br>hens  | 1                   | 4120              |
| RO423-HD                        | Breeding<br>hens  | 8                   | 46020             |
| RO315-IL                        | Breeding<br>hens  | 56                  | 205123            |
| RO125-MS                        | Breeding<br>hens  | 2                   | 11000             |
| RO316-PH                        | Breeding<br>hens  | 41                  | 196828            |
| RO115-SM                        | Breeding<br>hens  | 30                  | 150000            |
| RO216-VS                        | Breeding          | 12                  | 32733             |

|              |                          |            |                |
|--------------|--------------------------|------------|----------------|
|              | hens                     |            |                |
| <b>Total</b> | <b>Breeding<br/>hens</b> | <b>451</b> | <b>2374102</b> |

**Table . Number of flocks in the breeding hens sector**

**Sampling programmes**

The National Salmonella Control Programme encompasses the following serovars of zoonotic Salmonella: Salmonella enteritidis, Salmonella typhimurium Salmonella infantis, Salmonella virchow and Salmonella hadar .

The sampling programme will be in accordance to Regulation 2160/2003 EC and Regulation 200/2010 EC. We have also taken into account Regulation 1177/2006 EC.

#### 4.1. Summary of measures under the programme

Year :2013

- Control
- Testing
- Slaughter of positive animals
- Killing of positive animals
- Vaccination
- Treatment of animal products
- Disposal of products
- Monitoring or surveillance
- Other measures

#### 4.2. Designation of the central authority charged with supervising and coordinating the departments responsible for implementing the programme:

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in breeding hens flocks of *Gallus gallus* is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**

The departments responsible for implementing the National Control Programme of Salmonella in breeding hens are:

- for implementing at central level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary and Food Chain Safety General Directorate** ;
- In respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the checking of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**.
- Concerning the slaughter of flocks, the **Traceability and Food Safety of Animal Origin Directorate** is responsible for public health protection, through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.
- In respect of EC Regulation No **183/2005** on feed hygiene at the central level is **The Directorate for Technical Coordination of Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**

- At county level for implementing the Regulation (EC) No **2160/2003**, **882/2004** and **183/2005** there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the National Reference Laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

#### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

##### **Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România

Tel: 0374.322.013 / 0374.322.000

Fax: 0214.113.394

E-mail: [office@idah.ro](mailto:office@idah.ro)

- **H.I.P.V.H.** (Hygiene Institute for Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

#### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED**

##### **Hygiene Institute for Sanitary Veterinary Public Health**

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201

Tel: 021.252.46.51 Fax: 021.252.00.61 E-mail: [iispv @ iispv.ro](mailto:iispv@iispv.ro)

The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL in January 2012 is show in the link below:

[http://www.ansvsa.ro/documente/admin/lista%20laboratoarelor%202011%20-%2028.01.2011\\_11043ro.pdf](http://www.ansvsa.ro/documente/admin/lista%20laboratoarelor%202011%20-%2028.01.2011_11043ro.pdf)

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the serotype according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

### **1. N.S.V.F.S.A.**

- proposal of the plan of disease control.
- elaborate and submit to the EC the National Control Programme
- evaluate, coordinate the implementation of the National Control Programme in Romania and propose the update of this programme if is necessary (depending of the national situation)
- adoption of measures based on the disease situation in Romania
- submission of reports to the E.C.
- training the specialists from C.S.V.F.S.D

### **2. S.V.F.S.D.**(Sanitary Veterinary and Food Safety Direction).

- Coordination of the programme at the county level
- Official sampling is performed by the official veterinarian
- Propose the compensation costs to the Commission which will evaluate compensation documents.

**3. I.D.A.H.** Institute of Diagnosis and Animal Health is the National Reference Laboratory concerning animal health and was also designated as NRL for Salmonella in animals.

Responsibilities and tasks of the National Reference Laboratory for Salmonella (**I.D.A.H.**), pursuant to Directive 2003/99/EC and Regulation (EC) No 2160/2003, according to provision of N.S.V.F.S.A.. President Order no.160/2006 and 205/2007 are:

#### **1. General duties**

- (a) To collaborate with the Community reference laboratory in their area of competence.
- (b) To coordinate, as appropriate, the activities of laboratories responsible for the analysis of samples in accordance with, in particular, Articles 4, 5 and 7 of Directive 2003/99/EC.
- (c) To coordinate the activities of laboratories responsible for the analysis of samples in accordance with Article 12(1) of Regulation (EC) No 2160/2003/EC.
- (d) Where appropriate, to organise comparative tests between the laboratories referred to under (b) and (c) and to assure an appropriate follow-up of such comparative testing.
- (e) To ensure the dissemination to the competent authority and to the laboratories referred to under (b) and (c), of the information that the Community reference laboratory supplies.

- (f) To provide scientific and technical assistance to the National Sanitary Veterinary and Food Safety Authority in their area of competence.
- (g) Characterize the pathogen isolates, genetic typing of this agents
- (h) Keep in maxim security conditions, the isolates
- (i) Give to the C.E., CRL, OMS, and national reference Laboratory of other member states, with the accord of N.S.V.F.S.A. all the information required.
- (j) train of the specialists from the government and private laboratory.

## **2. Specific functions and duties**

- (a) To participate, as appropriate in the monitoring and control programme for Salmonella and related anti-microbial resistance pursuant to Directive 2003/99/EC and in the analysis and testing of *Salmonella* pursuant to Regulation (EC) No 2160/2003.
- (b) To inform, as appropriate, the Community reference laboratory on aspects related to Salmonella vaccine strains and other specific control methods.
- (d) To gather data and information on the activities developed and methods used in relevant laboratories and to inform the Community reference laboratory thereof.
- (e) To monitor the epidemiological evolution of salmonella in Roumania.

## **5. S.V.F.S.L.**

There are 41 county Sanitary Veterinary and Food Safety Laboratories. Only 27 of them apply quality assurance systems that conform to the requirements of the current EN/ISO standard and are designated by National Reference Laboratory for Salmonella in animals to perform bacteriological examinations in the framework of the programme under the supervision of IDAH-NRL for Salmonella.

## **6. H.I.P.S.V.H.**

Hygiene Institute for Public Sanitary Veterinary Health is the National Reference Laboratory concerning the expertise for food products of animal origin and feed and it is also designed as the National Reference Laboratory for Salmonella in food and feed –public veterinary health.



### **4.3. Description and delimitation of the geographical and administrative areas in which the programme is to be implemented:**

The National Control Programme will be implemented throughout Romania, covering all the national territory and will cover all breeding flocks of *Gallus gallus* consisting of at least 250 breeding poultry.

The administrative boundaries are the boundaries of the country. Roumania is administrative divided in 42 counties. There are 42 County Sanitary Veterinary and Food Safety Directorates and 41 County Sanitary veterinary and food Safety Laboratories.

### **4.4 Measures implemented under the programme**

#### **Bio-Security Measures**

Bio-security is a combination of practices, which are intended to prevent the spread of disease-causing organisms within the poultry farm. Where these are performed in parallel with the sanitation and disinfection procedures, bio-security measures could eradicate or, at least, reduce the level of pathogens to values, at which no hazard of infection would be likely.

The bio-security measures in commercial poultry farms are in accordance to the N.S.V.F.S.A. President Order 147/2006 regarding the bio-security measures in commercial poultry farms and the movements of live poultry, products and poultry by-products.

#### **Bio-security measures on holdings:**

- Health status of poultry
- On entering to all houses on the farm must be located disinfection barrier
- Control of movement of people
- Transport hygiene
- Feed hygiene
- Water hygiene
- Rodent, insect and bird control
- Cleaning and disinfecting of buildings
- Recording of all events and operations

According to the provision of Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program approved every year by N.S.V.F.S.A. President Order, after each cleaning and disinfections the owner is oblige to take samples to verify the efficient of the disinfection. The official

veterinarian take sample to verify the efficiency of the disinfection only in case of a positive flock.

Hygiene measures on poultry farms are also assessed during visits for the collection of official samples and during general visits to premises for other purpose.

#### **4.4.1. Measures and terms of legislation as regards the registration of holdings:**

All commercial poultry flocks in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centers / holdings of origin and means of transport for health and animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products. Poultry holdings shall be registered and sanitary veterinary approved in order to be able to operate.

The legal framework for carrying out commercial activity with poultry and poultry products consists of:

- **Order** of the President of the National Sanitary Veterinary and Food Safety Authority no **144/2006** for the approval of the Sanitary veterinary norm on animal health conditions governing intra-Community trade in, and imports from third countries of, poultry and hatching eggs transposing Council Directive 90/539/EEC on animal health conditions governing intra-Community trade in, and imports from third countries of poultry and hatching eggs.

In order to be sanitary veterinary approved, poultry holdings shall satisfy, in accordance with the legislation in force, the following conditions:

- Appropriate facilities and operation;
- Application of the “Program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” approved by Order of the National Sanitary Veterinary and Food Safety Authority President .
- At least one inspection visit per year by the official veterinarian;
- Additional checks to verify the compliance of the establishment with the hygiene measures and the operation of the establishments.

Each poultry holding receives a distinct approval number, number that can be the same with the one given in compliance with the **Council Regulation no. 2782/75/EEC** on the production and marketing of eggs for hatching and of farmyard poultry chicks.

#### **4.4.2. Measures and terms of legislation as regards the identification of animals:** Not applicable.

#### **4.4.3. Measures and terms of legislation as regards the notification of the disease:**

**Commission Decision 176/2005** of 1 March 2005 laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2006/924/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2010/160/EU** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2008/755/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2009/847/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

The internal notification is made under the provision of **N.S.V.F.S.A. President ORDER no. 79 /2008** to approve the sanitary and veterinary norm on domestic notification and official declaration of animal diseases.

#### **4.4.4. Measures and terms of legislation as regards the measures in case of a positive result:**

Whenever a flock is found positive by own-check sampling in the framework of the SNCP in breeding hens this flock is considered a suspect flock and movement restriction and other relevant restrictive measures are mandatory imposed on this flock.

When a breeding flock of *Gallus gallus* is suspected of being infected with *Salmonella* Enteritidis or *Salmonella* Typhimurium the flock is placed under official control by the Competent Authority. The flock is suspected of being infected when *S. Enteritidis* or *S. Typhimurium* is isolated from a sample of faeces, boot swabs, tissue or organs from the birds in the flock, or from dust in the environment. This applies to breeding flocks from day old through to end of production. If the flock is in the laying phase no further eggs may be sent for hatching and no birds or hatching eggs may leave the holding, except under license issued by the Competent Authority. If a flock is suspected of being infected with *Salmonella* Enteritidis, *Salmonella* Typhimurium, *Salmonella* Hadar, *Salmonella* Infantis or *Salmonella* Virchow, confirmatory samples consisting of a repeat of five pairs of boot swabs per flock will be taken (or two pooled 150g samples of faeces in the case of caged flocks) by a government inspector for confirmation. If *S. Enteritidis*, or *S. Typhimurium* is confirmed in these samples the flock will be slaughtered,

similar to the procedures which are currently in place. When infection with *Salmonella* Enteritidis or *Salmonella* Typhimurium has been confirmed, the owner is required to have the birds slaughtered in accordance with Community legislation on food hygiene. Infection with *S. Typhimurium* and *S. Enteritidis* will be confirmed by samples taken by the Competent Authority as detailed in with Annex 2.2.2.2 of Regulation (EC) 200/2010.

Also additional samples can be collected for the possible testing of antimicrobials or bacterial growth inhibitors as follows: birds shall be taken at random from within each poultry house of birds on the holding, normally up to five birds per house, unless the competent authority deems it necessary to sample a higher number of birds.

If the source of infection is not confirmed, antimicrobial testing shall be carried out or new bacteriological testing for the presence of the relevant *Salmonella* serotypes shall be carried out on the breeding flock or their progeny before trade restrictions are lifted.

A breeding flock shall be considered positive for the purpose of ascertaining the achievement of the Union target:

- when the presence of the relevant *Salmonella* serotypes (other than vaccine strains) has been detected in one or more samples taken in the flock, even if the relevant *Salmonella* serotypes is only detected in the dust sample, or
- when the confirmatory sampling as part of official controls in accordance with point 2.2.2.2(b) of Regulation 200/2010 does not confirm the detection of relevant *Salmonella* serotypes but antimicrobials or bacterial growth inhibitors have been detected in the flock.

This rule shall not apply in exceptional cases described in point 2.2.2.2(c) of Regulation 200/2010 where the initial *Salmonella* positive result from sampling at the initiative of the food business operator has not been confirmed by the sampling as part of official controls.

A positive breeding flock shall only be counted once regardless of how often the relevant *Salmonella* serotypes has been detected in this flock during the production period or whether the sampling was carried out at the initiative of the food business operator or by the competent authority. However, if sampling during the production period is spread over two calendar years, the result of each year shall be reported separately.

The owner or person responsible for the flock is required to clean and disinfect the building where the infected birds were kept, and provide evidence to the Competent Authority that the cleaning and disinfection has been satisfactory. Bacteriological test to verify the efficiency of disinfection of poultry houses after depopulation of a *Salmonella*-positive flock will be carried out by the Competent Authority. Re-stocking may not take place until the cleaning and disinfection has been carried out properly.

Hatching eggs present in the hatchery from the time the flock was suspected to be infected are removed and destroyed in accordance with Regulation (EC) No 1774/2002.

When a breeding flock is suspected of being infected with *S. Hadar*, *S. Infantis*, or *S. Virchow*, the operator/owner of the flock will be required to draw up a plan in consultation with his/her veterinarian and the Competent Authority for the monitoring and control of the infection. The plan should include an epidemiological investigation, a review of biosecurity measures, any additional monitoring procedures, and measures to be taken to reduce or eliminate the infection. As appropriate the flock may be placed under official control.

The operator/owner in consultation with his/her veterinarian may consider vaccination of the flock against *Salmonella* with a product which has a marketing authorisation in line with the requirements of Commission Regulation (EC) No **1177/2006** implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards requirements for the use of specific control methods in the framework of the national programmes for the control of salmonella in poultry.

Vaccination may only be used as a preventative measure; it is not an alternative to the requirements in Annex II C of Commission Regulation (EC) No 2160/2003 for the use of specific control methods in the framework of the national programmes for the control of *Salmonella*. 1.8.6 Antimicrobial treatment may not be used for the control of *Salmonella* in the national control programme except within the limits set by Commission Regulation **(EC) No 1177/2006** implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards requirements for the use of specific control methods in the framework of the national programmes for the control of salmonella in poultry.

Isolates of *Salmonella Enteritidis* and *Salmonella Typhimurium* will be examined to determine if they are vaccine strains according to the manufacturer's protocol. If vaccine strains are confirmed in samples the flock will not be classed as positive for the purposes of establishing the progress towards the target.

In exceptional cases when the competent authority has reason to question the results of the testing (such as false positive or false negative results), it may decide to repeat the testing in accordance with 2.2.2.2 point (b) from the provision Annex of Regulation 200/2010.

Operators will be invited to contact their veterinary for advice on how to reduce or eliminate the *Salmonella*. Advice will include recommendations on management, cleaning and disinfection, pest control, biosecurity, monitoring, and the potential use of vaccines.

In case of suspicion or confirmation of ***Salmonella enteritidis*** or ***Salmonella typhimurium*** the NRL shall notify immediately the N.SV.F.S. and local C.S.V.F.S.D..

## **Arrangements for implementing the specific requirements concerning flocks of breeding hens set out in Regulation 2160/2003 Annex II C.**

### ***Specific requirements concerning breeding flocks of Gallus gallus***

1. Starting with 1 January 2010 the measures laid down below in points 3 to 5 must be taken whenever the analysis of samples carried out in accordance with point 2.2.2.1 and 2.2.2.2 of Annex of the Regulation 200/2010 indicates the presence of *Salmonella enteritidis* or *Salmonella typhimurium* in a breeding flock of *Gallus gallus* in the circumstances set out in point 2.

2. (a) If the competent authority has approved the method of analysis used for samples taken in accordance with part B of the Regulation 2160/2003/EC , it may require that the measures laid down in points 3 to 5 be taken when such analysis detects the presence of *Salmonella enteritidis* or *Salmonella typhimurium*.

(b) Otherwise, the measures laid down in points 3 to 5 must be taken whenever the competent authority confirms a suspicion of the presence of *Salmonella enteritidis* or *Salmonella typhimurium* arising from the analysis of samples carried out in accordance with part B of the Regulation 2160/2003/EC.

#### **3. Non-incubated eggs from the flock must be destroyed.**

However, such eggs may be used for human consumption if they are treated in a manner that guarantees the elimination of *Salmonella enteritidis* and *Salmonella typhimurium* in accordance with Community legislation on food hygiene.

4. **All birds**, including day-old chicks, **in the flock must be slaughtered or destroyed so as to reduce as much as possible the risk of spreading Salmonella.** Slaughtering must be carried out in accordance with Community legislation on food hygiene. Products derived from such birds may be placed on the market for human consumption in accordance with Community legislation on food hygiene and, once applicable, part E of the Regulation 2160/2003. If not destined for human consumption, such products must be used or disposed of in accordance with Regulation (EC) No 1774/2002 of the European Parliament and of the Council, laying down health rules concerning animal by-products not intended for human consumption.

5. Where **eggs** for hatching from flocks in which *Salmonella enteritidis* or *Salmonella typhimurium* is present are still **present in a hatchery, they must be destroyed or treated** in accordance with Regulation (EC) No 1774/2002

### **Control of the use of feed antibiotics by official sampling**

According to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in

food safety field approved every year by N.S.V.F.S.A. President Order, feeding stuffs intended for poultry nutrition are checked in view to avoid the contamination with *Salmonella* spp. Also, in conformity with the same legislation the feed stuffs are checked in view to detect the use of antibiotics.

Residues examination is performed according to the Roumanian annual plan for examination for residues in live animals and animal origin products .

For broiler, hens, turkeys, other poultry a sample consists on one or more animals depending on the requirements of the analytical methods.

For each category of poultry considered, the minimum number of samples to be taken each year must be at least equal to one per 200 tones of annual production, with a minimum of 100 samples for each group of substances if the annual production of the category of birds considered is over 5 000 tones.

The following breakdown must be respected:

**Group A:** 50 % of the total samples. The equivalent of one fifth of these samples must be taken at farm level. Each sub-group of Group A must be checked each year using a minimum of 5 % of the total number of samples to be collected for Group A. The balance will be allocated according to the experience and background information of Roumania.

**Group B:** 50 % of the total samples, 30 % of samples must be checked for Group B 1 substances (antibiotics and sulfamides)

#### **4.4.6. Control procedures and in particular rules on the movement of animals liable to be affected or contaminated by a given disease and the regular inspection of the holdings or areas concerned.**

According to the provisions of N.S.V.F.S.A. President Order 147/2006, Regulation 2160/2003/EC, the following measures are to be adopted in order to prevent the dissemination of *Salmonella* spp. into commercial holdings. Animals from infected flocks belonging to commercial holdings are to be kept isolated and special conditions apply for removal of these animals. No bird may leave the house concerned unless the competent authority has authorized the slaughter or/and destruction under supervision of slaughter in a slaughterhouse designated by the competent authority. All the birds in the house must be slaughtered in accordance with the provisions of the **REGULATION (EC) No. 853/2004 laying down specific hygiene rules for food of animal origin in order** to reduce as much as possible the risk of spreading salmonella.

#### **4.4.7. Measures and terms of legislation as regards the control (testing, vaccination ...) of the disease.**

The legal basis is the provisions of the **Regulation 2160/2003/EC, Regulation 200/2010/EC and Regulation 1177/2006 EC.**

The vaccination against Salmonella in breeding hens it is not mandatory in Romania, but the CCA strongly recommender this. In case the owner of the flock wants to vaccinate against Salmonella he will send his vaccination programme to the local CA. If the vaccination programme is in accordance with the provision of Regulation 1177/2006 EC , the local CA approve it . Only the flocks which have an approved vaccination programme can vaccinate against Salmonella.

Measures and applicable legislation as regard the control of the disease are the measures outlined in point 4.4.4 above.

According to the provision of Regulation (EC) no. 1177/2006 CE in Romania the antimicrobials use for Salmonella control is forbidden.

A national *Salmonella* control programme was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 200/2010. The national control plan for *Salmonella* in breeding flocks come into effect in January 2007.

***The NCP defines a flock as a single group or multiple groups of chickens which share the same production unit (i.e. using the same air-space or range area). Where housing systems are not typical, the situation is likely to be assessed on a case by case basis. Multiple groups of chickens which have ‘beak-to-beak’ contact (inside or outside the house) are likely to be treated as a single flock.***

**Operators** will be required to implement the sampling programme in the Annex to EC Regulation 200/2010 (**self-control sampling**). For convenience the ‘Sampling protocol’ is repeated in **THE SAMPLING PROTOCOL**, showed below.

## **SAMPLING PROTOCOL.**

For each flock\*

### Sampling at the initiative of the food business operator (self-control sampling).

Sampling shall primarily consist of faecal samples and shall aim to detect a 1 % within flock prevalence, with a 95 % confidence limit. To that effect, the samples shall comprise one of the following:

(a) Pooled faeces made up of separate samples of fresh faeces each weighing not less than 1 g taken at random from a number of sites in the poultry house in which the breeding flock is kept, or where the breeding flock has free access to more than one poultry house on a particular holding, from each group of houses on the holding in which the breeding flock is kept. Faeces may be pooled for analysis up to a minimum of two pools.

The number of sites from which separate faeces samples are to be taken in order to make a pooled sample shall be as follows:

| Number of birds kept in the breeding flock | Number of faeces samples to be taken in the breeding flock |
|--|--|
| 250-349                                    | 200  |



|               |     |
|---------------|-----|
| 350-449       | 220 |
| 450-799       | 250 |
| 800-999       | 260 |
| 1 000 or more | 300 |

(b) Boot swabs and/or dust samples:

Boot swabs used shall be sufficiently absorptive to soak up moisture.

The surface of the boot swab shall be moistened using appropriate diluents approved by the competent authority). The samples shall be taken while walking through the house using a route that produces representative samples for all parts of the poultry house or the respective sector. It shall include littered and slatted areas provided that slats are safe to walk on. All separate pens within a poultry house shall be included in the sampling. On completion of the sampling in the chosen sector, boot swabs must be removed carefully so as not to dis-lodge adherent material.

The samples shall consist of:

- (i) five pairs of boot swabs, representing each about 20 % of the area of the poultry house; the swabs may be pooled for analysis into a minimum of two pools; or
- (ii) at least one pair of boot swabs representing the whole area of the poultry house and an additional dust sample collected from multiple places throughout the poultry house from surfaces with visible presence of dust; one or several moistened fabric swab(s) of at least 900 cm<sup>2</sup> surface area in total must be used to collect the dust sample.

(c) In cage breeding flocks, sampling may consist of naturally mixed faeces from dropping belts, scrapers or deep pits, depending on the type of house. Two samples of at least 150 g shall be collected to be tested individually:

- (i) droppings belts beneath each tier of cages which are run regularly and discharged into an auger or conveyor system;
- (ii) droppings pit system in which deflectors beneath the cages scrape into a deep pit beneath the house;
- (iii) droppings pit system in a step-cage poultry house when cages are offset and faeces fall directly into the pit.

There are normally several stacks of cages within a house. Pooled faeces from each stack shall be represented in the overall pooled sample. Two pooled samples shall be taken from each breeding flock as described in the following third to sixth subparagraphs.

In systems where there are belts or scrapers, these shall be run on the day of the sampling before sampling is carried out.

In systems where there are deflectors beneath cages and scrapers, pooled faeces that have lodged on the scraper after it has been run, shall be collected.

In step-cage poultry house systems where there is no belt or scraper system it is necessary to collect pooled faeces from throughout the deep pit.

Droppings belt systems: pooled faecal material from the discharge ends of the belts shall be collected.

### **Official sampling**

Official routine sampling shall be carried out on three occasions during the production cycle:

- (a) within four weeks following moving to laying phase or laying unit;
- (b) towards the end of the laying phase, not earlier than eight weeks before the end of the production cycle;
- (c) at any time during the production cycle which is sufficiently distant in time from the sampling referred to in points (a) and (b)

Official confirmatory sampling following the detection of the relevant *Salmonella* serotypes from sampling at the hatchery shall be performed as described in „Sampling at the initiative of the food business operator (self-control sampling)”

Additional samples can be collected for the possible testing of antimicrobials or bacterial growth inhibitors as follows: birds shall be taken at random from within each poultry house of birds on the holding, normally up to five birds per house, unless the competent authority deems it necessary to sample a higher number of birds.

If the source of infection is not confirmed, antimicrobial testing shall be carried out or new bacteriological testing for the presence of the relevant *Salmonella* serotypes shall be carried out on the breeding flock or their progeny before trade restrictions are lifted.

If antimicrobials or bacterial growth inhibitors are detected, the *Salmonella* infection shall be considered as confirmed.

#### Official samples for suspicion of false results

In exceptional cases where the competent authority has reason to question the results of the testing (such as false positive or false negative results), it may decide to repeat the testing in accordance with „Official confirmatory sampling”.

The use of antimicrobials (as defined in Regulation (EC) No 1177/2006) will be checked when the official sample is taken. If the flock is under antimicrobial medication for animal health or animal welfare reasons, the flock will be sampled again after the withdrawal period of the product specified in the Marketing Authorization. Flock owners are required to keep records of antimicrobial used and to make these records available to the competent authorities.

The records of samples harvest by the operator will be made available for inspection to the Competent Authority or its agent and will provide details of the identity of the flock sampled, date of sample, slaughter date, type of sample, laboratory carrying out the examination, and the result. The number of flocks on the holding and the number of birds present will be recorded.

When an official routine sample is taken it may replace the sample required to be taken by the operator.

#### 4. Measures of the submitted programme

##### Transport and preparation of the samples:

Samples shall preferably be sent by express mail or courier to the laboratories referred to in Articles 11 and 12 of Regulation (EC) No 2160/2003, within 24 hours after collection. If not sent within 24 hours, they shall be stored refrigerated. Transportation can be at ambient temperature as long as excessive heat (over 25 °C) and exposure to sunlight are avoided. At the laboratory samples shall be kept refrigerated until examination, which shall be started within 48 hours following receipt and within 96 hours after sampling.

##### *Boot swabs and dust samples:*

(a) The pair(s) of boot/sock swabs and dust sample (fabric swab) shall be carefully unpacked to avoid dislodging adherent faecal material or loose dust material and placed in 225 ml of BPW which has been pre-warmed to room temperature.

(b) The boot/socks and fabric swab shall be fully submersed in BPW to provide sufficient free liquid around the sample for migration of *Salmonella* away from the sample and therefore more BPW may be added, if necessary.

Separate preparations must be made of the boot swabs and the fabric swab.

(c) Where five pairs of boot/sock swabs are pooled into two samples, each pooled sample must be placed in of 225 ml of BPW, or more if necessary, to fully submerge the sample and provide sufficient free liquid around the sample for migration of *Salmonella* away from the sample.

(d) Swirl to fully saturate the sample and continue the culture by using the detection method described in point 3.2.

##### *Other faecal material samples:*

(a) The faeces samples shall be pooled and thoroughly mixed and a 25 g sub-sample shall be collected for culture.

(b) The 25 g sub-sample shall be added to 225 ml of BPW which has been pre-warmed to room temperature.

(c) The culture of the sample shall be continued by using the detection method described below.

##### **Detection method**

The detection of the relevant *Salmonella* serotypes shall be carried out according to Amendment 1 of EN/ISO 6579-2002/Amd1:2007. 'Microbiology of food and animal feeding stuffs – Horizontal method for the detection of *Salmonella* spp. — Amendment 1: Annex D: Detection of *Salmonella* spp. in animal faeces and in environmental samples from the primary production stage'.

As regards the boot swabs samples, dust samples and other faecal material samples, the incubated BPW enrichment broth for future culture may be pooled. To do so, incubate both samples in BPW . Take 1 ml of incubated broth from each

sample and mix thoroughly, then take 0,1 ml of the mixture and inoculate the modified semi-solid Rappaport-Vassiliadis (MSRV) plates. The samples in BPW must not be shaken, swirled or otherwise agitated after incubation as this releases inhibitory particulates and reduces subsequent isolation in MSRV.

### **Serotyping**

At least one isolate from each positive sample shall be serotyped, following the Kaufmann-White scheme.

#### **Storage of strains:**

At least one isolated strain per house and per year shall be collected by the NRL and stored for future phagotyping or anti-microbial susceptibility testing, using the normal methods for culture collection, which must ensure integrity of the strains for a minimum of two years.

#### **Use of vaccines**

Live Salmonella vaccines shall not be used in the framework of national control programme where the manufacturer does not provide an appropriate method to distinguish bacteriological wild-type strains of salmonella from vaccine strains.

Live salmonella vaccines shall not be used in the framework of national control programmes in breeding hens during production unless the safety of the use has been demonstrated and they are authorized for such purpose in accordance with Directive 2001/82/EC.

#### **4.4.8. Measures and terms of legislation as regards the compensation for owners of slaughtered and killed animals:**

In breeding flocks of *Gallus gallus* financial assistance is provided in the context of the control programme. This is set up by the **Gouvernement Decision no. 1214/2009** regarding the methodology for determining and paying compensation to be paid to owners of slaughtered animals, killed or otherwise affected by the rapid liquidation of outbreaks of animal diseases.

#### **4.4.9. Information and assessment on bio-security measures management and infrastructure in place in the flocks/holdings involved**

This information was given in the point 4.4.

### **5.General description of the benefits:**

#### **Costs of the programme**

This programme is in accordance with Council Decision 2009/470/ EEC, Commission Decision 2008/425/EC and Council Decision 92/65/ EEC. The detailed financial costs for the national control programme of Salmonella in breeding hens are laid down in point 8.

**Benefits of the programme**

The overall aim of the Salmonella National Control Programme is to control the occurrence of Salmonella in the eggs sector on a very low level and thereby protect humans against infection with food-borne salmonellas. It has been known that poultry often harbor latent infections with Salmonella, which may pose a serious human health risk.

The anticipated benefits of this programme reducing the relevant *Salmonella* are the minimising of human health problems and a consequent reduction in suffering, mortality and health service costs.

**6. Data on the epidemiological evolution during the last year:**

## 6.1.Evolution of zoonotic salmonellosis

### 6.1.1 Data on evolution of zoonotic salmonellosis

Year: 2009

Situation on date: 31.12.2009

Animal species: breeding hens (Gallus gallus)

Disease: zoonotic salmonellosis

| Region<br>(County+NU<br>TS CODE) | Type of<br>flocks(b) | Total<br>number<br>of flocks<br>© | Total number of<br>animals | Total number<br>of flocks<br>under the<br>programme | Total number<br>of animals<br>under the<br>programme | Number of<br>flocks<br>checked(d) | Number of positive<br>flocks (e)                                     |                           | Number of flocks<br>depopulated                                     |                       | Total number of animals<br>slaughtered or destroyed         |                           | Quantity of the eggs<br>destroyed (number<br>or kg)                 |                           | Quantity of eggs channelled to egg<br>products (number or kg) |                    |
|----------------------------------|----------------------|-----------------------------------|----------------------------|---|--|-----------------------------------|--|---------------------------|---|-----------------------|---|---------------------------|---|---------------------------|---|--------------------|
|                                  |                      |                                   |                            |   |  |                                   | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) | Serotyp<br>es<br>targeted<br>in the<br>control<br>progra<br>mme (f) | Other<br>serotypes(g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>serotyp<br>es(g) | Serotyp<br>es<br>targeted<br>in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) | Serotypes targeted<br>in the control<br>programme (f)         | Other serotypes(g) |
| RO121-<br><b>AB</b>              | Breeding<br>hens     | 18                                | 115000                     | 18  | 115000   | 18                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO311-<br><b>AG</b>              | Breeding<br>hens     | 24                                | 144594                     | 24  | 144594   | 24                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO211-<br><b>BC</b>              | Breeding<br>hens     | 47                                | 602000                     | 47  | 602000   | 47                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO111-<br><b>BH</b>              | Breeding<br>hens     | 11                                | 67646                      | 11  | 67646  | 11                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO221-<br><b>BR</b>              | Breeding<br>hens     | 5                                 | 43889                      | 5   | 43889  | 5                                 | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO122-<br><b>BV</b>              | Breeding<br>hens     | 45                                | 234791                     | 45  | 234791   | 45                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO222-<br><b>BZ</b>              | Breeding<br>hens     | 24                                | 128216                     | 12  | 77984  | 12                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO312-<br><b>CL</b>              | breeding<br>hens     | 26                                | 118000                     | 26  | 118000   | 26                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |
| RO223-<br><b>CT</b>              | Breeding<br>hens     | 12                                | 62660                      | 12  | 62660  | 12                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                  |

|              |               |            |                |            |                |            |          |          |             |          |          |          |          |          |          |          |
|--------------|---------------|------------|----------------|------------|----------------|------------|----------|----------|-------------|----------|----------|----------|----------|----------|----------|----------|
| RO313-DB     | Breeding hens | 24         | 90495          | 24         | 90495          | 24         | 0        | 3        | 0           | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO314-GR     | Breeding hens | 8          | 99622          | 8          | 99622          | 8          | 1*       |          | 1835**      | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO423-HD     | Breeding hens | 18         | 73518          | 18         | 73518          | 18         | 0        | 0        | 0           | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO315-IL     | Breeding hens | 20         | 64401          | 20         | 64401          | 20         | 1***     | 0        | 0           | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO316-PH     | Breeding hens | 25         | 127838         | 25         | 127838         | 25         | 0        | 0        | 0           | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO115-SM     | Breeding hens | 18         | 143950         | 18         | 143950         | 18         | 0        | 0        | 0           | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO226-VN     | Breeding hens | 12         | 62370          | 12         | 62370          | 12         | 0        | 0        | 0           | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| <b>Total</b> |               | <b>337</b> | <b>2178990</b> | <b>325</b> | <b>2128758</b> | <b>325</b> | <b>2</b> | <b>0</b> | <b>1835</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> |

\*2 strains of Salmonella enteritidis were suspected in a flock.

\*\* birds were removed from the flock

\*\*\* Salmonella enteritidis in faeces samples was suspected, but at confirmation tests were not identified any type of Salmonella spp.

Year: 2010

Situation on date: 31.12.2010

Animal species: breeding hens (Gallus gallus)

Disease: zoonotic salmonellosis

| Region<br>(County+<br>NUTS<br>CODE) | Type of<br>flocks(b) | Total<br>number<br>of flocks<br>© | Total<br>number of<br>animals | Total<br>number of<br>flocks<br>under the<br>programme | Total<br>number of<br>animals<br>under the<br>programme | Number<br>of flocks<br>checked<br>(d) | Number of positive<br>flocks (e)                                |                           | Number of flocks<br>depopulated                                |                           | Total number of animals<br>slaughtered or destroyed            |                       | Quantity of the eggs<br>destroyed (number or kg)                |                           | Quantity of eggs<br>channelled to egg<br>products (number or<br>kg) |                           |
|-------------------------------------|----------------------|-----------------------------------|-------------------------------|--|---|---------------------------------------|---|---------------------------|--|---------------------------|--|-----------------------|---|---------------------------|---|---------------------------|
|                                     |                      |                                   |                               |  |   |                                       | Serotypes<br>targeted<br>in the<br>control<br>programm<br>e (f) | Other<br>serotyp<br>es(g) | Serotypes<br>targeted<br>in the<br>control<br>programme<br>(f) | Other<br>serotypes<br>(g) | Serotypes<br>targeted<br>in the<br>control<br>programme<br>(f) | Other<br>serotypes(g) | Serotypes<br>targeted<br>in the<br>control<br>programm<br>e (f) | Other<br>serotypes<br>(g) | Serotypes<br>targeted<br>in the<br>control<br>programme<br>(f)      | Other<br>serotyp<br>es(g) |

|           |               |    |        |    |        |    |   |   |   |   |   |   |   |   |   |   |
|-----------|---------------|----|--------|----|--------|----|---|---|---|---|---|---|---|---|---|---|
| RO121-AB  | Breeding Hens | 18 | 108400 | 12 | 78000  | 12 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO421-AR  | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO311-AG  | Breeding Hens | 24 | 71561  | 24 | 71561  | 24 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO211-BC  | Breeding Hens | 47 | 116685 | 30 | 91858  | 30 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO111-BH  | Breeding Hens | 10 | 60000  | 10 | 60000  | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO112-BN  | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO212-BT  | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO221-BR  | Breeding Hens | 6  | 31843  | 6  | 31843  | 6  | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO122-BV  | Breeding Hens | 60 | 659512 | 36 | 390463 | 36 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO321-BUC | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO222-BZ  | Breeding Hens | 16 | 86828  | 16 | 86828  | 16 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO312-CL  | Breeding Hens | 18 | 90000  | 18 | 90000  | 18 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO422-CS  | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO113-CJ  | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO223-CT  | Breeding Hens | 18 | 110506 | 18 | 110506 | 18 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO123-CV  | Breeding Hens | 2  | 11500  | 2  | 11500  | 2  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO313-DB  | Breeding Hens | 24 | 118892 | 24 | 118892 | 24 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO411-DJ  | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO224-GL  | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |



|          |               |    |        |    |        |    |   |   |   |   |   |   |   |   |   |   |
|----------|---------------|----|--------|----|--------|----|---|---|---|---|---|---|---|---|---|---|
| RO314-GR | Breeding Hens | 4  | 34119  | 4  | 34119  | 2  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO412-GJ | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO124-HR | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO423-HD | Breeding Hens | 8  | 52000  | 3  | 19500  | 3  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO315-IL | Breeding Hens | 30 | 283330 | 30 | 283330 | 30 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO213-IS | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO322-IF | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO114-MM | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO413-MH | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO125-MS | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO214-NT | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO414-OT | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO316-PH | Breeding Hens | 37 | 124748 | 37 | 124748 | 37 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO116-SJ | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO115-SM | Breeding Hens | 19 | 160000 | 19 | 160000 | 19 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO126-SB | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO215-SV | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RO317-TR | Breeding Hens | 0  | 0      | 0  | 0      | 0  | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

|              |               |            |                |            |                |            |          |          |          |          |          |          |          |          |          |          |
|--------------|---------------|------------|----------------|------------|----------------|------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| RO424-TM     | Breeding Hens | 0          | 0              | 0          | 0              | 0          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO225-TL     | Breeding Hens | 0          | 0              | 0          | 0              | 0          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO415-VL     | Breeding Hens | 0          | 0              | 0          | 0              | 0          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO226-VN     | Breeding Hens | 18         | 66947          | 12         | 30663          | 12         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO216-VS     | Breeding Hens | 6          | 47538          | 6          | 47538          | 5          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| <b>Total</b> | Breeding Hens | <b>365</b> | <b>2234409</b> | <b>307</b> | <b>1841349</b> | <b>304</b> | <b>0</b> | <b>4</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> |

Year: 2011

Situation on date: 31.12.2011

Animal species: breeding hens (Gallus gallus)

Disease: zoonotic salmonellosis

| Region<br>(County+NUTS<br>CODE) | Type of<br>flocks(b) | Total<br>numbe<br>r of<br>flocks<br>© | Total<br>number of<br>animals | Total<br>numbe<br>r of<br>flocks<br>under<br>the<br>progra<br>mme | Total<br>number of<br>animals<br>under the<br>programme | Numbe<br>r of<br>flocks<br>checke<br>d(d) | Number of positive<br>flocks (e)                                    |                           | Number of flocks<br>depopulated                                     |                           | Total number of animals<br>slaughtered or destroyed          |                           | Quantity of the eggs<br>destroyed (number or kg)             |                           | Quantity of eggs channelled<br>to egg products (number or<br>kg) |                       |
|---------------------------------|----------------------|---------------------------------------|-------------------------------|---|---|---|---|---------------------------|---|---------------------------|--|---------------------------|--|---------------------------|--|-----------------------|
|                                 |                      |                                       |                               |   |   |   | Serotype<br>s<br>targeted<br>in the<br>control<br>program<br>me (f) | Other<br>serotype<br>s(g) | Serotype<br>s<br>targeted<br>in the<br>control<br>program<br>me (f) | Other<br>serotypes(<br>g) | Serotypes<br>targeted in<br>the control<br>programm<br>e (f) | Other<br>serotypes(<br>g) | Serotypes<br>targeted in<br>the control<br>programm<br>e (f) | Other<br>serotypes(<br>g) | Serotypes<br>targeted in<br>the control<br>programm<br>e (f)     | Other<br>serotypes(g) |
| RO121-<br>AB                    | Breeding<br>Hens     | 18                                    | 110600                        | 12  | 72000   | 12  | 0   | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         | 0  | 0                     |
| RO311-<br>AG                    | Breeding<br>Hens     | 24                                    | 150734                        | 24  | 150734  | 24  | 0   | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         | 0  | 0                     |
| RO211-<br>BC                    | Breeding<br>Hens     | 36                                    | 308872                        | 36  | 308872  | 36  | 0   | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         | 0  | 0                     |
| RO111-<br>BH                    | Breeding<br>Hens     | 10                                    | 85350                         | 10  | 85350   | 10  | 0   | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         | 0  | 0                     |
| RO221-                          | Breeding<br>Hens     | 7                                     | 47626                         | 7   | 47626   | 7   | 0   | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         | 0  | 0                     |

|              |               |            |                |            |                |            |          |          |          |          |          |          |          |          |          |          |
|--------------|---------------|------------|----------------|------------|----------------|------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| BR           |               |            |                |            |                |            |          |          |          |          |          |          |          |          |          |          |
| RO122-BV     | Breeding Hens | 46         | 159641         | 28         | 159641         | 28         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO222-BZ     | Breeding Hens | 24         | 143032         | 12         | 143032         | 12         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO312-CL     | Breeding Hens | 36         | 201333         | 36         | 201333         | 36         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO223-CT     | Breeding Hens | 30         | 216776         | 30         | 216776         | 30         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO123-CV     | Breeding Hens | 2          | 7484           | 2          | 7486           | 2          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO313-DB     | Breeding Hens | 61         | 250000         | 61         | 250000         | 61         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO224-GL     | Breeding Hens | 1          | 4120           | 1          | 0              | 0          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO423-HD     | Breeding Hens | 8          | 46020          | 8          | 46020          | 8          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO315-IL     | Breeding Hens | 56         | 205123         | 43         | 113792         | 43         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO125-MS     | Breeding Hens | 2          | 11000          | 2          | 11000          | 2          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO316-PH     | Breeding Hens | 41         | 196828         | 36         | 150532         | 36         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO115-SM     | Breeding Hens | 30         | 150000         | 30         | 150000         | 30         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO226-VN     | Breeding Hens | 12         | 32733          | 12         | 32733          | 12         | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| RO216-VS     | Breeding Hens | 7          | 46830          | 7          | 46830          | 7          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| IDAH         | Breeding Hens | 0          | 0              | 0          | 0              | 0          | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        | 0        |
| <b>Total</b> | Breeding Hens | <b>451</b> | <b>2374102</b> | <b>397</b> | <b>2193757</b> | <b>396</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> |

## 6.4 Data on vaccination programme

Data are not available-In Romania the vaccination is not mandatory.

## 7. Targets

### 7.1. Targets related to testing :to investigate the presence of Salmonella in breeding flocks

**Disease (a): zoonotic Salmonella**

**Animal species:**

| Region (b)  | Type of the test (c) | Target population (d) | Type of sample (e)     | Objective (f) | Number of planned tests |
|-------------|----------------------|-----------------------|------------------------|---------------|-------------------------|
| All regions | Bacteriological      | Breeding hens         | Faeces boot/sock swabs | Surveillance  | 2000                    |
|             | Bacteriological      | Breeding hens         | Faeces boot/sock swabs | Confirmation  | 100                     |
| Total       | -                    | -                     | -                      | -             |                         |

(a) Disease and species if necessary

(b) Region as defined in the approved eradication programme of the Member State

(c) Description of the test

(d) Specification of the targeted species and the categories of targeted animals

(e) Description of the sample (faeces)

(f) Description of the objective ( e.g. qualification, surveillance, confirmation of suspected cases, monitoring of campaigns, seroconversion, control on deleted vaccines, testing of vaccine , control of vaccination

### 7.1.2. Targets on testing of breeding flocks

Year: 2013

Animal species:Gallus gallus

Disease/infection(a):Zoonotic Salmonella

| Region      | Type of flock <sup>(b)</sup> | Total number of flocks <sup>(c)</sup> | Total number of animals | Total n° of flocks under the programme | Total n° of animals under the programme | Expected n° of flocks to be checked <sup>(d)</sup> | Number of flocks <sup>(d)</sup> expected to be positive <sup>(a)</sup> |      |      | Number of flocks expected to be depopulated |      | Total n° of animals expected to be slaughtered or destroyed <sup>(f)</sup> |      | Expected quantity of eggs to be destroyed (number or kg) <sup>(d)</sup> |    | Expected quantity of eggs channelled to egg products (number or kg.) <sup>(d)</sup> |      |
|-------------|------------------------------|---------------------------------------|-------------------------|--|---|--|--|------|------|---|------|--|------|---|----|---|------|
|             |                              |                                       |                         |  |   |  | (a1)   | (a2) | (a3) | a4  | (a3) | (a4)   | (a3) | (a4)  | a3 | (a4)  | (a3) |
| All regions | Breeding hens                | 400                                   | 2500000                 | 400                                    | 2500000                                 | 400  | 2  | 2    | 50   | 10  | 50   | 8000   | 0    | 10000   | 0  | 10000   | 0    |
| Total       | Breeding hens                | 400                                   | 2500000                 | 400                                    | 2500000                                 | 400  | 2  | 2    | 50   | 10  | 50   | 8000   | 0    | 10000   | 0  | 10000   | 0    |

- (a) For zoonotic salmonella indicate the serotypes covered by the control programmes: (a1) for Salmonella Enteritidis,(a2) for Salmonella Typhimurium, (a3) for other serotypes-specify as appropriate, (a4) for Salmonella Enteritidis or Salmonella Typhimurium.
- (b) For example, breeding flocks (rearing,adult flocks),production flocks, laying hen flocks,etc. Flocks equals herds or as appropriate.
- (c) Total number of flocks existing in the region including eligible flocks and non-eligible flocks for the programme.
- (d) Check means to perform a flock level test under the programme for the presence of Salmonella. In this column a flock should not be counted twice even if it has been checked more than once.
- (e) If a flock has been checked, in accordance with footnote (d), more than once, a positive sample should be taken into account only once.

## 7.2. Targets on vaccination

### 7.2.1. Targets on vaccination of Breeding hens flocks- Data not available

Disease <sup>(a)</sup>: Zoonotic Salmonella

Animal species: Breeding hens gallus gallus

| Region       | Total number of herds in vaccination or programme | Total number of animals in vaccination programme | No of herds in vaccination programme | Targets on vaccination programme      |   |  |
|--------------|---|--|--------------------------------------|---------------------------------------|---|--|
|              |   |  |                                      | No of herds expected to be vaccinated | No of animals expected to be vaccinated | No of doses of vaccine expected to be administered |
| All regions  | -   | -  | -                                    | -                                     | -                                       | -  |
| <b>Total</b> | -   | -  | -                                    | -                                     | -                                       | -  |

**The vaccination is not mandatory and the costs regarding purchase of vaccine doses and the vaccination are incurred by the business operators.**

## 8. Detailed analysis of the cost of the programme<sup>1</sup>

Costs mentioned below are estimated for a one-year period (1 of January 2013-31 of December 2013 )

| <b>Costs related to</b>             | <b>Specification</b>   | <b>Number of units</b> | <b>Unitary cost in €</b> | <b>Total amount in €</b> | <b>Community funding requested (yes/no)</b> |
|-------------------------------------|--|------------------------|--------------------------|--------------------------|---|
| <b>1. Testing</b>                   |  |                        |                          |                          |   |
| <b>1.1. Cost of the analysis</b>    |  |                        |                          |                          |   |
|                                     | Test: bacteriological exam for Salmonella spp. detection   | 2000                   | 16.70                    | 33400                    | YES   |
|                                     | Test : Salmonella spp. serotyping  | 100                    | 64.97                    | 6497                     | YES   |
|                                     | Test : the detection of antimicrobials or bacterial growth inhibitory effect in tissues from birds from flocks tested for Salmonella | 100                    | 47.44                    | 4744                     | YES   |
| <b>1.2. Cost of sampling</b>        | Official samples harvest   | 2000                   | 0,5                      | 1000                     | YES   |
|                                     | Disposable sterile containers for sampling of faeces   | 2000                   | 0,5                      | 1000                     | NO  |
|                                     | One use gloves-pairs   | 4000                   | 0,1                      | 400                      | NO  |
|                                     | Boot swabs   | 4000                   | 2                        | 8000                     | NO  |
| <b>1.3. Other costs</b>             | Overcoats  | 2000                   | 1                        | 2000                     | NO  |
| <b>2. Vaccination</b>               |  |                        |                          |                          |   |
| <b>2.1. Purchase of vaccine</b>     |  |                        |                          |                          |   |
| <b>2.2. Distribution costs</b>      |  |                        |                          |                          |   |
| <b>2.3. Administering costs</b>     |  |                        |                          |                          |   |
| <b>2.4. Control costs</b>           |  |                        |                          |                          |   |
| <b>3. Slaughter and destruction</b> |  |                        |                          |                          |   |

|  |   |              |            |                 |            |
|--|---|--------------|------------|-----------------|------------|
| <b>3.1. Compensation of animals</b>  | Compensation for the value of a parent breeding bird of <i>Gallus gallus</i> culled   | <b>8000</b>  | <b>8</b>   | <b>64000</b>    | <b>YES</b> |
|  | Compensation for hatching eggs of parent breeding <i>Gallus gallus</i> , per hatching egg destroyed                                     | <b>10000</b> | <b>0,4</b> | <b>4000</b>     | <b>YES</b> |
| <b>3.2. Transport costs</b>  |   |              |            |                 |            |
| <b>3.3. Destruction costs</b>  |   |              |            |                 |            |
| <b>3.4. Loss in case of slaughtering</b>   |   |              |            |                 |            |
| <b>3.5 Costs from treatment of products (eggs, hatching eggs, heat treat of broilers, etc)</b> |   |              |            |                 |            |
| <b>4. Cleaning and disinfection</b>  |   |              |            |                 |            |
|  | Test: Bacteriological test to verify the efficiency of disinfection of poultry houses after depopulation of a Salmonella-positive flock | <b>200</b>   | <b>10</b>  | <b>2000</b>     | <b>YES</b> |
| <b>5. Salaries (staff contracted for the programme only)</b>                                   |   |              |            |                 |            |
|  |   |              |            |                 |            |
| <b>6. Consumables and specific equipment</b>   |   |              |            |                 |            |
| <b>7. Other costs</b>  |   |              |            |                 |            |
| <b>TOTAL</b>   |   |              |            | <b>115641 €</b> | <b>YES</b> |





# Romanian National Control Programme for *Salmonella* in Broilers (*Gallus gallus*) 2013

## Part A

### a. Aim of the programme

To reduce or maintain the low prevalence of *Salmonellas* with public health significance in broiler flocks (*Gallus gallus*) in Romanian holdings producing broilers intended for human consumption at least to the target levels set out in Regulation (EC) No 646/2007 which is a maximum percentage of broiler flocks remaining positive for *Salmonella Enteritidis* and *Salmonella Typhimurium*, including also the monophasic strains of *S. Typhimurium*, for 1% or less.

The programme for the control of *Salmonella Enteritidis* and *Salmonella Typhimurium* in broiler flocks of *Gallus gallus* has been in operation in Romania since 2008. As a result, the number of *Salmonella Enteritidis* and *Salmonella Typhimurium* infected broiler flocks of *Gallus gallus* in Romania is currently very low. During 2011, a totally of 4910 broiler flocks were tested for *Salmonella* infection from which a no of 1535 were tested in official control and there were only 11 flocks positive for *Salmonella Typhimurium* and *Salmonella Enteritidis*. The prevalence for the target serotypes in broiler flock in 2011 was 0.7%, which is low and below the Community target.

### b. Sampling programme

The National Control Programme for *Salmonella* encompasses the following serovars of zoonotic *Salmonella*: *Salmonella enteritidis* and *Salmonella typhimurium*.

The sampling programme will be in accordance to Regulation **2160/2003 EC** and **Regulation 646/2007 EC**. We have also taken into account the Regulation 1177/2006 EC provisions and some amendments of Regulation 1168/2006 EC.

The programme was elaborated in compliance with the requirements of Council Regulation **(EC) No 646/2007** of 12 June 2007 implementing Regulation **(EC) No 2160/2003** of the European Parliament and of the Council as regards a Community target for the reduction of the prevalence of *Salmonella enteritidis* and *Salmonella typhimurium* in broilers and repealing Regulation **(EC) No 1091/2005**.

**c. Demonstrate the evidence that it complies with the specific requirements laid down in Parts E of Annex II to Regulation (EC) no 2160/2003**

***Specific requirement concerning fresh meat***

When the a broilers flock is confirmed with the presence of *Salmonella enteritidis* or *Salmonella typhimurium*:

- a. Fresh meat from broilers may be placed on the market on the condition that it meets the requirement of absence of Salmonella in 25 grams from the meat.
- b. The requirement laid down in point (a) does not apply to fresh poultry meat destined for heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene.
- c. The criteria laid down in point (a) does not apply to fresh poultry meat destined for industrial heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene.

**1. General**

**1.1 Summary referring to the occurrence of the salmonellosis in Romania**

In Romania in 2009 from broiler flocks were harvest 1411 official samples. Only 2 flocks were find positive for the zoonotic Salmonella serovars in the aim The National Control Programme for Salmonella.

The programme for the control of Salmonella Enteritidis and Salmonella Typhimurium in broiler flocks of Gallus gallus has been in operation in Romania since 2008. As a result, the number of Salmonella Enteritidis and Salmonella Typhimurium infected broiler flocks of Gallus gallus in Romani is currently very low. During 2011, a totally of 4910 broiler flocks were tested for Salmonella infection from wich a no of 1535 were tested in official control and there were only 11 flocks positive for Salmonella Typhimurium and Salmonella Enteritidis . The prevalence for the target serotypes in broiler flock in 2011 was 0.7%, which is low and below the Community target.

**The incidence of various serotypes of Salmonella in broilers , in 2011 in Roumania**

| NO.          | SEROTYPE      | NO. OF STRAINS<br>in broilers |
|--------------|---------------|-------------------------------|
| 1.           | S.Afula       | 1                             |
| 2.           | S.Agona       | 21                            |
| 3.           | S.Amsterdam   | 4                             |
| 4.           | S.Blockley    | 1                             |
| 5.           | S.Bredeney    | 3                             |
| 6.           | S.Enteritidis | 40                            |
| 7.           | S.Glostrup    | 7                             |
| 8.           | S.Hadar       | 5                             |
| 9.           | S.Infantis    | 342                           |
| 10.          | S.Inganda     | 2                             |
| 11.          | S.Irumu       | 1                             |
| 12.          | S.Kentucky    | 93                            |
| 13.          | S.Kottbus     | 3                             |
| 14.          | S.Liverpool   | 37                            |
| 15.          | S.Livingstone | 25                            |
| 16.          | S.Mbandaka    | 17                            |
| 17.          | S.Montevideo  | 1                             |
| 18.          | S.Saintpaul   | 3                             |
| 19.          | S.Senfteberg  | 7                             |
| 20.          | S.Taksony     | 35                            |
| 21.          | S.Tennessee   | 9                             |
| 22.          | S.Thompson    | 10                            |
| 23.          | S.Tilburg     | 4                             |
| 24.          | S.Uganda      | 2                             |
| <b>TOTAL</b> |               | <b>655</b>                    |

## 1.2 The structure and organization of the relevant Competent Authorities

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in flocks of broilers of *Gallus gallus* is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**

---

The departments responsible for implementing the National Control Programme of Salmonella in broilers are:

- for implementing at national level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary General Directorate** ;
- in respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the verification of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**
- Concerning the slaughter of broilers flocks, the **Hygiene and Veterinary Epidemiology Directorate** is responsible for public health protection, through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.
- In respect of EC Regulation No 183/2005 on feed hygiene at the central level is **The Directorate for Technical Coordination of Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**
- at county level for implementing the Regulation (EC) No 2160/2003,882/2004 and 183/2005 there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the national reference laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

#### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

**Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România

Tel: 0374.322.013 / 0374.322.000 Fax: 0214.113.394

E-mail: [office@idah.ro](mailto:office@idah.ro)

- **H.I.P.V.H.** (Hygiene Institute and Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

#### **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED**

**Hygiene Institute and Veterinary Public Health**

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201  
 Tel: 021.252.46.51 Fax: 021.252.00.61 E-mail: [iisppv @ iisppv.ro](mailto:iisppv@iisppv.ro)

### 1.3 Approved laboratories where samples collected within the programme are analyzed

**Diagnostics:** The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL is show below:

| Nr. Crt. | NAME                             | ADRESS   | TELEFON/ E-MAIL   |
|----------|----------------------------------|--|---|
| 1        | <b>I.D.S.A</b>                   | București, Str. Dr.Staicovici, nr.63, sector 5, Cod 050557         | 0374/322.013<br><a href="mailto:office@idah.ro">office@idah.ro</a>  |
| 2        | <b>L.S.V.S.A. Alba</b>           | Alba Iulia, Str. Lalelelor nr.7A, Cod 510217, Jud. Alba            | 0258/835.915<br><a href="mailto:office-alba@ansvsa.ro">office-alba@ansvsa.ro</a>  |
| 3        | <b>L.S.V.S.A. Bacău</b>          | Bacau, Str Bucovinei, Nr 21, Jud. Bacău                            | 0234/586.233, int.111<br><a href="mailto:office-bacau@ansvsa.ro">office-bacau@ansvsa.ro</a>   |
| 4        | <b>L.S.V.S.A.Bistrița-Năsăud</b> | Bistrita, Str. Tarpiului 26, Cod 420062, Jud. Bistrița Năsăud      | 0263/206.027; 0263/224.974<br><a href="mailto:office-bistrita-nasaud@ansvsa.ro">office-bistrita-nasaud@ansvsa.ro</a>  |
| 5        | <b>L.S.V.S.A. Botoșani</b>       | Botosani , Str Tudor Vladimirescu Nr.17, Jud. Botoșani             | 0231/512.766<br><a href="mailto:office-botosani@ansvsa.ro">office-botosani@ansvsa.ro</a>  |
| 6        | <b>L.S.V.S.A. Brașov</b>         | Brașov, Str. Calea Feldioarei Nr. 20 A, Cod 500450, Jud.Brașov     | 0268/440.257<br><a href="mailto:office-brasov@ansvsa.ro">office-brasov@ansvsa.ro</a><br><a href="mailto:dsvbv@rdslink.ro">dsvbv@rdslink.ro</a>                    |
| 7        | <b>L.S.V.S.A. Brăila</b>         | Brăila, str. Calea Galați, nr.344, Cod 810385, Jud. Brăila         | 0239/610.689<br>dsv@braila.rdsnet.ro<br><a href="mailto:office-braila@ansvsa.ro">office-braila@ansvsa.ro</a>  |
| 8        | <b>L.S.V.S.A. Buzău</b>          | Buzau, Str. Horticolei, nr. 58 bis, Jud. Buzau                     | 0238-725001<br>0238-725002<br><a href="mailto:office-buzau@ansv.ro">office-buzau@ansv.ro</a>  |
| 9        | <b>L.S.V.S.A. Călărași</b>       | Călărași, Str. Prelungirea Dobrogei , nr. 4, Jud.Călărași          | 0242/313.676,<br><a href="mailto:office-calarasi@ansvsa.ro">office-calarasi@ansvsa.ro</a> ;<br><a href="mailto:office_dsv@satline.ro">office_dsv@satline.ro</a> ; |
| 10       | <b>L.S.V.S.A Constanța</b>       | Constanta, Sos. Mangaliei Nr. 78, Jud. Constanța                   | 0241-682-417,<br>email: <a href="mailto:dsvet@rdsect.ro">dsvet@rdsect.ro</a>  |
| 11       | <b>L.S.V.S.A. Cluj</b>           | Cluj-Napoca, Piata Marasti Nr. 1, Jud. Cluj                        | 0264-445729, 0264-448177<br><a href="mailto:office-cluj@ansvsa.ro">office-cluj@ansvsa.ro</a><br><a href="mailto:dsvcj@rdslink.ro">dsvcj@rdslink.ro</a>            |
| 12       | <b>L.S.V.S.A. Dâmbovița</b>      | Târgoviște, Str. I.C. Brătianu, nr. 35, Cod 130055, Jud. Dâmbovița | 0372/737818<br><a href="mailto:office-dambovita@ansvsa.ro">office-dambovita@ansvsa.ro</a>   |
| 13       | <b>L.S.V.S.A. Giurgiu</b>        | Giurgiu, Str. Bucuresti nr.72, Jud. Giurgiu                        | 0246/ 230.491<br><a href="mailto:office-giurgiu@ansvsa.ro">office-giurgiu@ansvsa.ro</a>   |
| 14       | <b>L.S.V.S.A. Gorj</b>           | Târgul-Jiu, Str.Ecaterina Teodoroiu Nr.523, Jud. Gorj              | 0253/226.033<br><a href="mailto:dsv@intergorj.ro">dsv@intergorj.ro</a><br><a href="mailto:office-gorj@ansvsa.ro">office-gorj@ansvsa.ro</a>                        |

|    |                             |  |  |
|----|-----------------------------|--|--|
| 15 | <b>L.S.V.S.A Hunedoara</b>  | Deva, Str. 22 Decembrie Nr. 226, Jud. Hunedoara                        | 0254-221145, 0254-230527<br><a href="mailto:office-hunedoara@ansvsa.ro">office-hunedoara@ansvsa.ro</a> |
| 16 | <b>L.S.V.S.A. Ialomița</b>  | Slobozia, str. Lacului, nr. 12 , Jud. Ialomița                         | 0243/ 232.069<br><a href="mailto:dsvsa-ialomita@ansvsa.ro">dsvsa-ialomita@ansvsa.ro</a>                |
| 17 | <b>L.S.V.S.A. Iași</b>      | Iași, Aleea Mihail Sadoveanu 10, Jud. Iași                             | 0232-267501<br><a href="mailto:office-iasi@ansvsa.ro">office-iasi@ansvsa.ro</a>                        |
| 18 | <b>L.S.V.S.A. Maramureș</b> | Baia Mare, Str. Vasile Alecsandri, Nr. 66, Jud. Maramureș              | 0262/ 224.031<br><a href="mailto:office-maramures@ansvsa.ro">office-maramures@ansvsa.ro</a>            |
| 19 | <b>L.S.V.S.A. Mureș</b>     | Târgu Mures, str. Podeni nr.10 cod 540253, Jud. Mureș                  | 0265/ 314.975<br><a href="mailto:office-mures@ansvsa.ro">office-mures@ansvsa.ro</a>                    |
| 20 | <b>L.S.V.S.A. Neamț</b>     | Piatra Neamț, Aleea Tiparului, nr. 12, cod 610263, Jud. Neamț          | 0233/ 223.259<br><a href="mailto:office-neamt@ansvsa.ro">office-neamt@ansvsa.ro</a>                    |
| 21 | <b>L.S.V.S.A. Prahova</b>   | Ploiesti, Str. Corlatesti, nr. 11, Jud. Prahova,                       | 0244 / 57.17.51<br><a href="mailto:office-prahova@ansvsa.ro">office-prahova@ansvsa.ro</a>              |
| 22 | <b>L.S.V.S.A. Satu Mare</b> | Satu Mare, Str. Lăcrimioarei, Nr. 37, cod poștal 440067, Jud.Satu Mare | 0261/ 715.956<br><a href="mailto:office-satu-mare@ansvsa.ro">office-satu-mare@ansvsa.ro</a>            |
| 23 | <b>L.S.V.S.A. Sibiu</b>     | Sibiu, str. Calea Șurii Mari, nr. 21, cod 550089, Jud. Sibiu           | 0269/223.069<br><a href="mailto:office-sibiu@ansvsa.ro">office-sibiu@ansvsa.ro</a>                     |
| 24 | <b>L.S.V.S.A. Suceava</b>   | Suceava, Str. Scurta Nr. 2, Jud Suceava                                | Tel: 0230-522848,<br><a href="mailto:dsvsv@suceava.rdsnet.ro">dsvsv@suceava.rdsnet.ro</a>              |
| 25 | <b>L.S.V.S.A. Timiș</b>     | Timisoara, Str. Surorile Martir Caceu Nr.4, Jud. Timiș                 | 0256-204911<br><a href="mailto:office-timis@ansv.ro">office-timis@ansv.ro</a>                          |
| 26 | <b>L.S.V.S.A. Vaslui</b>    | Bârlad, str. Trestiana, nr. 2, cod 731030, Jud.Vaslui                  | 0235-421121,<br>0235-421413<br><a href="mailto:office-vaslui@ansv.ro">office-vaslui@ansv.ro</a>        |
| 27 | <b>L.S.V.S.A. Vrancea</b>   | Focsani,B-dul Brailei, nr. 121 bis, cod 620122, Jud.Vrancea            | 0237-215561,<br>0237-232727<br><a href="mailto:office-vrancea@ansv.ro">office-vrancea@ansv.ro</a>      |

All Laboratories have to use the methods of the diagnostic presented at the point 1.4.

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the serotype according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

#### **1.4 Methods used in the examination of the samples in the framework of the programme**

Samples harvest by operators and samples harvest as official controls are prepared and tested in accordance with the requirements of the Commission Regulation (EC) No 646/2007 Annex, using the method recommended by the Community Reference Laboratory for *Salmonella* in Bilthoven, Netherlands. The method is described in the current version of ISO 6579:/A1:2007): 'Detection of *Salmonella* spp. in animal faeces and in samples of the primary production stage'. A semi-solid medium (modified semi-solid Rappaport-Vassiladis medium, MSRV) is used as the single selective enrichment medium. At least one isolate from each positive sample shall will be serotyped according to the Kaufmann-White scheme.

Isolates of *S. Enteritidis* and *S. Typhimurium* will also be phage typed. In general the sensitivity of a panel of 10 antimicrobials will be determined.

### **1.5 Official controls**

In accordance with EC Regulation No 646/2007 provisions, each **S.V.F.S.D** will select each year at least 10% of holdings with more than 5000 birds. The Competent Authority or its agent will select one flock at random on the holding that is within 3 weeks of slaughter expedition. The selected flock will be sampled (two pairs of boot swabs normally) in accordance with the Annex of EC Regulation No 646/2007 provisions.

In addition where there has been detection of *S. Enteritidis* or *S. Typhimurium* in operator sampling program from previous flocks on the holding, control advice will be provided including on-farm visits by experts in *Salmonella* control when appropriate. All flocks of the holding will be officially sampled when the next crop of birds is placed.

The use of antimicrobials (as defined in Regulation (EC) No 1177/2006) will be checked when the official sample is taken. If the flock is under antimicrobial medication for animal health or animal welfare reasons, the flock will be sampled again after the withdrawal period of the product specified in the Marketing Authorization. Flock owners are required to keep records of antimicrobial used and to make these records available to the competent authorities.

The records of samples harvest by the operator will be made available for inspection to the Competent Authority or its agent and will provide details of the identity of the flock sampled, date of sample, slaughter date, type of sample, laboratory carrying out the examination, and the result. The number of flocks on the holding and the number of birds present will be recorded.

### **Official controls at other stages of the food chain.**

Under the terms of the EC Feed Hygiene **Regulation 183/2005** feed businesses operators must be approved or registered by the Local Authority. Approvals/registrations are issued for the producers of compound feeds, feed materials, feed additives and premixtures. Approval requires a prior-inspection visit by the Local Authority to ensure that the establishments are in conformity with the required standards. The registration is followed by placing of premises on the list of feed business operators. The competent authority performs checks according to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order.



## **1.6 Measures taken by the Competent Authorities with regard to animals or products in which the presence of *Salmonella* spp. have been detected**

In case of suspicion or confirmation of *Salmonella enteritidis* or *Salmonella typhimurium* the NRL/C.S.V.F.S.L. shall notify immediately the N.S.V.F.S.A and the local C.S.V.F.S.D.

In case of suspicion of infection the local C.S.V.F.S.D and the relevant authorities:

- prohibited the movement of broilers
- take additional samples for confirmation of infection

When a broiler flock of *Gallus gallus* is suspected of being infected with *Salmonella Enteritidis* or *Salmonella Typhimurium* the flock will be investigated. The flock is suspected of being infected when *S. Enteritidis* or *S. Typhimurium* is isolated from a sample of faeces, or boot swabs, carried out privately or as required by either the operator or the Competent Authority as detailed in the Annex to Regulation (EC) No 646/2007. Tissue/organs may be taken from birds as part of the investigation of clinical disease by the veterinarian; these cases will be discussed and additional follow up investigation carried out as appropriate, along with advice on *Salmonella* control.

Competent Authority will notify the operator to clean and disinfect the building from which the infected flock originated. After cleaning and disinfecting of the building the operator may be required to take swabs from a number of sites in the building and submit them to an approved laboratory in view to be tested for *Salmonella* in order to check the efficiency of the hygiene measures taken. In cases where *S. Enteritidis* or *S. Typhimurium* was isolated the cleaning and disinfection may be checked by the Competent Authority or its agent.

If the results of post-cleaning and disinfection monitoring of *Salmonella* are positive for *S. Enteritidis* or *S. Typhimurium*, the next crop (cycle) will be monitored under supervision of the Competent Authority or its agent. If *Salmonella* is isolated in this subsequent crop of birds the holding will be placed under official control; re-stocking of the house will be permitted only if the supervised post-cleaning and disinfection samples from the house are negative.

When the broilers are confirmed for the presence of *Salmonella enteritidis* or *Salmonella typhimurium*:

1. Fresh meat from broilers will be placed on the market with the condition that it meets the requirement concerning the absence of *Salmonella* in 25 grams of meat.
2. The requirement laid down in point 1 does shall not apply to fresh poultry meat intended for heat treatment or another treatment to eliminate *Salmonella* in accordance with Community legislation on food hygiene.

3. The criteria laid down in point 1 shall not apply to fresh poultry meat intended for industrial heat treatment or another treatment to eliminate Salmonella in accordance with Community legislation on food hygiene.

### **1.7 Relevant national legislation.**

- N.S.V.F.S.A President **Order no. 34** to approve the sanitary veterinary norm regarding to monitoring zoonoses and zoonotic agents

- Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order

- N.S.V.F.S.A President **Order no. 205** for the approval of national reference laboratories and duties

- N.S.V.F.S.A President **Order no. 160** for the approval of the sanitary veterinary norm regarding the community reference laboratory for the zoonoses epidemiology and Salmonella and the national reference laboratory for Salmonella

### **1.8 Financial assistance provided to food and feed businesses operators in the context of the National Control Programme**

In broilers flocks of *Gallus gallus* financial assistance is provided in the context of the control programme. This is set up by the **Government Decision no. 1214/2009** regarding the methodology for for determining and paying compensation to be paid to owners of slaughtered animals, killed or otherwise affected by the rapid liquidation of outbreaks of animal diseases

The amount to compensate the animal owners in the context SNCP is determined in accordance with the provision of Government Decision No. 1214/2009 and is established by an evaluation committee consisting of:

a) the representative of the county sanitary veterinary and food safety directorate (local CA)

b) local representative of the Ministry of Agriculture and Rural Development with responsibilities in determining the genetic value of the animal

c) the representative of the decision local unit from the local disease control center, designated by the Prefect,

d) the local mayor or the person designated by him.

Convening of the evaluation committee is made by the Prefect at the written request of the county sanitary veterinary and food safety directorate (local CA).

On the proposal of the local county sanitary veterinary and food safety directorate the compensation committee considers and approves the replacement value of the animals according to the genetic value of animal, zootechnical value, sex, age, weight, physiological status, category production, at market price at the

time when the liquidation of animals from the outbreak disease took place, and the average unit value based on the total amount of compensation for animals or products concerned according to Art. 4 of Regulation (EC) no. 349/2005 of the Commission, with the following amendments.

Romania applied to the Commission for co-financing for certain aspects of the control programme within the terms of Council Decision **2009/470/EC** on expenditure in the veterinary field.

## **2. Food and Feed businesses covered by the programme.**

The programme for the control of *Salmonella* in broilers of Gallus gallus is part of the controls along the whole food chain.

### **2.1 The structure of the production of the given species and products thereof.**

The structure of the broilers flocks in Romania in 2011 is presented in the table below

| Region                         | Type of flocks(b) | Total number of flocks | Total number of animals | Total number of holdings |
|--------------------------------|-------------------|------------------------|-------------------------|--------------------------|
| <b>Roumania (all counties)</b> | <b>Broiler</b>    | <b>6218</b>            | <b>123515372</b>        | <b>268</b>               |

**In Romania in 2011 there were 268 holdings with 6218 broiler flocks in total.**

### **2.2 The structure of the production of food**

At the end of the growing period the birds are slaughtered. In accordance with Regulation 853/2004 provisions, poultry meat intended for human consumption must be obtained in approved slaughterhouses. There are **40** approved slaughterhouses **in Romania**.

Broilers sold at retail level within Romania are required to be marked with a code identifying the establishment (production site), and the county of origin. This mark can be applied direct to the product, the wrapping, packaging or be printed on a label affixed to the product, the wrapping or the packaging. This is in accordance with EU Regulation No. 853/2004, Annex II, Section I (Identification Marking)

Poultry feed is supplied to farms by a small number of manufacturers. The major manufacturers of poultry feed operate to assurance schemes, apply HACCP principles and monitor for Salmonella.

## 2.3 Relevant guides for good animal husbandry practices or other guidelines

Some of the relevant guides for good animal husbandry practices are presented at the below link:

[http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella\\_12404ro.pdf](http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella_12404ro.pdf)

[http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010\\_13247ro.pdf](http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010_13247ro.pdf)

[http://www.ansvsa.ro/documente/admin/community\\_guide\\_layers\\_hygiene\\_practice\\_pullet\\_egg\\_ro\\_17813ro.pdf](http://www.ansvsa.ro/documente/admin/community_guide_layers_hygiene_practice_pullet_egg_ro_17813ro.pdf)

Other relevant guides for good animal husbandry practices and other guidelines are under constructions.

All farm registered must have their individual plan for good farmer practices, which are subject to approval by the official veterinarian responsible for the control of the holding concerned

## 2.4 Routine veterinary supervision of farms

The owner is responsible for the health and welfare of the poultry on the holding, and for ensuring that a veterinarian is consulted on disease and welfare issues as appropriate. It is mandatory for each holding to have a contract with a private veterinarian who is responsible for veterinary care. A veterinarian on behalf of the the Competent Authority carries out inspections on farms for animal welfare reasons, to take samples for residues, and to check medicine records.

Also a veterinarian on behalf of the Competent Authority visit the farms and take official samples in the framework of Salmonella NCP according with the legislation in force.

It is mandatory for each county sanitary veterinary and food safety directorate (local CA) to report to the NSVFSA every month the number of samples and results of these tests for each flock. Also the Salmonella NRL has the obligation to notify immediately NSVFSA and CSVFSD each positive sample for the relevant Salmonella.

## 2.5 Registration of farms

All commercial poultry flocks in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centers / holdings of origin and means of transport for health and

animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products.

## **2.6 Record-keeping at farms**

a. All broiler flock operators are required to keep records of veterinary medicines use, including vaccines, which must be available for inspection.

b. Records relating to movement of flocks on to/off the holding must be kept.

c. Records giving details of sampling for *Salmonella* and results will be kept either at the holding or be readily available.

## **2.7 Documents to accompany animals when dispatched**

Operators wishing to export birds or hatching eggs to another EU Member State (or certain third countries) must comply with **EU Directive 2009/158/EC** and ensure that the consignment is accompanied by a completed and signed Intra-trade Animal Health Certificate (ITAHC) for poultry breeding and production. This must be completed and signed by the Official Veterinarian as well as the operator to confirm compliance with the relevant articles of Directive.

The ITAHC will also require the reference number of the operator's poultry health certificate. The ITAHC is amended to include the results of the last test for Salmonella as is required in the Commission Regulation (EC) 2160/2003 Art 9.1 prior to any dispatching of the live animals, or hatching eggs, from the food business operators place of origin. The date and the result of testing are included in the relevant health certificate provided in the Community legislation.

## **2.8 Other relevant measures to ensure the traceability of animals.**

It is require to the operators of poultry flocks to keep records of poultry or hatching eggs entering or leaving the premises. The records must contain information on the number, date, and origin or destination. These records must be available to the Competent Authority for inspection.

All official veterinary health certificates issued for the intra community trade of poultry and hatching eggs are recorded on the Trade Control and Expert System (TRACES). This system allows tracking of exports of live animals and hatching eggs accompanied by veterinary health certification. TRACES generate ITAHCs issued for intra-Community movements. TRACES are an internet-based service which is owned and maintained by the Commission.

## **PART B**

### **1. Identification of the programme**

Member State: **Romania**  
Disease: **Salmonella Enteritidis and Typhimurium in broilers poultry (GALLUS gallus)**

Year of implementation: **2013**  
Reference of this document: **National Sanitary Veterinary and Food Safety Authority  
No. /**

Contact : **Dr. Nicolae LAZĂR,  
Director,  
phone: +4 021 315 78 75;  
fax: +4021 31249 67,  
e-mail: [lazar.niculae@ansvsa.ro](mailto:lazar.niculae@ansvsa.ro)**

**Date sent to the Commission : 27.04.2012**

## 2. Historical data on the epidemiological evolution of the disease

*Salmonella* has been recognized as an important zoonotic pathogen for many years. *Salmonella Enteritidis* and *Salmonella Typhimurium* have accounted for the majority of cases of human salmonellosis and have consistently been the most commonly implicated pathogens in general outbreaks of food-borne disease.

This National Control Programme of *Salmonella* was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 646/2007. The national control programme for *Salmonella* broilers flock was coming into effect in January 2009.

In line with Regulation EC No 1003/2005 since 01 January 2007 the programme has been enhanced and includes the control of *Salmonella* Enteritidis, *Salmonella Typhimurium* S. Hadar, S. Infantis and S. Virchow in breeding flocks. As a result of the control programme the number of *Salmonella Enteritidis* and *Salmonella Typhimurium* infected breeding flocks of *Gallus gallus* in Roumania is currently low. Of the other three *Salmonella* serovars, *Salmonella* Hadar, *Salmonella* Infantis and *Salmonella* Virchow, the occurrence is likewise at very low levels. Breeding flocks which are confirmed to be infected with zoonotic *Salmonella* are compulsorily slaughtered.

The success of the control programme in breeding flocks means that the day old chicks of broilers which are placed on farms should be free of *S. Enteritidis* and *S. Typhimurium*.

In Roumania in 2009 from broiler flocks were harvest 1411 official samples. Only 2 flocks were find positive for the zoonotic *Salmonella* serovars in the aim The National Control Programme for *Salmonella*.

The programme for the control of *Salmonella* Enteritidis and *Salmonella* Typhimurium in broiler flocks of *Gallus gallus* has been in operation in Romania since 2008. As a result, the number of *Salmonella* Enteritidis and *Salmonella* Typhimurium infected broiler flocks of *Gallus gallus* in Romani is currently very low. During 2011, a totally of 4910 broiler flocks were tested for *Salmonella* infection from wich a no of 1535 were tested in official control and there were only 11 flocks positive for *Salmonella* Typhimurium and *Salmonella* Enteritidis . The prevalence for the target serotypes in broiler flock in 2011 was 0.7%, which is low and below the Community target.

## The incidence of various serotypes of *Salmonella* in broilers flocks, in 2011

| NO.          | SEROTYPE      | NO. OF STRAINS in broilers |
|--------------|---------------|----------------------------|
| 21.          | S.Afula       | 1                          |
| 22.          | S.Agona       | 21                         |
| 23.          | S.Amsterdam   | 4                          |
| 24.          | S.Blockley    | 1                          |
| 25.          | S.Bredeney    | 3                          |
| 26.          | S.Enteritidis | 40                         |
| 27.          | S.Glostrup    | 7                          |
| 28.          | S.Hadar       | 5                          |
| 29.          | S.Infantis    | 342                        |
| 30.          | S.Inganda     | 2                          |
| 31.          | S.Irumu       | 1                          |
| 32.          | S.Kentucky    | 93                         |
| 33.          | S.Kottbus     | 3                          |
| 34.          | S.Liverpool   | 37                         |
| 35.          | S.Livingstone | 25                         |
| 36.          | S.Mbandaka    | 17                         |
| 37.          | S.Montevideo  | 1                          |
| 38.          | S.Saintpaul   | 3                          |
| 39.          | S.Senfenberg  | 7                          |
| 40.          | S.Taksony     | 35                         |
| 21           | S.Tennessee   | 9                          |
| 22           | S.Thompson    | 10                         |
| 23           | S.Tilburg     | 4                          |
| 24           | S.Uganda      | 2                          |
| <b>TOTAL</b> |               | <b>655</b>                 |

It is a statutory requirement for all laboratories which isolate *Salmonella* from a flock of chickens or its environment to report the finding and supply the isolate to the National Reference Laboratory (NRL) for *Salmonella*. The isolates are serotyped, phage-typed, where appropriate, and tested for antimicrobial sensitivity by the NRL. This information is recorded and analyzed. The number of reports received depends on the level and sensitivity of monitoring which is undertaken by



the producers. The reports provide useful information on the serovars which are most common in poultry, and indicate trends.

### 3. **Description of the submitted programme**

#### **Objectives**

The Community target for the reduction of *Salmonella enteritidis* and *Salmonella typhimurium* in broilers flocks of *Gallus gallus* shall be a reduction of the maximum percentage of remaining positive flocks to 1% or less by 31 December. The main objective of our programme for the reduction of *Salmonella enteritidis* and *Salmonella typhimurium including also the monophasic strains of S. Typhimurium* in broilers flocks of *Gallus gallus* shall be a reduction of the maximum percentage of positive flocks to 1 % or less by 31 December 2011

#### **Target animal population**

The National Control Programme for *Salmonella* in broilers will be held in all holdings of broiler flocks consisting of at least 500 poultry of *Gallus gallus*. Broilers holdings which have between 500 and 5,000 of birds will not be the subject of official testing, but will perform tests on the initiative of operators (self-control) within 3 weeks prior to depopulation and sending the birds abattoir.

Small flocks that are reared to supply meat for private domestic use, or small quantities of primary product supplied directly by the producer to the final consumer, or to local retail establishments directly supplying the primary product to the final consumer, will be exempt, as permitted in Regulation (EC) No 2160/2003 Article 1.3.

The number of holdings and broiler flocks in the broiler sector at the end 2011 are shown **in the table below.**

**Table . Number of flocks in the broiler sector**

| Region<br>(County+NUTS<br>CODE) | Type of<br>flocks(b) | Total<br>number<br>of<br>flocks | Total number<br>of animals |
|---------------------------------|----------------------|---------------------------------|----------------------------|
| RO121-AB                        | <b>broiler</b>       | 77                              | 954700                     |
| RO421-AR                        | <b>broiler</b>       | 8                               | 112000                     |
| RO311-AG                        | <b>broiler</b>       | 62                              | 989460                     |
| RO211-BC                        | <b>broiler</b>       | 127                             | 3140803                    |
| RO111-BH                        | <b>broiler</b>       | 88                              | 3002534                    |
| RO112-BN                        | <b>broiler</b>       | 18                              | 225000                     |
| RO212-BT                        | <b>broiler</b>       | 33                              | 500000                     |
| RO221-BR                        | <b>broiler</b>       | 0                               | 0                          |

|              |                |             |                  |
|--------------|----------------|-------------|------------------|
| RO122-BV     | <b>broiler</b> | 132         | 1195244          |
| RO321-BUC    | <b>broiler</b> | 0           | 0                |
| RO222-BZ     | <b>broiler</b> | 470         | 17204969         |
| RO312-CL     | <b>broiler</b> | 1017        | 19174974         |
| RO422-CS     | <b>broiler</b> | 47          | 576674           |
| RO113-CJ     | <b>broiler</b> | 214         | 5450000          |
| RO223-CT     | <b>broiler</b> | 276         | 5839200          |
| RO123-CV     | <b>broiler</b> | 26          | 437332           |
| RO313-DB     | <b>broiler</b> | 325         | 7500000          |
| RO411-DJ     | <b>broiler</b> | 66          | 2343729          |
| RO224-GL     | <b>broiler</b> | 16          | 256721           |
| RO314-GR     | <b>broiler</b> | 541         | 11128534         |
| RO412-GJ     | <b>broiler</b> | 132         | 3845190          |
| RO124-HR     | <b>broiler</b> | 2           | 32500            |
| RO423-HD     | <b>broiler</b> | 164         | 2210000          |
| RO315-IL     | <b>broiler</b> | 554         | 12246725         |
| RO213-IS     | <b>broiler</b> | 320         | 7318410          |
| RO322-IF     | <b>broiler</b> | 63          | 1117761          |
| RO114-MM     | <b>broiler</b> | 85          | 1754050          |
| RO413-MH     | <b>broiler</b> | 0           | 0                |
| RO125-MS     | <b>broiler</b> | 84          | 1512000          |
| RO214-NT     | <b>broiler</b> | 18          | 394971           |
| RO414-OT     | <b>broiler</b> | 0           | 0                |
| RO316-PH     | <b>broiler</b> | 143         | 1202620          |
| RO116-SJ     | <b>broiler</b> | 82          | 1870845          |
| RO115-SM     | <b>broiler</b> | 354         | 1600000          |
| RO126-SB     | <b>broiler</b> | 159         | 2347205          |
| RO215-SV     | <b>broiler</b> | 0           | 0                |
| RO317-TR     | <b>broiler</b> | 60          | 2160000          |
| RO424-TM     | <b>broiler</b> | 6           | 52620            |
| RO225-TL     | <b>broiler</b> | 0           | 0                |
| RO415-VL     | <b>broiler</b> | 221         | 2477768          |
| RO226-VN     | <b>broiler</b> | 160         | 298523           |
| RO216-VS     | <b>broiler</b> | 68          | 1042310          |
| <b>Total</b> | <b>broiler</b> | <b>6218</b> | <b>123515372</b> |

### **Sampling programmes**

The National Salmonella Control Programme encompasses the following serovars of zoonotic Salmonella: Salmonella enteritidis and Salmonella typhimurium .

The sampling programme will be in accordance to Regulation 2160/2003 EC and Regulation 646/2007 EC. We have also taken into account Regulation 1177/2006 EC and some amendments of Regulation 1168/2006 EC.

## 4. Measures of the submitted programme

### 4.1. Summary of measures under the programme

Year: 2013

- Control
- Testing
- Slaughter of positive animals
- Killing of positive animals
- Vaccination
- Treatment of animal products
- Disposal of products
- Monitoring or surveillance
- Other measure

### 4.2. Designation of the central authority charged with supervising and coordinating the departments responsible for implementing the programme:

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in flocks of broilers of *Gallus gallus* is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**

The departments responsible for implementing the National Control Programme of Salmonella in broilers are:

- for implementing at central level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary General Directorate** ;
- In respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the checking of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**.
- Concerning the slaughter of broilers flocks, the **Hygiene and Veterinary Epidemiology Directorate** is responsible for public health protection, through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.
- In respect of EC Regulation No **183/2005** on feed hygiene at the central level is **The Directorate for Technical Coordination of**

**Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**

- At county level for implementing the Regulation (EC) No **2160/2003**, **882/2004** and **183/2005** there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the National Reference Laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

**NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

**Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România

Tel: 0374.322.013 / 0374.322.000

Fax: 0214.113.394

E-mail: [office@idah.ro](mailto:office@idah.ro)

- **H.I.P.V.H.** (Hygiene Institute for Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

**NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED**

**Hygiene Institute for Sanitary Veterinary Public Health**

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201

Tel: 021.252.46.51 Fax: 021.252.00.61 E-mail: [iispv @ iispv.ro](mailto:iispv@iispv.ro)

The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL at February 2011 is show in the link below:

[http://www.ansvsa.ro/documente/admin/lista%20Lab%20Salm%20Site\\_11043ro.pdf](http://www.ansvsa.ro/documente/admin/lista%20Lab%20Salm%20Site_11043ro.pdf)

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the serotype according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

**1. N.S.V.F.S.A.**

- proposal of the plan of disease control.
- elaborate and submit to the EC the National Control Programme
- evaluate, coordinate the implementation of the National Control Programme in Romania and propose the update of this programme if is necessary (depending of the national situation)
- adoption of measures based on the disease situation in Romania
- submission of reports to the E.C.
- training the specialists from C.S.V.F.S.D

## **2. S.V.F.S.D.**(Sanitary Veterinary and Food Safety Direction).

- Coordination of the programme at the county level
- Official sampling is performed by the official veterinarian
- Propose the compensation costs to the Commission which will evaluate compensation documents.

**3. I.D.A.H.** Institute of Diagnosis and Animal Health is the National Reference Laboratory concerning animal health and was also designated as NRL for Salmonella in animals.

Responsibilities and tasks of the National Reference Laboratory for Salmonella (**I.D.A.H.**), pursuant to Directive 2003/99/EC and Regulation (EC) No 2160/2003, according to provision of N.S.V.F.S.A.. President Order no.160/2006 and 205/2007 are:

### **1. General duties**

- (a) To collaborate with the Community reference laboratory in their area of competence.
- (b) To coordinate, as appropriate, the activities of laboratories responsible for the analysis of samples in accordance with, in particular, Articles 4, 5 and 7 of Directive 2003/99/EC.
- (c) To coordinate the activities of laboratories responsible for the analysis of samples in accordance with Article 12(1) of Regulation (EC) No 2160/2003/EC.
- (d) Where appropriate, to organise comparative tests between the laboratories referred to under (b) and (c) and to assure an appropriate follow-up of such comparative testing.
- (e) To ensure the dissemination to the competent authority and to the laboratories referred to under (b) and (c), of the information that the Community reference laboratory supplies.

- (f) To provide scientific and technical assistance to the National Sanitary Veterinary and Food Safety Authority in their area of competence.
- (g) Characterize the pathogen isolates, genetic typing of this agents
- (h) Keep in maxim security conditions, the isolates
- (i) Give to the C.E., CRL, OMS, and national reference Laboratory of other member states, with the accord of N.S.V.F.S.A. all the information required.
- (j) train of the specialists from the government and private laboratory.

## **2. Specific functions and duties**

- (a) To participate, as appropriate in the monitoring and control programme for Salmonella and related anti-microbial resistance pursuant to Directive 2003/99/EC and in the analysis and testing of *Salmonella* pursuant to Regulation (EC) No 2160/2003.
- (b) To inform, as appropriate, the Community reference laboratory on aspects related to Salmonella vaccine strains and other specific control methods.
- (d) To gather data and information on the activities developed and methods used in relevant laboratories and to inform the Community reference laboratory thereof.
- (e) To monitor the epidemiological evolution of salmonella in Roumania.

## **5. S.V.F.S.L.**

There are 41 county Sanitary Veterinary and Food Safety Laboratories. Only 27 of them apply quality assurance systems that conform to the requirements of the current EN/ISO standard and are designated by National Reference Laboratory for Salmonella in animals to perform bacteriological examinations in the framework of the programme under the supervision of IDAH-NRL for Salmonella.

## **6. H.I.P.S.V.H.**

Hygiene Institute for Public Sanitary Veterinary Health is the National Reference Laboratory concerning the expertise for food products of animal origin and feed and it is also designed as the National Reference Laboratory for Salmonella in food and feed –public veterinary health.

### **4.3. Description and delimitation of the geographical and administrative areas in which the programme is to be implemented:**

The National Control Programme will be implemented throughout Romania, covering all the national territory and will cover all broiler flocks of *Gallus gallus* .

The administrative boundaries are the boundaries of the country. Roumania is administrative divided in 42 counties. There are 42 County Sanitary Veterinary and Food Safety Directorates and 41 County Sanitary veterinary and food Safety Laboratories.

### **4.4 Measures implemented under the programme**

#### **Bio-Security Measures**

Bio-security is a combination of practices, which are intended to prevent the spread of disease-causing organisms within the poultry farm. Where these are performed in parallel with the sanitation and disinfection procedures, bio-security measures could eradicate or, at least, reduce the level of pathogens to values, at which no hazard of infection would be likely.

The bio-security measures in commercial poultry farms are in accordance to the NSVFSA President Order 147/2006 regarding the bio-security measures in commercial poultry farms and the movements of live poultry, products and poultry by-products.

#### **Bio-security measures on holdings:**

- Health status of poultry
- On entering to all houses on the farm must be located disinfection barrier
- Control of movement of people
- Transport hygiene
- Feed hygiene
- Water hygiene
- Rodent, insect and bird control
- Cleaning and disinfecting of buildings
- Recording of all events and operations

According to the provision of Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program approved every year by N.S.V.F.S.A. President Order, after each cleaning and disinfections the owner is oblige to take samples to verify the efficient of the disinfection. The official veterinarian take sample to verify the efficiency of the disinfection only in case of a positive flock.

Hygiene measures on poultry farms are also assessed during visits for the collection of official samples and during general visits to premises for other purpose.

#### **4.4.1. Measures and terms of legislation as regards the registration of holdings:**

All commercial poultry flocks in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centres / holdings of origin and means of transport for health and animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products. Poultry holdings shall be registered and sanitary veterinary approved in order to be able to operate.

The legal framework for carrying out commercial activity with poultry and poultry products consists of:

- Order of the President of the National Sanitary Veterinary and Food Safety **Authority no 144/2006** for the approval of the Sanitary veterinary norm on animal health conditions governing intra-Community trade in, and imports from third countries of, poultry and hatching eggs transposing Council Directive 90/539/EEC on animal health conditions governing intra-Community trade in, and imports from third countries of poultry and hatching eggs.

In order to be sanitary veterinary approved, poultry holdings shall satisfy, in accordance with the legislation in force, the following conditions:

- Appropriate facilities and operation;
- Application of the “Program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” approved by Order of the National Sanitary Veterinary and Food Safety Authority President .
- At least one inspection visit per year by the official veterinarian;
- Additional checks to verify the compliance of the establishment with the hygiene measures and the operation of the establishments.

Each poultry holding receives a distinct approval number, number that can be the same with the one given in compliance with the **Council Regulation no. 2782/75/EEC** on the production and marketing of eggs for hatching and of farmyard poultry chicks.

#### **4.4.2.Measures and terms of legislation as regards the identification of animals:** Not applicable.

#### **4.4.3. Measures and terms of legislation as regards the notification of the disease:**



**Commission Decision 176/2005** of 1 March 2005 laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2006/924/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2010/160/EU** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2008/755/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2009/847/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

The internal notification is made under the provision of **N.S.V.F.S.A. President ORDER no. 79 /2008** to approve the sanitary and veterinary norm on domestic notification and official declaration of animal diseases.

#### **4.4.4. Measures and terms of legislation as regards the measures in case of a positive result:**

In case of suspicion or conformation of ***Salmonella enteritidis*** or ***Salmonella typhimurium*** the NRL shall notify immediately the N.S.V.F.S. and local C.S.V.F.S.D..

In case of suspicion of infection the local C.S.V.F.S.D. and the relevant authorities:

- prohibited the movement of broilers
- take additional samples for conformation of infection

When the broilers are confirmed for the presence of ***Salmonella enteritidis*** or ***Salmonella typhimurium***:

1. Fresh meat from broilers may be placed on the market on the condition that it meets the requirement of absence of Salmonella in 25 grams from the meat.
2. The requirement laid down in point 1 does not apply to fresh poultry meat destined for heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene.
3. The criterion laid down in point 1 does not apply to fresh poultry meat destined for industrial heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene

When a broiler flock of *Gallus gallus* is suspected of being infected with *Salmonella Enteritidis* or *Salmonella Typhimurium* the flock will be investigated. The flock is suspected of being infected when *S. Enteritidis* or *S. Typhimurium* is isolated from a sample of faeces, or boot swabs, carried out privately or as required by either the operator or the Competent Authority as detailed in the Annex to Regulation (EC) No 646/2007. Tissue/organs may be taken from birds as part of the investigation of clinical disease by the veterinarian; these cases will be discussed and additional follow up investigation carried out as appropriate, along with advice on *Salmonella* control.

Competent Authority will notify the operator to clean and disinfect the building from which the infected flock originated. After cleaning and disinfecting of the building the operator may be required to take swabs from a number of sites in the building and submit them to an approved laboratory in view to be tested for *Salmonella* in order to check the efficiency of the hygiene measures taken. In cases where *S. Enteritidis* or *S. Typhimurium* was isolated the cleaning and disinfection may be checked by the Competent Authority or its agent.

If the results of post-cleaning and disinfection monitoring of *Salmonella* are positive for *S. Enteritidis* or *S. Typhimurium*, the next crop (cycle) will be monitored under supervision of the Competent Authority or its agent. If *Salmonella* is isolated in this subsequent crop of birds the holding will be placed under official control; re-stocking of the house will be permitted only if the supervised post-cleaning and disinfection samples from the house are negative.

For the purposes of establishing the progress towards the target if *S. Enteritidis* or *S. Typhimurium* is isolated from either an operator sample or an official sample the flock is classed as positive. A flock positive for a specific serotype will be recorded only once for that serotype.

Operators with a flock which is positive for *S. Enteritidis* or *S. Typhimurium* will be contacted by the Competent Authority for advice on how to reduce or eliminate the *Salmonella*. Advice on the control of *Salmonella* in broilers will be available from government experts on *Salmonella* control. Advice may include recommendations on management, cleaning and disinfection, pest control, biosecurity, monitoring, and the potential use of other aids in the control of *Salmonella*.

In further future a Code of Practice for the control of *Salmonella* on broiler farms will be available.

### **Control of the use of feed antibiotics by official sampling**

According to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order, feeding

stuffs intended for poultry nutrition are checked in view to avoid the contamination with *Salmonella* spp. Also, in conformity with the same legislation the feed stuffs are checked in view to detect the use of antibiotics.

Residues examination is performed according to the Roumanian annual plan for examination for residues in live animals and animal origin products. For broiler, hens, turkeys, other poultry a sample consists on one or more animals depending on the requirements of the analytical methods.

For each category of poultry considered, the minimum number of samples to be taken each year must be at least equal to one per 200 tones of annual production, with a minimum of 100 samples for each group of substances if the annual production of the category of birds considered is over 5 000 tones.

The following breakdown must be respected:

**Group A:** 50 % of the total samples. The equivalent of one fifth of these samples must be taken at farm level. Each sub-group of Group A must be checked each year using a minimum of 5 % of the total number of samples to be collected for Group A. The balance will be allocated according to the experience and background information of Roumania.

**Group B:** 50 % of the total samples, 30 % of samples must be checked for Group B 1 substances (antibiotics and sulfamides)

**4.4.5. Measures and terms of legislation as regards the different qualifications of animals and herds:** Not applicable.

**4.4.6. Control procedures and in particular rules on the movement of animals liable to be affected or contaminated by a given disease and the regular inspection of the holdings or areas concerned.**

According to the provisions of N.S.V.F.S.A. President Order 147/2006, Regulation 2160/2003/EC, the following measures are to be adopted in order to prevent the dissemination of *Salmonella enteritidis*, *Salmonella typhimurium*, into commercial holdings. Animals from infected flocks belonging to commercial holdings are to be kept isolated and special conditions apply for removal of these animals. No bird may leave the house concerned unless the competent authority has authorized the slaughter or/and destruction under supervision of slaughter in a slaughterhouse designated by the competent authority. All the birds in the house must be slaughtered in accordance with the provisions of the **REGULATION (EC) No. 853/2004 laying down specific hygiene rules for food of animal origin in order** to reduce as much as possible the risk of spreading salmonella.

#### 4.4.7. Measures and terms of legislation as regards the control (testing, vaccination ...) of the disease.

The legal basis is the provisions of the **Regulation 2160/2003/EC**, **Regulation 646/2007/EC** and **Regulation 1177/2006 EC**.

According to the provision of Regulation (EC) no. 1177/2006 CE in Roumania the antimicrobials use for Salmonella control is forbidden.

A national *Salmonella* control programme was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 646/2007. The national control plan for *Salmonella* broilers come into effect in January 2009.

***The NCP defines a flock as a single group or multiple groups of chickens which share the same production unit (i.e. using the same air-space or range area). Where housing systems are not typical, the situation is likely to be assessed on a case by case basis. Multiple groups of chickens which have 'beak-to-beak' contact (inside or outside the house) are likely to be treated as a single flock.***

All flocks of broiler will be included in the national control programme unless exempted in Regulation (EC) No 2160/2003 under Article 1.3, i.e. birds produced for private domestic consumption, or where there is direct supply of small quantities of products to the final consumer or to local retail establishments directly supplying the primary products to the final consumer.

**Operators** will be required to implement the sampling programme in the Annex to EC Regulation 646/2007 (**self-control sampling**). For convenience the 'Sampling protocol' is repeated in **THE SAMPLING PROTOCOL**, showed below . Two pairs of boot sock/swabs will be taken by the operator within the period of **three weeks** before the birds are due for slaughter. The samples will be taken in sufficient time for the laboratory results to be known before the birds are transported to the slaughter house. It is important to know the *Salmonella* status of the flock before the first birds are slaughtered. Samples will be submitted to a laboratory authorised by the Competent Authority and which applies quality assurance systems that conform to the requirements of the current EN/ISO standard.

Each year at least 10% of holdings with more than 5,000 birds will be selected and at least one flock on the holding will be sampled by Animal Health, or other authorized agent, acting on behalf of the Competent Authority, who will take an '**official sample**'. In addition, attention will also be given to flocks where there have been previously positive *Salmonella* findings in the samples taken by the operators. Particular attention will be given to holdings where *S. Enteritidis* or *S. Typhimurium* has been isolated from samples. Sampling to verify the achievement of the target will be as detailed in 'Sampling protocol' in the Annex to Commission Regulation (EC) No 646/2007 and the procedure for implementing the Salmonella

National Control Programme in broilers flocks, issued by the Sanitary Veterinary General Directorate .

When an official sample is taken it may replace the sample required to be taken by the operator.

In accordance with Regulation (EC) No. 646/2007 Annex point 1 (c) the operator of a broiler holding may make an application to the Competent Authority for a derogation not to sample all flocks on the holding. The Competent Authority will assess the application for derogation against the criteria listed in the Annex -

(i) an all in/all out system is used;

(ii) the same management applies to all flocks;

(iii) feed and water supply is common to all flocks;

(iv) during one year and at least six crops of flocks, samples were taken for and tested for *Salmonella* spp according to the standard monitoring scheme in all flocks on the holding. Samples of all flocks of at least one crop were taken by the Competent Authority; and no SE or ST was found in any sample from this period of enhanced testing

The Competent Authority may approve the derogation if satisfied.

## **Sampling protocol.**

For each flock\*

At least two pairs of boot/sock swabs shall be taken. All boot/sock swabs must be pooled into one sample. For free range broiler flocks, samples shall only be collected in the area inside the house.

Before using the boot/sock swabs, their surface shall be moistened with deionised water, or sterile water or any other diluents approved by the national reference laboratory referred to in Article 11 of Regulation (EC) No 2160/2003. The use of farm water containing antimicrobials or additional disinfectants shall be prohibited. The recommended way to moisten boot swabs shall be to pour the liquid inside before putting them on. It shall be ensured that all sections in a house are represented in the sampling in a proportionate way and that at least 100 steps are taken with each pair of boot swabs. Each pair should cover about 50 % of the area of the house.

On completion of sampling the boot/sock swabs shall be carefully removed so as not to dislodge adherent material. Boot swabs may be inverted to retain material. They shall be placed in a bag or pot and labelled to identify the flock sampled, and the date the samples were taken.

## **Transport and preparation of the samples:**

Samples shall be sent by express mail or courier to the approved laboratories **within 25 hours after collection**. At the laboratory samples shall be kept refrigerated until examination, which shall be carried out **within 48 hours** following receipt. If they are tested after 48 hours the results are invalidate.

The pair of boot/sock swabs shall be carefully unpacked to avoid dislodging adherent faecal material, pooled and placed in 225 ml buffered peptone water (BPW) which has been pre-warmed to room temperature. The sample shall be swirled to fully saturate it and culture shall be continued by using the detection method, described below.

### **Detection method:**

The Salmonella spp. detection from faeces of animals made by the NRL for Salmonella in animals or C.S.V.F.S.L. is made according to PSO 001 - "Isolation and identification of the Salmonellosis" done by RENAR in 2006 (accreditation certificate no. 222 L/2006) and revised in May 2009. The procedure is in accordance with SR EN ISO 6579:2003 / A1: 2007 and with the standard procedure of CRL - Salmonella, Bilthoven, Netherlands.

In that detection method, a semi-solid medium (modified semi-solid Rappaport-Vassiladis medium, MSRV) is used as the single selective enrichment medium.

### **Serotyping**

At least one isolate from each positive sample shall be serotyped, following the Kaufmann-White scheme.

### **Storage of strains:**

At least one isolated strain per house and per year shall be collected by the NRL and stored for future phagotyping or anti-microbial susceptibility testing, using the normal methods for culture collection, which must ensure integrity of the strains for a minimum of two years.

### **Use of vaccines**

Live Salmonella vaccines shall not be used in the framework of national control programme where the manufacturer does not provide an appropriate method to distinguish bacteriological wild-type strains of salmonella from vaccine strains.

Live salmonella vaccines shall not be used in the framework of national control programmes in laying hens during production unless the safety of the use has been demonstrated and they are authorised for such purpose in accordance with Directive 2001/82/EC.

Although **vaccines against *Salmonella*** are not currently used in **broilers**, this practice may change in the future if suitable vaccines are developed.

#### **4.4.8. Measures and terms of legislation as regards the compensation for owners of slaughtered and killed animals:**

In broilers flocks of *Gallus gallus* financial assistance is provided in the context of the control programme. This is set up by the **Guverment Decision no. 1214/2009** regarding the methodology for determining and paying compensation to be paid to owners of slaughtered animals, killed or otherwise affected by the rapid liquidation of outbreaks of animal diseases.

#### **4.4.9. Information and assessment on bio-security measures management and infrastructure in place in the flocks/holdings involved**

This information was given in the point 4.4.

## **5. General description of the benefits:**

### **Costs of the programme**

This programme is in accordance with Council Decision 2009/470/ EEC, Commission Decision 2008/425/EC and Council Decision 92/65/ EEC. The detailed financial costs for the national control programme of Salmonella in laying hens are laid down in point 8. The detailed financial costs for the national control programme of Salmonella in broiler are laid down in point 8.

### **Benefits of the programme**

The overall aim of the Salmonella National Control Programme is to control the occurrence of Salmonella in the poultry sector on a very low level and thereby protect humans against infection with food-borne salmonellas. It has been known that poultry often harbour latent infections with Salmonella, which may pose a serious human health risk.

The anticipated benefits of this programme reducing the *Salmonella enteritidis* and *Salmonella typhimurium* are the minimising of human health problems and a consequent reduction in suffering, mortality and health service costs.

## **6. Data on the epidemiological evolution during the last years:**



## 6.1.Evolution of zoonotic salmonellosis

### 6.1.1 Data on evolution of zoonotic salmonellosis

Year: 2009

Situation on date: 31.12.2009

Animal species: broiler (Gallus gallus)

Disease: zoonotic salmonellosis

| Region<br>(County+<br>NUTS<br>CODE) | Type of<br>flocks(b) | Total<br>number<br>of<br>flocks © | Total<br>number of<br>animals | Total<br>number of<br>flocks<br>under the<br>programme | Total<br>number of<br>animals<br>under the<br>programme | Number of<br>flocks<br>checked(d) | Number of positive<br>flocks (e)                            |                           | Number of flocks<br>depopulated                             |                           | Total number of animals<br>slaughtered or<br>destroyed      |                           | Quantity of the<br>eggs destroyed<br>(number or kg)         |                                       | Quantity of eggs<br>channelled to egg products<br>(number or kg) |                       |
|-------------------------------------|----------------------|-----------------------------------|-------------------------------|--|---|-----------------------------------|---|---------------------------|---|---------------------------|---|---------------------------|---|---------------------------------------|--|-----------------------|
|                                     |                      |                                   |                               |  |   |                                   | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>serotyp<br>es(g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>seroty<br>pes(g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>serotypes<br>(g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Oth<br>er<br>ser<br>oty<br>pes(<br>g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f)      | Other<br>serotypes(g) |
| RO121-AB                            | BROILER              | 77                                | 1150000                       | 77   | 1150000   | 77                                | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO421-AR                            | BROILER              | 0                                 | 0                             | 0  | 0   | 0                                 | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO311-AG                            | BROILER              | 151                               | 7945670                       | 151  | 7945670   | 151                               | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO211-BC                            | BROILER              | 682                               | 15710000                      | 682  | 15710000  | 682                               | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO111-BH                            | BROILER              | 0                                 | 0                             | 0  | 0   | 0                                 | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO112-BN                            | BROILER              | 7                                 | 104700                        | 7  | 104700  | 7                                 | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO212-BT                            | BROILER              | 6                                 | 148780                        | 6  | 148780  | 6                                 | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO221-BR                            | BROILER              | 0                                 | 0                             | 0  | 0   | 0                                 | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO122-BV                            | BROILER              | 173                               | 1602249                       | 173  | 1602249   | 173                               | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO321-BUC                           | BROILER              | 0                                 | 0                             | 0  | 0   | 0                                 | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO222-BZ                            | BROILER              | 438                               | 12845086                      | 438  | 12845086  | 438                               | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO312-CL                            | BROILER              | 227                               | 24000000                      | 227  | 24000000  | 22                                | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO422-CS                            | BROILER              | 29                                | 4539000                       | 29   | 4539000   | 29                                | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO113-CJ                            | BROILER              | 51                                | 815000                        | 51   | 815000  | 15                                | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO223-CT                            | BROILER              | 43                                | 906000                        | 39   | 870000  | 32                                |   | 1 (Senftenberg)           | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO123-CV                            | BROILER              | 26                                | 191000                        | 26   | 191000  | 26                                | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO313-DB                            | BROILER              | 84                                | 10540000                      | 84   | 10540000  | 84                                | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |
| RO411-DJ                            | BROILER              | 0                                 | 0                             | 0  | 0   | 0                                 | 0   | 0                         | 0   | 0                         | 0   | 0                         | 0   | 0                                     | 0  | 0                     |

Revizie : 24.08.2012

|              |                |             |                  |             |                  |             |          |              |          |            |          |                |          |          |          |          |
|--------------|----------------|-------------|------------------|-------------|------------------|-------------|----------|--------------|----------|------------|----------|----------------|----------|----------|----------|----------|
| RO224-GL     | BROILER        | 6           | 417491           | 6           | 413210           | 6           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO314-GR     | BROILER        | 523         | 9676221          | 523         | 9676221          | 523         | 0        | 148          | 0        | 148        | 0        | 3026563        | 0        | 0        | 0        | 0        |
| RO412-GJ     | BROILER        | 36          | 132332           | 36          | 132332           | 36          | 0        | 1(Infant is) | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO124-HR     | BROILER        | 6           | 25500            | 6           | 25500            | 6           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO423-HD     | BROILER        | 170         | 2140000          | 170         | 2140000          | 170         | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO315-IL     | BROILER        | 412         | 10605550         | 412         | 10605550         | 21          | 2        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO213-IS     | BROILER        | 90          | 10340955         | 90          | 10340955         | 8           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO322-IF     | BROILER        | 0           | 0                | 0           | 0                | 0           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO114-MM     | BROILER        | 19          | 312245           | 19          | 312245           | 19          | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO413-MH     | BROILER        | 0           | 0                | 0           | 0                | 0           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO125-MS     | BROILER        | 12          | 960000           | 12          | 960000           | 12          | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO214-NT     | BROILER        | 1           | 20000            | 1           | 20000            | 1           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO414-OT     | BROILER        | 2           | 60000            | 2           | 60000            | 2           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO316-PH     | BROILER        | 136         | 917822           | 109         | 917822           | 109         | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO116-SJ     | BROILER        | 3           | 322002           | 3           | 294575           | 2           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO115-SM     | BROILER        | 57          | 1500000          | 57          | 1500000          | 57          | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO126-SB     | BROILER        | 157         | 2440000          | 157         | 2440000          | 157         | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO215-SV     | BROILER        | 0           | 0                | 0           | 0                | 0           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO317-TR     | BROILER        | 12          | 1800000          | 12          | 1800000          | 12          | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO424-TM     | BROILER        | 11          | 558500           | 11          | 558500           | 11          | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO225-TL     | BROILER        | 0           | 0                | 0           | 0                | 0           | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO415-VL     | BROILER        | 196         | 1897793          | 196         | 1897793          | 196         | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO226-VN     | BROILER        | 195         | 3362050          | 195         | 3362050          | 22          | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| RO216-VS     | BROILER        | 70          | 628943           | 70          | 628943           | 48          | 0        | 0            | 0        | 0          | 0        | 0              | 0        | 0        | 0        | 0        |
| <b>Total</b> | <b>BROILER</b> | <b>4108</b> | <b>128614889</b> | <b>4077</b> | <b>128547181</b> | <b>3160</b> | <b>2</b> | <b>150</b>   | <b>0</b> | <b>148</b> | <b>0</b> | <b>3026563</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> |

Year: 2010

Situation on date: 31.12.2010

Animal species: broiler (Gallus gallus)

Disease: zoonotic salmonellosis

| Region<br>(County+N<br>UTS CODE) | Type of<br>flocks(b) | Total<br>number<br>of flocks<br>© | Total number<br>of animals | Total<br>number of<br>flocks<br>under the<br>programme | Total number<br>of animals<br>under the<br>programme | Number<br>of flocks<br>official<br>checked(<br>d) | Number of positive<br>flocks (e)                                     |                           | Number of flocks<br>depopulated                                      |                                   | Total number of animals<br>slaughtered                               |                           | Quantity of the eggs<br>destroyed (number or kg)             |                       | Quantity of eggs channelled<br>to egg products (number or<br>kg) |                       |
|----------------------------------|----------------------|-----------------------------------|----------------------------|--|--|---|--|---------------------------|--|-----------------------------------|--|---------------------------|--|-----------------------|--|-----------------------|
|                                  |                      |                                   |                            |  |  |   | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Othe<br>r<br>serot<br>ypes(<br>g) | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotypes(<br>g) | Serotypes<br>targeted in<br>the control<br>programm<br>e (f) | Other<br>serotypes(g) | Serotypes<br>targeted in<br>the control<br>programm<br>e (f)     | Other<br>serotypes(g) |
| RO121-AB                         | broiler              | 83                                | 958000                     | 83   | 958000   | 83  | 0  | 13                        | 0  | 13                                | 0  | 150048                    | 0  | 0                     | 0  | 0                     |
| RO421-AR                         | broiler              | 20                                | 400000                     | 20   | 400000   | 4   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO311-AG                         | broiler              | 28                                | 705510                     | 28   | 705510   | 0   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO211-BC                         | broiler              | 486                               | 3288000                    | 486  | 3288000  | 48  | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO111-BH                         | broiler              | 41                                | 386000                     | 40   | 378000   | 9   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO112-BN                         | broiler              | 1                                 | 2000                       | 1  | 2000   | 1   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO212-BT                         | broiler              | 81                                | 364030                     | 81   | 364030   | 45  | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO221-BR                         | broiler              | 0                                 | 0                          | 0  | 0  | 0   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO122-BV                         | broiler              | 743                               | 13474060                   | 743  | 13474060   | 173   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO321-BUC                        | broiler              | 0                                 | 0                          | 0  | 0  | 0   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO222-BZ                         | broiler              | 410                               | 10216000                   | 410  | 10216000   | 410   | 1  | 29                        | 1  | 29                                | 13754  | 833855                    | 0  | 0                     | 0  | 0                     |
| RO312-CL                         | broiler              | 850                               | 32000000                   | 850  | 32000000   | 10  | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO422-CS                         | broiler              | 125                               | 4476472                    | 124  | 4476472  | 3   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO113-CJ                         | broiler              | 255                               | 3260000                    | 255  | 3260000  | 51  | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO223-CT                         | broiler              | 246                               | 5813400                    | 41   | 968900   | 39  | 0  | 16                        | 0  | 16                                | 0  | 92163                     | 0  | 0                     | 0  | 0                     |
| RO123-CV                         | broiler              | 104                               | 4476472                    | 6  | 258258   | 6   | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |
| RO313-DB                         | broiler              | 84                                | 1046105                    | 84   | 1046105  | 84  | 0  | 1                         | 0  | 1                                 | 0  | 12453                     | 0  | 0                     | 0  | 0                     |
| RO411-DJ                         | broiler              | 114                               | 2799355                    | 60   | 1186783  | 16  | 0  | 0                         | 0  | 0                                 | 0  | 0                         | 0  | 0                     | 0  | 0                     |

Revizie : 24.08.2012

|              |                |             |                  |             |                  |             |          |            |          |            |              |                |          |          |          |          |
|--------------|----------------|-------------|------------------|-------------|------------------|-------------|----------|------------|----------|------------|--------------|----------------|----------|----------|----------|----------|
| RO224-GL     | broiler        | 38          | 396000           | 6           | 9200             | 0           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO314-GR     | broiler        | 530         | 9342931          | 83          | 1502218          | 83          | 0        | 26         | 0        | 26         | 0            | 466598         | 0        | 0        | 0        | 0        |
| RO412-GJ     | broiler        | 132         | 3845190          | 132         | 3845190          | 28          | 1        | 6          | 1        | 6          | 29130        | 174781         | 0        | 0        | 0        | 0        |
| RO124-HR     | broiler        | 1           | 35000            | 0           | 0                | 0           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO423-HD     | broiler        | 63          | 610000           | 63          | 610000           | 6           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO315-IL     | broiler        | 492         | 11795885         | 492         | 11795885         | 58          | 0        | 15         | 0        | 15         | 0            | 359630         | 0        | 0        | 0        | 0        |
| RO213-IS     | broiler        | 480         | 9254678          | 200         | 7895254          | 92          | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO322-IF     | broiler        | 48          | 749825           | 44          | 731325           | 2           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO114-MM     | broiler        | 19          | 230137           | 19          | 230137           | 19          | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO413-MH     | broiler        | 0           | 0                | 0           | 0                | 0           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO125-MS     | broiler        | 78          | 1400000          | 78          | 1400000          | 14          | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO214-NT     | broiler        | 0           | 0                | 0           | 0                | 0           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO414-OT     | broiler        | 0           | 0                | 0           | 0                | 0           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO316-PH     | broiler        | 520         | 1395869          | 504         | 1385869          | 8           | 1        | 7          | 1        | 7          | 36067        | 259433         | 0        | 0        | 0        | 0        |
| RO116-SJ     | broiler        | 53          | 1055500          | 53          | 1055500          | 37          | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO115-SM     | broiler        | 336         | 1280000          | 336         | 1280000          | 56          | 1        | 3          | 1        | 3          | 3810         | 12429          | 0        | 0        | 0        | 0        |
| RO126-SB     | broiler        | 156         | 2365050          | 156         | 2365050          | 156         | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO215-SV     | broiler        | 0           | 0                | 0           | 0                | 0           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO317-TR     | broiler        | 60          | 245000           | 60          | 245000           | 2           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO424-TM     | broiler        | 20          | 188000           | 20          | 188000           | 2           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO225-TL     | broiler        | 0           | 0                | 0           | 0                | 0           | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO415-VL     | broiler        | 213         | 4491151          | 213         | 4491151          | 43          | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO226-VN     | broiler        | 199         | 372076           | 199         | 372076           | 43          | 0        | 0          | 0        | 0          | 0            | 0              | 0        | 0        | 0        | 0        |
| RO216-VS     | broiler        | 70          | 1157345          | 70          | 1157345          | 22          | 0        | 8          | 0        | 8          | 0            | 132268         | 0        | 0        | 0        | 0        |
| <b>Total</b> | <b>broiler</b> | <b>7179</b> | <b>133875041</b> | <b>6040</b> | <b>113541318</b> | <b>1653</b> | <b>4</b> | <b>124</b> | <b>4</b> | <b>124</b> | <b>82761</b> | <b>2493658</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> |

Revizie : 24.08.2012

Year: 2011

Situation on date: 31.12.2011

Animal species: broiler (Gallus gallus)

Disease: zoonotic salmonellosis

| Region<br>(County+NUTS<br>CODE) | Type of<br>flocks(<br>b) | Total<br>number<br>of<br>flocks<br>© | Total number<br>of animals | Total<br>number of<br>flocks<br>under the<br>programm<br>e | Total number<br>of animals<br>under the<br>programme | Number<br>of<br>flocks<br>checke<br>d(d) | Number of positive<br>flocks (e)                                     |                           | Number of<br>flocks<br>depopulated                                   |                                   | Total number of animals<br>slaughtered or<br>destroyed          |                           | Quantity of the eggs<br>destroyed<br>(number or kg)             |                           | Quantity of eggs<br>channelled to egg products<br>(number or kg) |                       |
|---------------------------------|--------------------------|--------------------------------------|----------------------------|--|--|--|--|---------------------------|--|-----------------------------------|---|---------------------------|---|---------------------------|--|-----------------------|
|                                 |                          |                                      |                            |  |  |  | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Othe<br>r<br>seroty<br>pes<br>(g) | Serotypes<br>targeted in<br>the<br>control<br>programm<br>e (f) | Other<br>serotypes(<br>g) | Serotypes<br>targeted in<br>the<br>control<br>programm<br>e (f) | Other<br>serotypes(<br>g) | Serotypes<br>targeted in<br>the<br>control<br>programm<br>e (f)  | Other<br>serotypes(g) |
| RO121-AB                        | Broiler                  | 77                                   | 954700                     | 74   | 954700   | 74                                       | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO421-AR                        | Broiler                  | 8                                    | 112000                     | 8  | 112000   | 2  | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO311-AG                        | Broiler                  | 62                                   | 989460                     | 59   | 944941   | 8  | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO211-BC                        | Broiler                  | 127                                  | 3140803                    | 127  | 3722543  | 30                                       | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO111-BH                        | Broiler                  | 88                                   | 3002534                    | 88   | 3002534  | 37                                       | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO112-BN                        | Broiler                  | 18                                   | 225000                     | 18   | 225000   | 2  | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO212-BT                        | Broiler                  | 33                                   | 500000                     | 29   | 439394   | 13                                       | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO221-BR                        | Broiler                  | 0                                    | 0                          | 0  | 0  | 0  | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO122-BV                        | Broiler                  | 132                                  | 1195244                    | 132  | 1195244  | 132                                      | 1  | 22                        | 1  | 22                                | 11650   | 315360                    | 0   | 0                         | 0  | 0                     |
| RO321-BUC                       | Broiler                  | 0                                    | 0                          | 0  | 0  | 0  | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO222-BZ                        | Broiler                  | 470                                  | 17204969                   | 470  | 17204969   | 470                                      | 1  | 26                        | 1  | 26                                | 14400   | 470342                    | 0   | 0                         | 0  | 0                     |
| RO312-CL                        | Broiler                  | 1017                                 | 19174974                   | 32   | 2554822  | 32                                       | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO422-CS                        | Broiler                  | 47                                   | 576674                     | 47   | 576674   | 24                                       | 2  | 1                         | 2  | 1                                 | 245340  | 12270                     | 0   | 0                         | 0  | 0                     |
| RO113-CJ                        | Broiler                  | 214                                  | 5450000                    | 214  | 5450000  | 80                                       | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO223-CT                        | Broiler                  | 276                                  | 5839200                    | 46   | 973200   | 46                                       | 0  | 13                        | 0  | 13                                | 0   | 275034                    | 0   | 0                         | 0  | 0                     |
| RO123-CV                        | Broiler                  | 26                                   | 437332                     | 26   | 437332   | 26                                       | 2  | 0                         | 2  | 0                                 | 34000   | 0                         | 0   | 0                         | 0  | 0                     |
| RO313-DB                        | Broiler                  | 325                                  | 7500000                    | 325  | 1500000  | 42                                       | 0  | 2                         | 0  | 2                                 | 0   | 9250                      | 0   | 0                         | 0  | 0                     |
| RO411-DJ                        | Broiler                  | 66                                   | 2343729                    | 6  | 214000   | 4  | 0  | 0                         | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |
| RO224-GL                        | Broiler                  | 16                                   | 256721                     | 16   | 256721   | 4  | 0  |                           | 0  | 0                                 | 0   | 0                         | 0   | 0                         | 0  | 0                     |

Revizie : 24.08.2012

|              |                |             |                  |             |                 |             |           |            |           |            |               |                |          |          |          |          |
|--------------|----------------|-------------|------------------|-------------|-----------------|-------------|-----------|------------|-----------|------------|---------------|----------------|----------|----------|----------|----------|
| RO314-GR     | Broiler        | 541         | 11128534         | 541         | 11128534        | 90          | 0         | 166        | 0         | 166        | 0             | 3096375        | 0        | 0        | 0        | 0        |
| RO412-GJ     | Broiler        | 132         | 3845190          | 132         | 3845190         | 7           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO124-HR     | Broiler        | 2           | 32500            | 2           | 32500           | 0           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO423-HD     | Broiler        | 164         | 2210000          | 164         | 180970          | 17          | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO315-IL     | Broiler        | 554         | 12246725         | 554         | 12246725        | 34          | 0         | 22         | 0         | 22         | 0             | 486332         | 0        | 0        | 0        | 0        |
| RO213-IS     | Broiler        | 320         | 7318410          | 320         | 7318410         | 38          | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO322-IF     | Broiler        | 63          | 1117761          | 63          | 1117761         | 10          | 0         | 1          | 0         | 1          | 0             | 17743          | 0        | 0        | 0        | 0        |
| RO114-MM     | Broiler        | 85          | 1754050          | 85          | 1754050         | 16          | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO413-MH     | Broiler        | 0           | 0                | 0           | 0               | 0           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO125-MS     | Broiler        | 84          | 1512000          | 84          | 1512000         | 14          | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO214-NT     | Broiler        | 18          | 394971           | 1           | 22000           | 1           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO414-OT     | Broiler        | 0           | 0                | 0           | 0               | 0           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO316-PH     | Broiler        | 143         | 1202620          | 137         | 1102640         | 108         | 1         | 24         | 1         | 24         | 37030         | 395649         | 0        | 0        | 0        | 0        |
| RO116-SJ     | Broiler        | 82          | 1870845          | 82          | 1870845         | 22          | 2         | 2          | 2         | 2          | 36000         | 36919          | 0        | 0        | 0        | 0        |
| RO115-SM     | Broiler        | 354         | 1600000          | 354         | 1600000         | 61          | 2         | 10         | 2         | 10         | 70000         | 300000         | 0        | 0        | 0        | 0        |
| RO126-SB     | Broiler        | 159         | 2347205          | 159         | 2347205         | 27          | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO215-SV     | Broiler        | 0           | 0                | 0           | 0               | 0           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO317-TR     | Broiler        | 60          | 2160000          | 60          | 2160000         | 6           | 0         | 34         | 0         | 34         | 0             | 1020000        | 0        | 0        | 0        | 0        |
| RO424-TM     | Broiler        | 6           | 52620            | 6           | 52620           | 0           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO225-TL     | Broiler        | 0           | 0                | 0           | 0               | 0           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO415-VL     | Broiler        | 221         | 2477768          | 221         | 2477768         | 26          | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO226-VN     | Broiler        | 160         | 298523           | 160         | 298523          | 25          | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| RO216-VS     | Broiler        | 68          | 1042310          | 68          | 1042310         | 7           | 0         | 7          | 0         | 7          | 0             | 107300         | 0        | 0        | 0        | 0        |
| IDAHA        | Broiler        | 0           | 0                | 0           | 0               | 0           | 0         | 0          | 0         | 0          | 0             | 0              | 0        | 0        | 0        | 0        |
| <b>Total</b> | <b>Broiler</b> | <b>6218</b> | <b>123515372</b> | <b>4910</b> | <b>91874125</b> | <b>1535</b> | <b>11</b> | <b>330</b> | <b>11</b> | <b>330</b> | <b>448420</b> | <b>6542574</b> | <b>0</b> | <b>0</b> | <b>0</b> | <b>0</b> |

Revizie : 24.08.2012

## 6.4 Data on vaccination programme

Not applicable

## 7. Targets

### 7.1. Targets related to testing :to investigate the presence of Salmonella in broiler flocks

Disease (a): zoonotic Salmonella

Animal species:

| Region (b)  | Type of the test (c) | Target population (d) | Type of sample (e)     | Objective (f) | Number of planned tests |
|-------------|----------------------|-----------------------|------------------------|---------------|-------------------------|
| All regions | Bacteriological      | Broiler               | Faeces boot/sock swabs | Surveillance  | 2000                    |
|             | Bacteriological      | Broiler               | Faeces boot/sock swabs | Confirmation  | 500                     |
| Total       | -                    | -                     | -                      | -             | 2500                    |

(a) Disease and species if necessary

(b) Region as defined in the approved eradication programme of the Member State

(c) Description of the test

(d) Specification of the targeted species and the categories of targeted animals

(e) Description of the sample (faeces)

(f) Description of the objective ( e.g. qualification, surveillance, confirmation of suspected cases, monitoring of campaigns, seroconversion, control on deleted vaccines, testing of vaccine , control of vaccination

Revizie : 24.08.2012

## 7.1.2. Targets on testing of broiler flocks

Year: 2013

Animal species: Gallus gallus

Disease/infection(a): Zoonotic Salmonella

| Region      | Type of flock <sup>(b)</sup> | Total number of flocks <sup>(c)</sup> | Total number of animals | Total n° of flocks under the programme | Total n° of animals under the programme | Expected n° of flocks to be checked <sup>(d)</sup> | Number of flocks <sup>(d)</sup> expected to be positive <sup>(a)</sup> |      |      | Number of flocks expected to be depopulated |      | Total n° of animals expected to be slaughtered or destroyed <sup>(f)</sup> |      | Expected quantity of eggs to be destroyed (number or kg) <sup>(d)</sup> |    | Expected quantity of eggs channelled to egg products (number or kg.) <sup>(d)</sup> |      |
|-------------|------------------------------|---------------------------------------|-------------------------|--|---|--|--|------|------|---|------|--|------|---|----|---|------|
|             |                              |                                       |                         |  |   |  | (a1)   | (a2) | (a3) | a4  | (a3) | (a4)   | (a3) | (a4)  | a3 | (a4)  | (a3) |
| All regions | Roumania                     | 6000                                  | 150.000.000             | 6000                                   | 150.000.000                             | 1700   | 15   | 15   | 300  | 30  | 300  | 20000000   | 0    | -   | -  | -   | -    |
| Total       | Roumania                     | 6000                                  | 150.000.000             | 6000                                   | 150.000.000                             | 1700   | 15   | 15   | 300  | 30  | 300  | 20000000   | 0    | -   | -  | -   | -    |

(a) For zoonotic salmonella indicate the serotypes covered by the control programmes: (a1) for Salmonella Enteritidis, (a2) for Salmonella Typhimurium, (a3) for other serotypes- specify as appropriate, (a4) for Salmonella Enteritidis or Salmonella Typhimurium.

(b) For example, breeding flocks (rearing, adult flocks), production flocks, laying hen flocks, etc. Flocks equals herds or as appropriate.

(c) Total number of flocks existing in the region including eligible flocks and non-eligible flocks for the programme.

(d) Check means to perform a flock level test under the programme for the presence of Salmonella. In this column a flock should not be counted twice even if it has been checked more than once.

(e) If a flock has been checked, in accordance with footnote (d), more than once, a positive sample should be taken into account only once.

Revizie : 24.08.2012



## 7.2. Targets on vaccination

### 7.2.1. Targets on vaccination of broiler flocks- not applicable

Disease <sup>(a)</sup>: Zoonotic Salmonella

Animal species: gallus gallus

| Region       | Total number of herds in vaccination or programme | Total number of animals in vaccination programme | Targets on vaccination programme     |                                       |   | No of doses of vaccine expected to be administered |
|--------------|---|--|--------------------------------------|---------------------------------------|---|--|
|              |   |  | No of herds in vaccination programme | No of herds expected to be vaccinated | No of animals expected to be vaccinated |  |
| All regions  | -   | -  | -                                    | -                                     | -                                       | -  |
| <b>Total</b> | -   | -  | -                                    | -                                     | -                                       | -  |

**The vaccination is not mandatory and the costs regarding purchase of vaccine doses and the vaccination are incurred by the business operators.**

## 8. Detailed analysis of the cost of the programme<sup>1</sup>

Costs mentioned below are estimated for a one-year period (1 of January 2013-31 of December 2013)

| <b>Costs related to</b>          | <b>Specification</b>   | <b>Number of units</b> | <b>Unitary cost in €</b> | <b>Total amount in €</b> | <b>Community funding requested (yes/no)</b> |
|----------------------------------|--|------------------------|--------------------------|--------------------------|---|
| <b>1. Testing</b>                |  |                        |                          |                          |   |
| <b>1.1. Cost of the analysis</b> | Test: bacteriological exam for Salmonella spp.detection  | <b>2000</b>            | <b>16,70</b>             | <b>33400</b>             | <b>YES</b>                                  |
|                                  | Test : Salmonella spp. serotyping  | <b>500</b>             | <b>64.97</b>             | <b>32485</b>             | <b>YES</b>                                  |
|                                  | Test : the detection of antimicrobials or bacterial growth inhibitory effect in tissues from birds from flocks tested for Salmonella | <b>100</b>             | <b>47.44</b>             | <b>4744</b>              | <b>YES</b>                                  |
| <b>1.2. Cost of sampling</b>     | Official sample harvest  | <b>2000</b>            | <b>0,5</b>               | <b>1000</b>              | <b>YES</b>                                  |
|                                  | Disposable sterile containers for sampling of faeces   | <b>2000</b>            | <b>0,5</b>               | <b>1000</b>              | <b>NO</b>                                   |
|                                  | One use gloves-pairs   | <b>2000</b>            | <b>0,1</b>               | <b>200</b>               | <b>NO</b>                                   |
|                                  | Boot swabs   | <b>4000</b>            | <b>2</b>                 | <b>8000</b>              | <b>NO</b>                                   |
| <b>1.3. Other costs</b>          | Overcoats  | <b>2000</b>            | <b>1</b>                 | <b>2000</b>              | <b>NO</b>                                   |
| <b>2. Vaccination</b>            |  |                        |                          |                          |   |
| <b>2.1. Purchase of vaccine</b>  |  |                        |                          |                          |   |
| <b>2.2. Distribution costs</b>   |  |                        |                          |                          |   |
| <b>2.3. Administering costs</b>  |  |                        |                          |                          |   |

|  |   |            |           |                |            |
|--|---|------------|-----------|----------------|------------|
| <b>2.4. Control costs</b>  |   |            |           |                |            |
| <b>3. Slaughter and destruction</b>  |   |            |           |                |            |
| <b>3.1. Compensation of animals</b>  |   |            |           |                |            |
| <b>3.2. Transport costs</b>  |   |            |           |                |            |
| <b>3.3. Destruction costs</b>  |   |            |           |                |            |
| <b>3.4. Loss in case of slaughtering</b>   |   |            |           |                |            |
| <b>3.5 Costs from treatment of products (eggs, hatching eggs, heat treat of broilers, etc)</b> |   |            |           |                |            |
| <b>4. Cleaning and disinfection</b>  |   |            |           |                |            |
|  | Test: Bacteriological test to verify the efficiency of disinfection of poultry houses after depopulation of a Salmonella-positive flock | <b>150</b> | <b>10</b> | <b>1500</b>    | <b>YES</b> |
| <b>5. Salaries (staff contracted for the programme only)</b>                                   |   |            |           |                |            |
|  |   |            |           |                |            |
| <b>6. Consumables and specific equipment</b>   |   |            |           |                |            |
| <b>7. Other costs</b>  |   |            |           |                |            |
| <b>TOTAL</b>   |   |            |           | <b>73129 €</b> | <b>YES</b> |



# Romanian National Control Programme for *Salmonella* in Turkey 2013

## a. Aim of the programme

To reduce or maintain the low prevalence of *Salmonellas* of public health significance in fattening and adult breeding turkey flocks on holdings in Romania, at least to the target levels set out in Regulation (EC) No 584/2008 which is a maximum percentage of fattening and adult breeding turkey flocks remaining positive for *Salmonella* Enteritidis and *Salmonella* Typhimurium, *including also the monophasic strains of S. Typhimurium*, to 1% or less by 31 December 2012. Because Roumania has less than 100 flocks of adult breeding or fattening turkeys, our target shall be that no more than one flock of adult breeding or fattening turkeys may remain positive by 31 December 2012.

## b. Sampling programme

The National Control Programme for *Salmonella* encompasses the following serovars of zoonotic *Salmonella*: *Salmonella enteritidis* and *Salmonella typhimurium*.

The sampling programme will be in accordance to Regulation **2160/2003 EC** and **Regulation EC 584/2008**. We have also taken into account the Regulation **1177/2006 EC** and **Regulation 213/2009**.

The programme was elaborated in compliance with the requirements of Council Regulation **(EC) No 584/2008** implementing Regulation (EC) No 2160/2003 of the European Parliament as regards a Community target for the reduction of the prevalence of *Salmonella enteritidis* and *Salmonella typhimurium* in turkeys.

Operators will be required to implement the sampling programme in the Annex to Regulation (EC) 584/2008 and Regulation (EC) 200/2010. For convenience the 'Sampling protocol' is repeated below.

For **fattening turkeys** at least two pairs of boot sock/swabs, or one pair of boot sock/swabs and a dust sample will be taken by the operator within the period of three weeks before the birds are due for slaughter. Where possible, the samples will be taken in sufficient time for the laboratory results to be known before the birds are transported to the slaughterhouse.

For **breeding turkeys**, samples for the detection of *Salmonella* will be taken from rearing turkey breeding flocks at day-old, at four weeks of age and two weeks before moving to the laying phase or laying unit. In adult breeding flocks, samples shall be taken at least every third week during the laying period at the holding or at

the hatchery. The samples in adult breeding flocks shall be taken in accordance with the provisions laid down in Annex to Regulation (EC) No 200/2010.

Each year official samples will be taken by the local Competent Authority, or other authorized agent, acting on behalf of the Competent Authority. When an official sample is taken it may replace the sample required to be taken by the operator. Sampling to verify the achievement of the target will be as detailed in "Sampling protocol" in the Annex to Commission Regulation (EC) No. 584/2008.

**c. Demonstrate the evidence that it complies with the specific requirements laid down in Parts C and E of Annex II to Regulation (EC) no 2160/2003**

### ***Specific requirement concerning fresh meat***

When a turkey flock is confirmed for the presence of *Salmonella enteritidis* or *Salmonella typhimurium*:

a. Fresh meat from turkey may be placed on the market on the condition that it meets the requirement of absence of *Salmonella* in 25 grams from the meat.

b. The requirement laid down in point (a) does not apply to fresh turkey meat destined for heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene.

c. The criteria laid down in point (a) does not apply to fresh turkey meat destined for industrial heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene.

### ***Specific requirements concerning breeding flocks of turkey***

1. Starting with 1 January 2010 the measures laid down below in points 3 to 5 must be taken whenever the analysis of samples carried out in accordance with point 2.2.2.1 and 2.2.2.2 of Annex of the Regulation 200/2010 indicates the presence of *Salmonella enteritidis* or *Salmonella typhimurium* in a breeding flock of turkey in the circumstances set out in point 2.

2. (a) If the competent authority has approved the method of analysis used for samples taken in accordance with part B of the Regulation 2160/2003/EC , it may require that the measures laid down in points 3 to 5 be taken when such analysis detects the presence of *Salmonella enteritidis* or *Salmonella typhimurium*.

(b) Otherwise, the measures laid down in points 3 to 5 must be taken whenever the competent authority confirms a suspicion of the presence of *Salmonella enteritidis* or *Salmonella typhimurium* arising from the analysis of samples carried out in accordance with part B of the Regulation 2160/2003/EC.

**3. Non-incubated eggs from the flock must be destroyed.**

However, such eggs may be used for human consumption if they are treated in a manner that guarantees the elimination of *Salmonella enteritidis* and *Salmonella typhimurium* in accordance with Community legislation on food hygiene.

4. **All birds**, including day-old chicks, **in the flock must be slaughtered or destroyed so as to reduce as much as possible the risk of spreading Salmonella**. Slaughtering must be carried out in accordance with Community legislation on food hygiene. Products derived from such birds may be placed on the market for human consumption in accordance with Community legislation on food hygiene and, once applicable, part E of the Regulation 2160/2003. If not destined for human consumption, such products must be used or disposed of in accordance with Regulation (EC) No 1774/2002 of the European Parliament and of the Council, laying down health rules concerning animal by-products not intended for human consumption.

5. Where **eggs** for hatching from flocks in which *Salmonella enteritidis* or *Salmonella typhimurium* is present are still **present in a hatchery, they must be destroyed or treated** in accordance with Regulation (EC) No 1774/2002.

## 1. General

### 1.1 Summary referring to the occurrence of the salmonellosis in Romania

The programme for the control of *Salmonella Enteritidis* and *Salmonella Typhimurium* in turkey fattening flocks has been in operation in Romania from 2010. As a result, the number of *Salmonella Enteritidis* and *Salmonella Typhimurium* infected turkey flocks is currently very low. During 2011, a totally of 99 flocks were tested for *Salmonella* infection from which a no of 40 were tested in official control and there were none positive flocks for *Salmonella Typhimurium* and *Salmonella Enteritidis*. The prevalence for the target serotypes in broiler flock in 2011 was 0%, which is low and below the Community target.

The incidence of various serotypes of *Salmonella* in turkey farms, in 2011 in Roumania

| NO. | SEROTYPE             | NO. OF STRAINS in turkeys |
|-----|----------------------|---------------------------|
| 1.  | <b>S. Agona</b>      | 1                         |
| 2.  | <b>S. Newport</b>    | 5                         |
| 3.  | <b>S. Saintpaul</b>  | 8                         |
| 4.  | <b>S.Senftenberg</b> | 1                         |
| 5.  | <b>TOTAL</b>         | 15                        |

### 1.2 The structure and organization of the relevant Competent Authorities

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in breeding turkey flocks is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**

---

The departments responsible for implementing the National Control Programme of Salmonella in turkey are:

- for implementing at national level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary General Directorate** ;
- in respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the verification of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**
- Concerning the slaughter of flocks, the **Hygiene and Veterinary Epidemiology Directorate** is responsible for public health protection, through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.
- In respect of EC Regulation No 183/2005 on feed hygiene at the central level is **The Directorate for Technical Coordination of Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**
- at county level for implementing the Regulation (EC) No 2160/2003,882/2004 and 183/2005 there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the national reference laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

**NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

**Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România

Tel: 0374.322.013 / 0374.322.000

Fax: 0214.113.394

E-mail: [office@idah.ro](mailto:office@idah.ro)



- **H.I.P.V.H.** (Hygiene Institute and Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

## NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED

### Hygiene Institute and Veterinary Public Health

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201

Tel: 021.252.46.51

Fax: 021.252.00.61

E-mail: [iispv @ iispv.ro](mailto:iispv@iispv.ro)

### 1.3 Approved laboratories where samples collected within the programme are analyzed

**Diagnostics:** The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL at January 2012 is show below:

| Nr. Crt. | NAME                      | ADRESS   | TELEPHON/ E-MAIL   |
|----------|---------------------------|--|--|
| 1        | I.D.S.A                   | București, Str. Dr.Staicovici, nr.63, sector 5, Cod 050557     | 0374/322.013<br><a href="mailto:office@idah.ro">office@idah.ro</a>   |
| 2        | L.S.V.S.A. Alba           | Alba Iulia, Str. Lalelelor nr.7A, Cod 510217, Jud. Alba        | 0258/835.915<br><a href="mailto:office-alba@ansvsa.ro">office-alba@ansvsa.ro</a>   |
| 3        | L.S.V.S.A. Bacău          | Bacau, Str Bucovinei, Nr 21, Jud. Bacău                        | 0234/586.233, int.111<br><a href="mailto:office-bacau@ansvsa.ro">office-bacau@ansvsa.ro</a>  |
| 4        | L.S.V.S.A.Bistrița-Năsăud | Bistrița, Str. Tarpiului 26, Cod 420062, Jud. Bistrița Năsăud  | 0263/206.027; 0263/224.974<br><a href="mailto:office-bistrita-nasaud@ansvsa.ro">office-bistrita-nasaud@ansvsa.ro</a>                           |
| 5        | L.S.V.S.A. Botoșani       | Botoșani , Str Tudor Vladimirescu Nr.17, Jud. Botoșani         | 0231/512.766<br><a href="mailto:office-botosani@ansvsa.ro">office-botosani@ansvsa.ro</a>   |
| 6        | L.S.V.S.A. Brașov         | Brasov, Str. Calea Feldioarei Nr. 20 A, Cod 500450, Jud.Brașov | 0268/440.257<br><a href="mailto:office-brasov@ansvsa.ro">office-brasov@ansvsa.ro</a><br><a href="mailto:dsvbv@rdslink.ro">dsvbv@rdslink.ro</a> |

|    |                             |  |   |
|----|-----------------------------|--|---|
| 7  | <b>L.S.V.S.A. Brăila</b>    | Brăila, str. Calea Galați, nr.344, Cod 810385, Jud. Brăila             | 0239/610.689<br>dsv@braila.rdsnet.ro<br><a href="mailto:office-braila@ansvsa.ro">office-braila@ansvsa.ro</a>  |
| 8  | <b>L.S.V.S.A. Buzău</b>     | Buzau, Str. Horticolei, nr. 58 bis, Jud. Buzau                         | 0238-725001<br>0238-725002<br><a href="mailto:office-buzau@ansv.ro">office-buzau@ansv.ro</a>  |
| 9  | <b>L.S.V.S.A. Călărași</b>  | Călărași, Str. Prelungirea Dobrogei , nr. 4, Jud.Călărași              | 0242/313.676,<br><a href="mailto:office-calarasi@ansvsa.ro">office-calarasi@ansvsa.ro</a> ;<br><a href="mailto:office_dsv@satline.ro">office_dsv@satline.ro</a> ; |
| 10 | <b>L.S.V.S.A Constanța</b>  | Constanta, Sos. Mangaliei Nr. 78, Jud. Constanța                       | 0241-682-417,<br>email: <a href="mailto:dsvct@rdset.ro">dsvct@rdset.ro</a>  |
| 11 | <b>L.S.V.S.A. Cluj</b>      | Cluj-Napoca, Piata Marasti Nr. 1, Jud. Cluj                            | 0264-445729, 0264-448177<br><a href="mailto:office-cluj@ansvsa.ro">office-cluj@ansvsa.ro</a><br><a href="mailto:dsvcj@rdslink.ro">dsvcj@rdslink.ro</a>            |
| 11 | <b>L.S.V.S.A. Dâmbovița</b> | Târgoviște, Str. I.C. Brătianu, nr. 35, Cod 130055, Jud. Dâmbovița     | 0372/737818<br><a href="mailto:office-dambovita@ansvsa.ro">office-dambovita@ansvsa.ro</a>   |
| 13 | <b>L.S.V.S.A. Giurgiu</b>   | Giurgiu, Str. Bucuresti nr.72, Jud. Giurgiu                            | 0246/ 230.491<br><a href="mailto:office-giurgiu@ansvsa.ro">office-giurgiu@ansvsa.ro</a>   |
| 14 | <b>L.S.V.S.A. Gorj</b>      | Târgul-Jiu, Str.Ecaterina Teodoroiu Nr.523, Jud. Gorj                  | 0253/226.033<br><a href="mailto:dsv@intergorj.ro">dsv@intergorj.ro</a><br><a href="mailto:office-gorj@ansvsa.ro">office-gorj@ansvsa.ro</a>                        |
| 15 | <b>L.S.V.S.A Hunedoara</b>  | Deva, Str. 22 Decembrie Nr. 226, Jud. Hunedoara                        | 0254-221145, 0254-230527<br><a href="mailto:office-hunedoara@ansvsa.ro">office-hunedoara@ansvsa.ro</a>  |
| 16 | <b>L.S.V.S.A. Ialomița</b>  | Slobozia, str. Lacului, nr. 12 , Jud. Ialomița                         | 0243/ 232.069<br><a href="mailto:dsvsa-ialomita@ansvsa.ro">dsvsa-ialomita@ansvsa.ro</a>   |
| 17 | <b>L.S.V.S.A. Iași</b>      | Iași, Aleea Mihail Sadoveanu 10, Jud. Iași                             | 0232-267501<br><a href="mailto:office-iasi@ansvsa.ro">office-iasi@ansvsa.ro</a>   |
| 18 | <b>L.S.V.S.A. Maramureș</b> | Baia Mare, Str. Vasile Alecsandri, Nr. 66, Jud. Maramureș              | 0262/ 224.031<br><a href="mailto:office-maramures@ansvsa.ro">office-maramures@ansvsa.ro</a>   |
| 19 | <b>L.S.V.S.A. Mureș</b>     | Târgu Mures, str. Podeni nr.10 cod 540253, Jud. Mureș                  | 0265/ 314.975<br><a href="mailto:office-mures@ansvsa.ro">office-mures@ansvsa.ro</a>   |
| 20 | <b>L.S.V.S.A. Neamț</b>     | Piatra Neamț, Aleea Tiparului, nr. 12, cod 610263, Jud. Neamț          | 0233/ 223.259<br><a href="mailto:office-neamt@ansvsa.ro">office-neamt@ansvsa.ro</a>   |
| 21 | <b>L.S.V.S.A. Prahova</b>   | Ploiesti, Str. Corlatesti, nr. 11, Jud. Prahova,                       | 0244 / 57.17.51<br><a href="mailto:office-prahova@ansvsa.ro">office-prahova@ansvsa.ro</a>   |
| 22 | <b>L.S.V.S.A. Satu Mare</b> | Satu Mare, Str. Lăcrimioarei, Nr. 37, cod poștal 440067, Jud.Satu Mare | 0261/ 715.956<br><a href="mailto:office-satu-mare@ansvsa.ro">office-satu-mare@ansvsa.ro</a>   |
| 23 | <b>L.S.V.S.A. Sibiu</b>     | Sibiu, str. Calea Șurii Mari, nr. 21, cod 550089, Jud. Sibiu           | 0269/223.069<br><a href="mailto:office-sibiu@ansvsa.ro">office-sibiu@ansvsa.ro</a>  |
| 24 | <b>L.S.V.S.A. Suceava</b>   | Suceava, Str. Scurta Nr. 2, Jud Suceava                                | Tel: 0230-522848,<br><a href="mailto:dsvsv@suceava.rdsnet.ro">dsvsv@suceava.rdsnet.ro</a>   |
| 25 | <b>L.S.V.S.A. Timiș</b>     | Timisoara, Str. Surorile Martir Caceu Nr.4, Jud. Timiș                 | 0256-204911<br><a href="mailto:office-timis@ansv.ro">office-timis@ansv.ro</a>   |
| 26 | <b>L.S.V.S.A. Vaslui</b>    | Bârlad, str. Trestiana, nr. 2, cod 731030, Jud.Vaslui                  | 0235-421121,<br>0235-421413<br><a href="mailto:office-vaslui@ansv.ro">office-vaslui@ansv.ro</a>   |
| 27 | <b>L.S.V.S.A. Vrancea</b>   | Focsani,B-dul Brailei, nr. 121 bis, cod 620122, Jud.Vrancea            | 0237-215561,<br>0237-232727<br><a href="mailto:office-vrancea@ansv.ro">office-vrancea@ansv.ro</a>   |

All Laboratories have to use the diagnostic methods presented at the point 1.4.

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the serotype

according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

#### **1.4 Methods used in the examination of the samples in the framework of the programme**

Samples harvest by operators and samples harvest as official controls are prepared and tested in accordance with the requirements of the Commission Regulation (EC) No 200/2010, using the method recommended by the Community Reference Laboratory for *Salmonella* in Bilthoven, Netherlands. The method is described in the current version of ISO 6579:/A1:2007): 'Detection of *Salmonella* spp. in animal faeces and in samples of the primary production stage'. A semi-solid medium (modified semi-solid Rappaport-Vassiladis medium, MSR/V) is used as the single selective enrichment medium. At least one isolate from each positive sample shall be sero-typed according to the Kaufmann-White scheme. In general the sensitivity of a panel of 10 antimicrobials will be determined.

#### **1.5 Official controls**

In accordance with Regulation (EC) No 584/2008 the Competent Authority will select each year at random:

- All flocks on at least 10% of holdings with more than 500 **fattening turkeys**.
- All flocks on least 10% of holdings with at least 250 adult **breeding turkeys** between 30 and 45 weeks of age, but including all holdings where *S. Enteritidis* or *S. Typhimurium* has been detected during the previous 12 months and all holdings with elite, great grandparents and grandparent breeding turkeys. This sampling may also take place at the hatchery
- Each time the Competent Authority considers it necessary

In the case of fattening turkeys the flocks should be tested within 3 weeks of going for slaughter. The flocks will be sampled in accordance with the Annex in Regulation (EC) No 584/2008..

The use of antimicrobials (as defined in Regulation (EC) No 1177/2006) will be checked when the official sample is taken. If the flock is under antimicrobial medication for animal health or animal welfare reasons the flock will be sampled again after the period of withdrawal considered suitable by the Competent Authority and taking into account the product Marketing Authorisation. Flock owners are required to keep records of antimicrobial use and to make these records available .

The records of samples taken by the operator will be made available for inspection to the Competent Authority or its agent and will provide details of the identity of the flock sampled, date of sample, slaughter date, type of sample,

laboratory carrying out the examination, and the result. The number of flocks on the holding and the number of birds.

Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs requires poultry abattoirs and establishments handling processed meat to demonstrate compliance with microbiological criteria for *Salmonella* that includes undertaking microbiological testing for *Salmonella* and corrective action when the criteria are not met. The sampling frequencies are prescribed in the Regulation but do not apply to establishments producing small quantities.

### **Official controls at other stages of the food chain.**

Under the terms of the EC Feed Hygiene **Regulation 183/2005** feed businesses operators must be approved or registered by the Local Authority. Approvals/registrations are issued for the producers of compound feeds, feed materials, feed additives and premixtures. Approval requires a prior-inspection visit by the Local Authority to ensure that the establishments are in conformity with the required standards. The competent authority performs checks according to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and the program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order.

### **1.6 Measures taken by the Competent Authorities with regard to animals or products in which the presence of *Salmonella* spp. have been detected**

In case of suspicion or confirmation of *Salmonella enteritidis* or *Salmonella typhimurium* the NRL/C.S.V.F.S.L. shall notify immediately the N.S.V.F.S.A and the local C.S.V.F.S.D.

When a turkey flock is suspected of being infected with *Salmonella* Enteritidis or *Salmonella* Typhimurium the flock will be investigated. The flock is suspected of being infected when *S. Enteritidis* or *S. Typhimurium* is isolated from samples, carried out privately or as required by either the operator or the Competent Authority as detailed in the Annex to Regulation (EC) No 584/2008. Tissue/organs may be taken from birds as part of the investigation of clinical disease; these cases will be discussed and additional follow up investigation carried out as appropriate, along with advice on *Salmonella* control. Neck skins are sampled at abattoirs under Regulation (EC) No 1441/2007. The potential for cross contamination make the results from abattoir sampling difficult to interpret in relation to a particular flock.

Isolates of *S. Enteritidis* and *S. Typhimurium* will be examined to determine if they are vaccine strains according to the manufacturer's specification. If vaccine

strains are confirmed in samples the flock will not be classed as positive for the purposes of establishing the progress towards the target. Although vaccines against *Salmonella* are not currently used in meat turkeys in Romania.

### **Fattening turkeys**

In most cases it will not be possible to carry out an investigation of the birds in the flock from which the *Salmonella* was isolated as the birds will have been slaughtered shortly after the results become available. A notice may be served by the Competent Authority as necessary requiring the operator to clean and disinfect the building from which the infected flock originated. After cleaning and disinfecting the building the operator may be required to take swabs from a number of sites in the building and submit them to an approved laboratory to be tested for *Salmonella* in order to check on the efficiency of the hygiene measures taken. In cases where *S. Enteritidis* or *S. Typhimurium* was isolated, the cleaning and disinfection should be checked by the Competent Authority or its agent. When *S. Enteritidis* or *S. Typhimurium* is detected in a flock of fattening turkeys, sampling by the Competent Authority will take place in all flocks on the holding in circumstances as laid out in Regulation (EC) No 584/2008.

When a fattening flock is positive for *Salmonella* in the samples taken during the period 3 weeks before slaughter the food business operator for the abattoir where the birds are to be slaughtered should be informed so that arrangements can be made to take measures to reduce the possibility of cross-contamination of other batches, for example arranging slaughter to take place at the end of the day, or before a break, etc.

### **Breeding Turkeys**

When a breeding turkey flock is suspected of being infected with *Salmonella* Enteritidis or *Salmonella* Typhimurium, they will be subject to controls as laid out in Regulation (EC) No 2160/2003 and Community legislation on food hygiene. This applies to breeding flocks from day old through to end of production.

The owner or person responsible for the flock is required to clean and disinfect the building where the infected birds were kept, and provide evidence to the Competent Authority that the cleaning and disinfection has been satisfactory.

Sampling by the Competent Authority will be carried out on all flocks on holdings in case of detection of *Salmonella* Enteritidis or *Salmonella* Typhimurium from samples taken at the hatchery by food business operators or within the frame of official controls, to investigate the origin of infection. Official sampling will be carried out on all holdings/hatcheries where *S. Enteritidis* or *S. Typhimurium* was detected during the previous 12 months as part of the framework of annual official sampling to be carried out under the control of the Competent Authority. This sampling will take place according to the procedure in Annex to Regulation (EC) No 200/2010.

## General

Operators with a flock which is positive for *S. Enteritidis* or *S. Typhimurium* will be contacted by the Competent Authority and invited to contact their veterinary adviser for advice on how to reduce or eliminate the *Salmonella*. Advice may include recommendations on management, cleaning and disinfection, pest control, biosecurity, monitoring, and the potential use of other aids in the control of *Salmonella*.

The operator/owner in consultation with his/her veterinarian may consider vaccination of the flock against *Salmonella* with a product which has a marketing authorisation in line with the requirements of Commission Regulation (EC) No. 1177/2006 as regards requirements for the use of specific control methods in the framework of the national programmes for the control of *Salmonella*. Vaccination may only be used as a preventative measure; it is not an alternative to the requirements in Annex II C of Commission Regulation (EC) No 2160/2003 for the use of specific control methods in the framework of the national programmes for the control of *Salmonella*. Antimicrobial treatment may not be used for the control of *Salmonella* in the national control programme except within the limits set by Commission Regulation (EC) No.1177/2006. For the purposes of establishing the progress towards the target if *S. Enteritidis* or *S. Typhimurium* is isolated from either an operator sample or an official sample the flock is classed as positive. A flock positive for a specific serotype will be recorded only once for that serotype.

### 1.7 Relevant national legislation.

- N.S.V.F.S.A President **Order no. 34** to approve the sanitary veterinary norm regarding to monitoring zoonoses and zoonotic agents

- Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order

- N.S.V.F.S.A President **Order no. 205** for the approval of national reference laboratories and duties

- N.S.V.F.S.A President **Order no. 160** for the approval of the sanitary veterinary norm regarding the community reference laboratory for the zoonoses epidemiology and Salmonella and the national reference laboratory for Salmonella.

### 1.8 Financial assistance provided to food and feed businesses

In turkey flocks no financial assistance is provided in the context of the control programme. Romania applied to the Commission for co-financing for certain aspects of the control programme within the terms of Council Decision **2009/470/EC** on expenditure in the veterinary field.

## 2. Food and Feed businesses covered by the programme.

The programme for the control of *Salmonella* in turkeys is part of the controls along the whole food chain.

### 2.1 The structure of the production of the given species and products thereof.

The structure of the turkey fattening flocks and holdings in Romania at the beginning of 2011 is presented below:

| Region<br>(County+NUTS CODE) | Type of<br>flocks   | Number of flocks |
|------------------------------|---------------------|------------------|
| RO122-BV                     | Fattening<br>turkey | 96               |
| RO125-MS                     | Fattening<br>turkey | 3                |
| RO411-DJ                     | Fattening<br>turkey | 12               |
| <b>Total</b>                 |                     | <b>111</b>       |

### 2.2 The structure of the production of food

At the end of production period the birds are slaughtered and may go for human consumption. In accordance with Regulation 853/2004 provisions, poultry meat intended for human consumption must be obtained in approved slaughterhouses. There are **40** approved slaughterhouses in Romania.

Poultry feed is supplied to farms by a small number of manufacturers. The major manufacturers of poultry feed operate to assurance schemes, apply HACCP principles and monitor for *Salmonella*.

### 2.3 Relevant guides for good animal husbandry practices or other guidelines

Some of the relevant guides for good animal husbandry practices are presented at the below link:

[http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella\\_12404ro.pdf](http://www.ansvsa.ro/documente/admin/ghid%20rozatoare%20-%20salmonella_12404ro.pdf)

[http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010\\_13247ro.pdf](http://www.ansvsa.ro/documente/admin/GHID%20Sall%20ferma%20broiler%20-%202002.10.2010_13247ro.pdf)

[http://www.ansvsa.ro/documente/admin/community\\_guide\\_layers\\_hygiene\\_practice\\_pullet\\_egg\\_ro\\_17813ro.pdf](http://www.ansvsa.ro/documente/admin/community_guide_layers_hygiene_practice_pullet_egg_ro_17813ro.pdf)

Other relevant guides for good animal husbandry practices and other guidelines are under constructions.

## **2.4 Routine veterinary supervision of farms**

The owner is responsible for the health and welfare of the poultry on the holding, and for ensuring that a veterinarian is consulted on disease and welfare issues as appropriate. It is mandatory for each holding to have a contract with a private veterinarian who is responsible for veterinary care. A veterinarian on behalf of the Competent Authority carries out inspections on farms for animal welfare reasons, to take samples for residues to administer and enforce the legislation on marketing of eggs and to check medicine records.

Also a veterinarian on behalf of the Competent Authority visit the farms and take official samples in the framework of Salmonella NCP according with the legislation in force.

It is mandatory for each county sanitary veterinary and food safety directorate (local CA) to report to the NSVFSA every month the number of samples and results of these tests for each flock. Also the Salmonella NRL has the obligation to notify immediately NSVFSA and CSVFSD each positive sample for the relevant Salmonella.

## **2.5 Registration of farms**

All commercial poultry holdings in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centers / holdings of origin and means of transport for health and animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products. All commercial holdings are entered into our national data base.

All broiler turkey flocks of more than 500 birds and all breeding turkey flock with more than 250 birds are included in the Salmonella National Control Programme.

## **2.6 Record-keeping at farms**

- a. All turkey flock operators are required to keep records of veterinary medicines use, including vaccines, which must be available for inspection.
- b. Records relating to movement of flocks on to/off the holding must be kept.
- c. Records giving details of sampling for *Salmonella* and results will be kept either at the holding or be readily available.



## **2.7 Documents to accompany animals when dispatched**

Operators wishing to export birds or hatching eggs to another EU Member State (or certain third countries) must comply with **EU Directive 2009/158/EC** and ensure that the consignment is accompanied by a completed and signed Intra-trade Animal Health Certificate (ITAHC) for poultry breeding and production. This must be completed and signed by the Official Veterinarian as well as the operator to confirm compliance with the relevant articles of Directive.

The ITAHC will also require the reference number of the operator's poultry health certificate. The ITAHC is amended to include the results of the last test for Salmonella as is required in the Commission Regulation (EC) 2160/2003 Art 9.1 prior to any dispatching of the live animals, or hatching eggs, from the food business operators place of origin. The date and the result of testing are included in the relevant health certificate provided in the Community legislation.

## **2.8 Other relevant measures to ensure the traceability of animals.**

It is require to the operators of poultry flocks to keep records of poultry or hatching eggs entering or leaving the premises. The records must contain information on the number, date, and origin or destination. These records must be available to the Competent Authority for inspection.

All official veterinary health certificates issued for the intra community trade of poultry and hatching eggs are recorded on the Trade Control and Expert System (TRACES). This system allows tracking of exports of live animals and hatching eggs accompanied by veterinary health certification. TRACES generate ITAHCs issued for intra-Community movements. TRACES are an internet-based service which is owned and maintained by the Commission.

## **PART B**

### **1. Identification of the programme**

Member State: **Romania**  
Disease: **Salmonella enteritidis and Salmonella typhimurium in turkey**

Year of implementation: **2013**  
Reference of this document: **National Sanitary Veterinary and Food Safety Authority**  
**No. /**

Contact : **Dr. Nicolae LAZĂR,**  
**Director,**  
**phone: +4 021 315 78 75;**  
**fax: +4021 31249 67,**  
**e-mail: [lazar.niculae@ansvsa.ro](mailto:lazar.niculae@ansvsa.ro)**

**Date sent to the Commision : 29.04.2012**

*Salmonella* has been recognized as an important zoonotic pathogen for many years. *Salmonella Enteritidis* and *Salmonella Typhimurium* have accounted for the majority of cases of human salmonellosis and have consistently been the most commonly implicated pathogens in general outbreaks of food-borne disease.

A National Control Programme of *Salmonella* in turkey flocks was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 584/2006. The national control programme for *Salmonella* in turkey flocks was coming into effect in January 2010.

The programme for the control of *Salmonella Enteritidis* and *Salmonella Typhimurium* in turkey fattening flocks has been in operation in Romania from 2010. As a result, the number of *Salmonella Enteritidis* and *Salmonella Typhimurium* infected turkey flocks is currently very low. During 2011, a totally of 99 flocks were tested for *Salmonella* infection from which a no of 40 were tested in official control and there were none positive flocks for *Salmonella Typhimurium* and *Salmonella Enteritidis*. The prevalence for the target serotypes in broiler flock in 2011 was 0%, which is low and below the Community target.

#### **The incidence of various serotypes of *Salmonella* in turkeys flocks, in 2011**

| <b>NO.</b> | <b>SEROTYPE</b>   | <b>NO. OF STRAINS in turkeys</b> |
|------------|-------------------|----------------------------------|
| 6.         | <b>S. Agona</b>   | 1                                |
| 7.         | <b>S. Newport</b> | 5                                |

|     |                      |           |
|-----|----------------------|-----------|
| 8.  | <b>S. Saintpaul</b>  | <b>8</b>  |
| 9.  | <b>S.Senftenberg</b> | <b>1</b>  |
| 10. | <b>TOTAL</b>         | <b>15</b> |

It is a statutory requirement for all laboratories which isolate *Salmonella* from a flock or its environment to report the finding and supply the isolate to the National Reference Laboratory (NRL) for *Salmonella*. The isolates are sero-typed, phage-typed (Cantacuzino Institute), where appropriate, and tested for antimicrobial sensitivity by the NRL. This information is recorded and analyzed. The number of reports received depends on the level and sensitivity of monitoring which is undertaken by the producers. The reports provide useful information on the serovars which are most common in poultry, and indicate trends.

### **3. Description of the submitted programme**

#### **Objectives**

The main objective of our programme is to reduce or maintain the low prevalence of *Salmonellas* of public health significance in fattening and adult breeding turkey flocks on holdings in Romania, at least to the target levels set out in Regulation (EC) No 584/2008 which is a maximum percentage of fattening and adult breeding turkey flocks remaining positive for *Salmonella* Enteritidis and *Salmonella* Typhimurium to 1% or less by 31 December 2012. Because Roumania has less than 100 flocks fattening turkeys, our target shall be that no more than one flock of fattening turkeys may remain positive by 31 December 2012.

The subsidiary objectives of the programme include further reduction of the incidence of infected table eggs, the reduction of incidence of meat and eventually the reduction of the occurrence of human salmonellosis and a consequent reduction in suffering, mortality and health service costs. This programme is in accordance to Council Directive 1999/74/EC, Commission Decision 2008/425/EC, Council Decision 92/65/ EEC, Council Decision 2009/470/ EEC, Regulation 2160/2003/EC, Regulation 584/2008/EC and Regulation 1177/2006/ EC.

A national *Salmonella* control programme was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 584/2008. The National Control Programme for *Salmonella* in fattening and breeding turkeys come into effect in January 2010.

#### **Target animal population**

All flocks of 250 or more breeding turkeys and all fattening turkeys flocks will be included in the National Control Programme unless exempted in Regulation (EC) No. 2160/2003 under Article 1.3 (birds produced for private domestic consumption, or where there is direct supply of small quantities of products to the final consumer or to local retail establishments directly supplying the primary products to the final consumer).

The number of holdings and flocks in turkeys flock rearing for meat sector at the beginning of 2011 are shown **in the table below**.

**Table . Number of holdings and flocks in turkey sector at the beginning of 2011**

| Region<br>(County+NUTS CODE) | Type of<br>flocks   | Number of flocks |
|------------------------------|---------------------|------------------|
| RO122-BV                     | Fattening<br>turkey | 96               |
| RO125-MS                     | Fattening<br>turkey | 3                |
| RO411-DJ                     | Fattening<br>turkey | 12               |
| <b>Total</b>                 |                     | <b>111</b>       |

### **Sampling programmes**

The National Salmonella Control Programme encompasses the following serovars of zoonotic Salmonella: *Salmonella enteritidis* and *Salmonella typhimurium* .

The sampling programme will be in accordance to Regulation 2160/2003 EC and Regulation 584/2008 EC. We have also taken into account Regulation 1177/2006 EC.

#### **4.1. Summary of measures under the programme**

##### **First year: 2013**

- Control
- Testing
- Slaughter of positive animals
- Killing of positive animals
- Vaccination
- Treatment of animal products
- Disposal of products
- Monitoring or surveillance
- Other measures

**4.2. Designation of the central authority charged with supervising and coordinating the departments responsible for implementing the programme:**

The Central Competent Authority for the National Control Programme in respect of EC Regulation 2160/2003 for the control of *Salmonella* in breeding hens flocks of *Gallus gallus* is:

**AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ  
ȘI PENTRU SIGURANȚA ALIMENTELOR**

---

The departments responsible for implementing the National Control Programme of Salmonella in breeding hens are:

- for implementing at central level of Regulation (EC) No 2160/2003 provisions regarding animal health status is **Sanitary Veterinary General Directorate** ;
- In respect of Regulation (EC) No 882/2004 provisions concerning official controls performed in view to ensure the checking of compliance with the feed and food law the main body at central level is **Inspection and Control Directorate**.
- Concerning the slaughter of flocks, the **Hygiene and Veterinary Epidemiology Directorate** is responsible for public health protection, through proportionate enforcement of legislation in meat processing plants and sanitary veterinary authorization.
- In respect of EC Regulation No **183/2005** on feed hygiene at the central level is **The Directorate for Technical Coordination of Reference Institutes, Sanitary Veterinary and Food Safety Laboratories, Pharmacovigilance and Animal Nutrition**
- At county level for implementing the Regulation (EC) No **2160/2003, 882/2004 and 183/2005** there are 42 county Sanitary Veterinary and Food Safety Directorates (**S.V.F.S.D**) and 41 county Sanitary Veterinary and Food Safety Laboratories
- **I.D.A.H.** (Institute for Diagnosis and Animal Health) is the National Reference Laboratory concerning animal health and in its structure is the NRL for Salmonella in live animals.

**NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN ANIMALS**

**Institute for Diagnosis and Animal Health**

Dr. Staicovici street, no. 63, district 5, cod 050557, Bucharest, România

Tel: 0374.322.013 / 0374.322.000

Fax: 0214.113.394

E-mail: [office@idah.ro](mailto:office@idah.ro)

- **H.I.P.V.H.** (Hygiene Institute for Veterinary Public Health) is the national reference laboratory concerning the expertise of food and feed and in its structure is the NRL for Salmonella in food and feed

## **NATIONAL REFERENCE LABORATORY FOR SALMONELLOSIS IN FOOD AND FEED**

### **Hygiene Institute for Sanitary Veterinary Public Health**

Campul Mosilor street, no. 5, district 2, Bucharest, Romania, cod 021201

Tel: 021.252.46.51

Fax: 021.252.00.61

E-mail: [iispv @ iispv.ro](mailto:iispv@iispv.ro)

The laboratory examinations for *Salmonellosis* in animals are performed by the National Reference Laboratory for *Salmonellosis* in animals at the Institute for Diagnosis and Animals Health– Bucharest at national level and at the county level at the County Sanitary Veterinary and Food Safety Laboratory (CSVFSL). A list of the CSVFSL at January 2012 is show in the link below:

[http://www.ansvsa.ro/documente/admin/lista%20laboratoarelor%202011%20-%2028.01.2011\\_11043ro.pdf](http://www.ansvsa.ro/documente/admin/lista%20laboratoarelor%202011%20-%2028.01.2011_11043ro.pdf)

At the National Reference Laboratory for Salmonella in animals from the Institute for Diagnosis and Animals Health– Bucharest is performing the sero-type according to the Kaufmann-White scheme from each positive sample found in Romania. Also here, is test the sensitivity of antimicrobials from each isolate.

### **1. N.S.V.F.S.A.**

- proposal of the plan of disease control.
- elaborate and submit to the EC the National Control Programme
- evaluate, coordinate the implementation of the National Control Programme in Romania and propose the update of this programme if is necessary (depending of the national situation)
- adoption of measures based on the disease situation in Romania
- submission of reports to the E.C.
- training the specialists from C.S.V.F.S.D

### **2. S.V.F.S.D.(Sanitary Veterinary and Food Safety Direction).**

- Coordination of the programme at the county level
- Official sampling is performed by the official veterinarian

- Propose the compensation costs to the Commission which will evaluate compensation documents.

**3. I.D.A.H.** Institute of Diagnosis and Animal Health is the National Reference Laboratory concerning animal health and was also designated as NRL for Salmonella in animals.

Responsibilities and tasks of the National Reference Laboratory for Salmonella (I.D.A.H.), pursuant to Directive 2003/99/EC and Regulation (EC) No 2160/2003, according to provision of N.S.V.F.S.A.. President Order no.160/2006 and 205/2007 are:

### **1. General duties**

(a) To collaborate with the Community reference laboratory in their area of competence.

(b) To coordinate, as appropriate, the activities of laboratories responsible for the analysis of samples in accordance with, in particular, Articles 4, 5 and 7 of Directive 2003/99/EC.

(c) To coordinate the activities of laboratories responsible for the analysis of samples in accordance with Article 12(1) of Regulation (EC) No 2160/2003/EC.

(d) Where appropriate, to organise comparative tests between the laboratories referred to under (b) and (c) and to assure an appropriate follow-up of such comparative testing.

(e) To ensure the dissemination to the competent authority and to the laboratories referred to under (b) and (c), of the information that the Community reference laboratory supplies.

(f) To provide scientific and technical assistance to the National Sanitary Veterinary and Food Safety Authority in their area of competence.

(g) Characterize the pathogen isolates, genetic typing of this agents

(h) Keep in maxim security conditions, the isolates

(i) Give to the C.E., CRL, OMS, and national reference Laboratory of other member states, with the accord of N.S.V.F.S.A. all the information required.

(j) train of the specialists from the government and private laboratory.

### **2. Specific functions and duties**



(a) To participate, as appropriate in the monitoring and control programme for Salmonella and related anti-microbial resistance pursuant to Directive 2003/99/EC and in the analysis and testing of *Salmonella* pursuant to Regulation (EC) No 2160/2003.

(b) To inform, as appropriate, the Community reference laboratory on aspects related to Salmonella vaccine strains and other specific control methods.

(d) To gather data and information on the activities developed and methods used in relevant laboratories and to inform the Community reference laboratory thereof.

(e) To monitor the epidemiological evolution of salmonella in Roumania.

#### **5. S.V.F.S.L.**

There are 41 county Sanitary Veterinary and Food Safety Laboratories. Only 27 of them apply quality assurance systems that conform to the requirements of the current EN/ISO standard and are designated by National Reference Laboratory for Salmonella in animals to perform bacteriological examinations in the framework of the programme under the supervision of IDAH-NRL for Salmonella.

#### **6. H.I.P.S.V.H.**

Hygiene Institute for Public Sanitary Veterinary Health is the National Reference Laboratory concerning the expertise for food products of animal origin and feed and it is also designed as the National Reference Laboratory for Salmonella in food and feed –public veterinary health.

#### **4.3. Description and delimitation of the geographical and administrative areas in which the programme is to be implemented:**

The National Control Programme will be implemented throughout Romania, covering all the national territory and will cover all turkey flocks of consisting of at least 250 poultry.

The administrative boundaries are the boundaries of the country. Roumania is administrative divided in 42 counties. There are 42 County Sanitary Veterinary and Food Safety Directorates and 41 County Sanitary veterinary and food Safety Laboratories.

#### **4.4 Measures implemented under the programme**

##### **Bio-Security Measures**

Bio-security is a combination of practices, which are intended to prevent the spread of disease-causing organisms within the poultry farm. Where these are performed in parallel with the sanitation and disinfection procedures, bio-security measures could eradicate or, at least, reduce the level of pathogens to values, at which no hazard of infection would be likely.

The bio-security measures in commercial poultry farms are in accordance to the N.S.V.F.S.A. President Order 147/2006 regarding the bio-security measures in commercial poultry farms and the movements of live poultry, products and poultry by-products.

**Bio-security measures on holdings:**

- Health status of poultry
- On entering to all houses on the farm must be located disinfection barrier
- Control of movement of people
- Transport hygiene
- Feed hygiene
- Water hygiene
- Rodent, insect and bird control
- Cleaning and disinfecting of buildings
- Recording of all events and operations

According to the provision of Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program approved every year by N.S.V.F.S.A. President Order, after each cleaning and disinfections the owner is oblige to take samples to verify the efficient of the disinfection. The official veterinarian takes sample to verify the efficiency of the disinfection only in case of a positive flock with relevant serotypes.

Hygiene measures on poultry farms are also assessed during visits for the collection of official samples and during general visits to premises for other purpose.

**4.4.1. Measures and terms of legislation as regards the registration of holdings:**

All commercial poultry flocks in Romania are registered at the local competent authority according to the provision of National Sanitary Veterinary and Food Safety Authority President **Order no. 16/2010** to approve the sanitary-veterinary procedure of sanitary-veterinary registration/authorization of units/collection centers / holdings of origin and means of transport for health and animal welfare propose, of the units involved in storage and neutralization of animal which are not intended for human consumption and processed products. Poultry holdings shall be registered and sanitary veterinary approved in order to be able to operate.

The legal framework for carrying out commercial activity with poultry and poultry products consists of:

- **Order** of the President of the National Sanitary Veterinary and Food Safety Authority no **144/2006** for the approval of the Sanitary veterinary norm on animal health conditions governing intra-Community trade in, and imports from third countries of, poultry and hatching eggs transposing Council Directive 90/539/EEC on animal health conditions governing intra-Community trade in, and imports from third countries of poultry and hatching eggs.

In order to be sanitary veterinary approved, poultry holdings shall satisfy, in accordance with the legislation in force, the following conditions:

- Appropriate facilities and operation;
- Application of the “Program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” approved by Order of the National Sanitary Veterinary and Food Safety Authority President .
- At least one inspection visit per year by the official veterinarian;
- Additional checks to verify the compliance of the establishment with the hygiene measures and the operation of the establishments.

Each poultry holding receives a distinct approval number, number that can be the same with the one given in compliance with the **Council Regulation no. 2782/75/EEC** on the production and marketing of eggs for hatching and of farmyard poultry chicks.

**4.4.2.Measures and terms of legislation as regards the identification of animals:** Not applicable.

**4.4.3.Measures and terms of legislation as regards the notification of the disease:**

**Commission Decision 176/2005** of 1 March 2005 laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2006/924/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2010/160/EU** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2008/755/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

**COMMISSION DECISION 2009/847/EC** amending Decision 2005/176/EC laying down the codified form and the codes for the notification of animal diseases pursuant to Council Directive 82/894/EEC

The internal notification is made under the provision of **N.S.V.F.S.A. President ORDER no. 79 /2008** to approve the sanitary and veterinary norm on domestic notification and official declaration of animal diseases.

#### **4.4.4. Measures and terms of legislation as regards the measures in case of a positive result:**

In case of suspicion or confirmation of *Salmonella enteritidis* or *Salmonella typhimurium* the NRL/C.S.V.F.S.L. shall notify immediately the N.S.V.F.S.A and the local C.S.V.F.S.D.

When a turkey flock is suspected of being infected with *Salmonella Enteritidis* or *Salmonella Typhimurium* the flock will be investigated. The flock is suspected of being infected when *S. Enteritidis* or *S. Typhimurium* is isolated from samples, carried out privately or as required by either the operator or the Competent Authority as detailed in the Annex to Regulation (EC) No 584/2008. Tissue/organs may be taken from birds as part of the investigation of clinical disease; these cases will be discussed and additional follow up investigation carried out as appropriate, along with advice on *Salmonella* control. Neck skins are sampled at abattoirs under Regulation (EC) No 1441/2007. The potential for cross contamination make the results from abattoir sampling difficult to interpret in relation to a particular flock.

Isolates of *S. Enteritidis* and *S. Typhimurium* will be examined to determine if they are vaccine strains according to the manufacturer's specification. If vaccine strains are confirmed in samples the flock will not be classed as positive for the purposes of establishing the progress towards the target. Although vaccines against *Salmonella* are not currently used in meat turkeys in Romania.

#### **Fattening turkeys**

In most cases it will not be possible to carry out an investigation of the birds in the flock from which the *Salmonella* was isolated as the birds will have been slaughtered shortly after the results become available. A notice may be served by the Competent Authority as necessary requiring the operator to clean and disinfect the building from which the infected flock originated. After cleaning and disinfecting the building the operator may be required to take swabs from a number of sites in the building and submit them to an approved laboratory to be tested for *Salmonella* in order to check on the efficiency of the hygiene measures taken. In cases where *S. Enteritidis* or *S. Typhimurium* was isolated, the cleaning and disinfection may be checked by the Competent Authority or its agent. When *S. Enteritidis* or *S. Typhimurium* is detected in a flock of fattening turkeys, sampling by the Competent Authority will take place in all flocks on the holding in circumstances as laid out in Regulation (EC) No 584/2008.

When a fattening flock is positive for *Salmonella* in the samples taken during the period 3 weeks before slaughter the food business operator for the abattoir where the birds are to be slaughtered should be informed so that arrangements can be made to take measures to reduce the possibility of cross-contamination of other batches, for example arranging slaughter to take place at the end of the day, or before a break, etc.

### **Breeding Turkeys**

When a breeding turkey flock is suspected of being infected with *Salmonella* Enteritidis or *Salmonella* Typhimurium, they will be subject to controls as laid out in Regulation (EC) No 2160/2003 and Community legislation on food hygiene. This applies to breeding flocks from day old through to end of production.

The owner or person responsible for the flock is required to clean and disinfect the building where the infected birds were kept, and provide evidence to the Competent Authority that the cleaning and disinfection has been satisfactory.

Sampling by the Competent Authority will be carried out on all flocks on holdings in case of detection of *Salmonella* Enteritidis or *Salmonella* Typhimurium from samples taken at the hatchery by food business operators or within the frame of official controls, to investigate the origin of infection. Official sampling will be carried out on all holdings/hatcheries where *S. Enteritidis* or *S. Typhimurium* was detected during the previous 12 months as part of the framework of annual official sampling to be carried out under the control of the Competent Authority. This sampling will take place according to the procedure in Annex to Regulation (EC) No 200/2010.

### **General**

Operators with a flock which is positive for *S. Enteritidis* or *S. Typhimurium* will be contacted by the Competent Authority and invited to contact their veterinary adviser for advice on how to reduce or eliminate the *Salmonella*. Advice may include recommendations on management, cleaning and disinfection, pest control, biosecurity, monitoring, and the potential use of other aids in the control of *Salmonella*.

The operator/owner in consultation with his/her veterinarian may consider vaccination of the flock against *Salmonella* with a product which has a marketing authorisation in line with the requirements of Commission Regulation (EC) No. 1177/2006 as regards requirements for the use of specific control methods in the framework of the national programmes for the control of *Salmonella*. Vaccination may only be used as a preventative measure; it is not an alternative to the requirements in Annex II C of Commission Regulation (EC) No 2160/2003 for the use of specific control methods in the framework of the national programmes for the control of *Salmonella*. Antimicrobial treatment may not be used for the control

of *Salmonella* in the national control programme except within the limits set by Commission Regulation (EC) No.1177/2006. For the purposes of establishing the progress towards the target if *S. Enteritidis* or *S. Typhimurium* is isolated from either an operator sample or an official sample the flock is classed as positive. A flock positive for a specific serotype will be recorded only once for that serotype.

In case of suspicion or confirmation of ***Salmonella enteritidis*** or ***Salmonella typhimurium*** the NRL shall notify immediately the N.SV.F.S. and local C.S.V.F.S.D..

### ***Specific requirement concerning fresh meat***

When the turkey are confirmed for the presence of *Salmonella enteritidis* or *Salmonella typhimurium*:

- a. Fresh meat from turkey may be placed on the market on the condition that it meets the requirement of absence of *Salmonella* in 25 grams from the meat.
- b. The requirement laid down in point (a) does not apply to fresh poultry meat destined for heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene.
- c. The criteria laid down in point (a) does not apply to fresh poultry meat destined for industrial heat treatment or another treatment to eliminate salmonella in accordance with Community legislation on food hygiene.

### ***Specific requirements concerning breeding flocks of turkey***

1. Starting with 1 January 2010 the measures laid down below in points 3 to 5 must be taken whenever the analysis of samples carried out in accordance with point 2.2.2.1 and 2.2.2.2 of Annex of the Regulation 200/2010 indicates the presence of *Salmonella enteritidis* or *Salmonella typhimurium* in a breeding flock of turkey in the circumstances set out in point 2.

2. (a) If the competent authority has approved the method of analysis used for samples taken in accordance with part B of the Regulation 2160/2003/EC , it may require that the measures laid down in points 3 to 5 be taken when such analysis detects the presence of *Salmonella enteritidis* or *Salmonella typhimurium*.

(b) Otherwise, the measures laid down in points 3 to 5 must be taken whenever the competent authority confirms a suspicion of the presence of *Salmonella enteritidis* or *Salmonella typhimurium* arising from the analysis of samples carried out in accordance with part B of the Regulation 2160/2003/EC.

### **3. Non-incubated eggs from the flock must be destroyed.**

However, such eggs may be used for human consumption if they are treated in a manner that guarantees the elimination of *Salmonella enteritidis* and *Salmonella typhimurium* in accordance with Community legislation on food hygiene.

4. **All birds**, including day-old chicks, **in the flock must be slaughtered or destroyed so as to reduce as much as possible the risk of spreading Salmonella**. Slaughtering must be carried out in accordance with Community legislation on food hygiene. Products derived from such birds may be placed on the market for human consumption in accordance with Community legislation on food hygiene and, once applicable, part E of the Regulation 2160/2003. If not destined for human consumption, such products must be used or disposed of in accordance with Regulation (EC) No 1774/2002 of the European Parliament and of the Council, laying down health rules concerning animal by-products not intended for human consumption.

5. Where **eggs** for hatching from flocks in which *Salmonella enteritidis* or *Salmonella typhimurium* is present are still **present in a hatchery, they must be destroyed or treated** in accordance with Regulation (EC) No 1774/2002.

### **Control of the use of feed antibiotics by official sampling**

According to the Romanian program of surveillance, prevention and animal disease control, of the diseases transmissible from animals to humans, animal protection and environment protection” and program for surveillance and control in food safety field approved every year by N.S.V.F.S.A. President Order, feeding stuffs intended for poultry nutrition are checked in view to avoid the contamination with *Salmonella* spp. Also, in conformity with the same legislation the feed stuffs are checked in view to detect the use of antibiotics.

Residues examination is performed according to the Roumanian annual plan for examination for residues in live animals and animal origin products .

For broiler, hens, turkeys, other poultry a sample consists on one or more animals depending on the requirements of the analytical methods.

For each category of poultry considered, the minimum number of samples to be taken each year must be at least equal to one per 200 tones of annual production, with a minimum of 100 samples for each group of substances if the annual production of the category of birds considered is over 5 000 tones.

The following breakdown must be respected:

**Group A:** 50 % of the total samples. The equivalent of one fifth of these samples must be taken at farm level. Each sub-group of Group A must be checked each year using a minimum of 5 % of the total number of samples to be collected for Group A. The balance will be allocated according to the experience and background information of Roumania.

**Group B:** 50 % of the total samples, 30 % of samples must be checked for Group B 1 substances (antibiotics and sulfamides)

#### **4.4.6. Control procedures and in particular rules on the movement of animals liable to be affected or contaminated by a given disease and the regular inspection of the holdings or areas concerned.**

According to the provisions of N.S.V.F.S.A. President Order 147/2006, Regulation 2160/2003/EC, the following measures are to be adopted in order to prevent the dissemination of *Salmonella spp.* into commercial holdings. Animals from infected flocks belonging to commercial holdings are to be kept isolated and special conditions apply for removal of these animals. No bird may leave the house concerned unless the competent authority has authorized the slaughter or/and destruction under supervision of slaughter in a slaughterhouse designated by the competent authority. All the birds in the house must be slaughtered in accordance with the provisions of the **REGULATION (EC) No. 853/2004 laying down specific hygiene rules for food of animal origin** in order to reduce as much as possible the risk of spreading salmonella.

#### **4.4.7. Measures and terms of legislation as regards the control (testing, vaccination ...) of the disease.**

The legal basis is the provisions of the **Regulation 2160/2003/EC, Regulation 584/2008/EC** and **Regulation 1177/2006 EC**.

According to the provision of Regulation (EC) no. 1177/2006 CE in Roumania the antimicrobials use for *Salmonella* control is forbidden.

A national *Salmonella* control programme was implemented to comply with Regulation (EC) No 2160/2003 and Regulation (EC) No 584/2008. The national control plan for *Salmonella* in turkey come into effect in January 2010.

***The NCP defines a flock as a single group or multiple groups of chickens which share the same production unit (i.e. using the same air-space or range area). Where housing systems are not typical, the situation is likely to be assessed on a case by case basis. Multiple groups of chickens which have 'beak-to-beak' contact (inside or outside the house) are likely to be treated as a single flock.***

**Operators** will be required to implement the sampling programme in the Annex to EC Regulation 584/2008 (**self-control sampling**). For convenience the 'Sampling protocol' is repeated in **THE SAMPLING PROTOCOL**, showed below.

The sampling programme will be in accordance to Regulation **2160/2003 EC** and **Regulation EC 584/2008**. We have also taken into account the Regulation **1177/2006 EC** and **Regulation 213/2009**.

The programme was elaborated in compliance with the requirements of Council Regulation **(EC) No 584/2008** implementing Regulation (EC) No 2160/2003 of the European Parliament as regards a Community target for the reduction of the prevalence of *Salmonella enteritidis* and *Salmonella typhimurium* in turkeys.



Operators will be required to implement the sampling programme in the Annex to Regulation (EC) 584/2008 and Regulation (EC) 200/2010. For convenience the 'Sampling protocol' is repeated below.

For **fattening turkeys** at least two pairs of boot sock/swabs, or one pair of boot sock/swabs and a dust sample will be taken by the operator within the period of three weeks before the birds are due for slaughter. Where possible, the samples will be taken in sufficient time for the laboratory results to be known before the birds are transported to the slaughterhouse.

For **breeding turkeys**, samples for the detection of *Salmonella* will be taken from rearing turkey breeding flocks at day-old, at four weeks of age and two weeks before moving to the laying phase or laying unit. In adult breeding flocks, samples shall be taken at least every third week during the laying period at the holding or at the hatchery. The samples in adult breeding flocks shall be taken in accordance with the provisions laid down in Annex to Regulation (EC) No 200/2010.

Each year official samples will be taken by the local Competent Authority, or other authorized agent, acting on behalf of the Competent Authority. When an official sample is taken it may replace the sample required to be taken by the operator. Sampling to verify the achievement of the target will be as detailed in "Sampling protocol" in the Annex to Commission Regulation (EC) No. 584/2008.

## **SAMPLING PROTOCOL.**

For each flock\*

### ***Sampling at the hatchery***

Sampling shall occur at the hatchery in accordance with the provisions laid down in point 2.2.1 of the Annex to Regulation (EC) No 584/2008.

### ***Sampling at the holding for each flock\****

#### **Breeding turkeys**

Samples from adult turkey breeding flocks shall be taken in accordance with the provisions laid down in Annex to Regulation (EC) No. 584/2008.

| <b>Phase of production</b>                      | <b>Type of sample</b>  |
|---|--|
| Every third week from each adult breeding flock | 5 pairs of boot swabs from each flock pooled into a minimum of two composite samples.<br>OR<br>One pair of boot swabs and a dust sample to be analysed as 2 pools. |

Samples shall also be taken at the initiative of the food business operator during the rearing phase at day old, at 4 weeks of age and at two weeks before moving to the laying phase or laying unit.

### **Fattening turkeys**

At least two pairs of boot/sock swabs or one boot/sock swab and one dust sample shall be taken. For free range flocks of turkeys, samples shall only be collected in the area inside the house. All boot/sock swabs may be pooled into one sample. In flocks with less than 100 turkeys, where it is not possible to use boot/sock swabs as access to the houses is not possible, they may be replaced by hand drag swabs, where the boot swabs or socks are worn over gloved hands and rubbed over surfaces contaminated with fresh faeces, or if not feasible, by other sampling techniques for faeces fit for the intended purpose. Before putting on the boot/sock swabs, their surface shall be moistened with sterile water or any other diluent approved by the national reference laboratory referred to in Article 11 of Regulation (EC) No 2160/2003. The use of farm water containing antimicrobials or additional disinfectants shall be prohibited. The recommended way to moisten boot swabs shall be to pour the liquid inside before putting them on.

It shall be ensured that all sections in a house are represented in the sampling in a proportionate way. If 2 pairs of boot swabs are taken, each pair should cover about 50 % of the area of the house and that at least 100 steps should be taken with each pair of boot swabs. Alternatively, if one pair of boot swabs is taken, the sampling must be carried out so as to ensure covering 100 % of the area of the house if combined with a dust sample, collected from multiple places throughout the house from surfaces with visible presence of dust. On completion of sampling the boot/sock swabs shall be carefully removed so as not to dislodge adherent material. Boot swabs may be inverted to retain material. They shall be placed in a bag or pot and labeled. The Competent Authority shall supervise education of the food business operators to guarantee the correct application of the sampling protocol. In the case of sampling by the Competent Authority because of suspicion of *Salmonella* infection in a flock on that holding and in any other case considered appropriate, the Competent Authority shall satisfy itself by conducting further tests as appropriate so that the results of examinations for *Salmonella* in flocks of turkeys are not affected by the use of antimicrobials in those flocks. Where the presence of *Salmonella* Enteritidis and *Salmonella* Typhimurium is not detected but antimicrobials or bacterial growth inhibitory effect are detected it shall be considered as an infected flock of turkeys for the purpose of the Community target referred to in Article 1 of Commission Regulation 584/2008.

### **Official sampling**

In accordance with Regulation (EC) No 584/2008 the Competent Authority will select each year at random:

- All flocks on at least 10% of holdings with more than 500 **fattening turkeys**.
- All flocks on at least 10% of holdings with at least 250 adult **breeding turkeys** between 30 and 45 weeks of age, but including all holdings where *S. Enteritidis* or *S. Typhimurium* has been detected during the previous 12 months and all holdings with elite, great grandparents and grandparent breeding turkeys. This sampling may also take place at the hatchery
- Each time the Competent Authority considers it necessary

In the case of fattening turkeys the flock should be within 3 weeks of going for slaughter. The flocks will be sampled in accordance with the Annex in Regulation (EC) No 584/2008..

The use of antimicrobials (as defined in Regulation (EC) No 1177/2006) will be checked when the official sample is taken. If the flock is under antimicrobial medication for animal health or animal welfare reasons the flock will be sampled again after the period of withdrawal considered suitable by the Competent Authority and taking into account the product Marketing Authorisation. Flock owners are required to keep records of antimicrobial use and to make these records available .

The records of samples taken by the operator will be made available for inspection to the Competent Authority or its agent and will provide details of the identity of the flock sampled, date of sample, slaughter date, type of sample, laboratory carrying out the examination, and the result. The number of flocks on the holding and the number of birds.

Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs requires poultry abattoirs and establishments handling processed meat to demonstrate compliance with microbiological criteria for *Salmonella* that includes undertaking microbiological testing for *Salmonella* and corrective action when the criteria are not met. The sampling frequencies are prescribed in the Regulation but do not apply to establishments producing small quantities.

#### **4. Measures of the submitted programme**

Samples taken as above shall preferably be sent by express mail or courier within 24 hours of collection to a authorized laboratory for the detection of *Salmonella*. If not sent within 24 hours the sample may be stored refrigerated for up to a maximum of 24 hours and then sent to the laboratory so as to arrive within 48 hours of the time of collection of the sample At the laboratory samples shall be kept refrigerated until examination unless this examination is begun within 3 hours of arrival of the sample at the laboratory. Examination shall be commenced within 48 hours following receipt.

### *Boot swabs and dust samples:*

(a) The pair(s) of boot/sock swabs and dust sample (fabric swab) shall be carefully unpacked to avoid dislodging adherent faecal material or loose dust material and placed in 225 ml of BPW which has been pre-warmed to room temperature.

(b) The boot/socks and fabric swab shall be fully submersed in BPW to provide sufficient free liquid around the sample for migration of *Salmonella* away from the sample and therefore more BPW may be added, if necessary.

Separate preparations must be made of the boot swabs and the fabric swab.

(c) Where five pairs of boot/sock swabs are pooled into two samples, each pooled sample must be placed in of 225 ml of BPW, or more if necessary, to fully submerge the sample and provide sufficient free liquid around the sample for migration of *Salmonella* away from the sample.

(d) Swirl to fully saturate the sample and continue the culture by using the detection method described in point 3.2.

### *Other faecal material samples:*

(a) The faeces samples shall be pooled and thoroughly mixed and a 25 g sub-sample shall be collected for culture.

(b) The 25 g sub-sample shall be added to 225 ml of BPW which has been pre-warmed to room temperature.

(c) The culture of the sample shall be continued by using the detection method described below.

## **Detection method**

The detection of the relevant *Salmonella* serotypes shall be carried out according to Amendment 1 of EN/ISO 6579-2002/Amd1:2007. 'Microbiology of food and animal feeding stuffs – Horizontal method for the detection of *Salmonella* spp. — Amendment 1: Annex D: Detection of *Salmonella* spp. in animal faeces and in environmental samples from the primary production stage'.

As regards the boot swabs samples, dust samples and other faecal material samples, the incubated BPW enrichment broth for future culture may be pooled. To do so, incubate both samples in BPW . Take 1 ml of incubated broth from each sample and mix thoroughly, then take 0,1 ml of the mixture and inoculate the modified semi-solid Rappaport-Vassiliadis (MSRV) plates.

The samples in BPW must not be shaken, swirled or otherwise agitated after incubation as this releases inhibitory particulates and reduces subsequent isolation in MSRV.

## **Serotyping**

At least one isolate from each positive sample shall be serotyped, following the Kaufmann-White scheme.

#### **Storage of strains:**

At least one isolated strain per house and per year shall be collected by the NRL and stored for future phagetyping or antimicrobial susceptibility testing, using the normal methods for culture collection, which must ensure integrity of the strains for a minimum of two years.

#### **Use of vaccines**

Live Salmonella vaccines shall not be used in the framework of national control programme where the manufacturer does not provide an appropriate method to distinguish bacteriological wild-type strains of salmonella from vaccine strains.

Live salmonella vaccines shall not be used in the framework of national control programmes in breeding hens during production unless the safety of the use has been demonstrated and they are authorized for such purpose in accordance with Directive 2001/82/EC.

#### **4.4.8. Measures and terms of legislation as regards the compensation for owners of slaughtered and killed animals:**

There is no financial assistance provided in the context of the control programme.

#### **4.4.9. Information and assessment on bio-security measures management and infrastructure in place in the flocks/holdings involved**

This information was given in the point 4.4.

#### **Costs of the programme**

This programme is in accordance with Council Decision 2009/470/ EEC, Commission Decision 2008/425/EC and Council Decision 92/65/ EEC. The detailed financial costs for the national control programme of Salmonella in breeding hens are laid down in point 8.

#### **Benefits of the programme**

The overall aim of the Salmonella National Control Programme is to control the occurrence of Salmonella in turkey sector on a very low level and thereby protect humans against infection with food-borne salmonellas. It has been known that poultry often harbor latent infections with Salmonella, which may pose a serious human health risk.

The anticipated benefits of this programme reducing the relevant *Salmonella* are the minimising of human health problems and a consequent reduction in suffering, mortality and health service costs.

**5. Data on the epidemiological evolution during the last year:**

**6.1. Evolution of zoonotic salmonellosis**

### 6.1.1 Data on evolution of zoonotic salmonellosis

Year: 2010

Situation on date: 31.12.2010

Animal species: turkey rearing for meat

Disease: zoonotic salmonellosis

| Region<br>(County+NUTS CODE) | Type of<br>flocks(b)           | Total<br>number<br>of flocks<br>© | Total number of<br>animals | Total number<br>of flocks<br>under the<br>programme | Total number<br>of animals<br>under the<br>programme | Number of<br>flocks<br>checked(d) | Number of positive<br>flocks (e)                                     |                           | Number of flocks<br>depopulated                                     |                       | Total number of animals<br>slaughtered or destroyed         |                           | Quantity of the eggs<br>destroyed (number<br>or kg)                 |                           | Quantity of eggs<br>channelled to egg<br>products (number or<br>kg) |                           |
|------------------------------|--------------------------------|-----------------------------------|----------------------------|---|--|-----------------------------------|--|---------------------------|---|-----------------------|---|---------------------------|---|---------------------------|---|---------------------------|
|                              |                                |                                   |                            |   |  |                                   | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotype<br>s(g) | Serotype<br>s<br>targeted<br>in the<br>control<br>program<br>me (f) | Other<br>serotypes(g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>serotype<br>s(g) | Serotype<br>s<br>targeted<br>in the<br>control<br>program<br>me (f) | Other<br>serotyp<br>es(g) | Serotyp<br>es<br>targeted<br>in the<br>control<br>progra<br>mme (f) | Other<br>serotype<br>s(g) |
| BRASOV-RO122                 | Turkeys<br>rearing<br>for meat | 52                                | 299047                     | 52  | 299047   | 35                                | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                         |
| DOLJ-RO411                   | Turkeys<br>rearing<br>for meat | 3                                 | 15300                      | 1   | 5100   | 1                                 | 0  | 0                         | 0   | 0                     | 0   | 0                         | 0   | 0                         | 0   | 0                         |
| MURES-RO125                  | Turkeys<br>rearing<br>for meat | 18                                | 80000                      | 18  | 80,000   | 18                                | 0  | 7                         | 0   | 7                     | 0   | 0                         | 0   | 0                         | 0   | 0                         |
| <b>TOTAL</b>                 |                                | <b>73</b>                         | <b>394347</b>              | <b>71</b>   | <b>384147</b>  | <b>54</b>                         | <b>0</b>   | <b>7</b>                  | <b>0</b>  | <b>7</b>              | <b>0</b>  | <b>0</b>                  | <b>0</b>  | <b>0</b>                  | <b>0</b>  | <b>0</b>                  |

Year: 2011

Situation on date: 31.12.2011

Animal species: turkey rearing for meat

Disease: zoonotic salmonellosis

| Region<br>(County+NUTS<br>CODE) | Type of<br>flocks(b) | Total<br>number<br>of flocks<br>© | Total<br>number<br>of<br>animals | Total<br>number of<br>flocks<br>under the<br>programme | Total<br>number of<br>animals<br>under the<br>programme | Number of<br>flocks<br>checked(d) | Number of positive flocks<br>(e)                            |                           | Number of flocks<br>depopulated                                      |                           | Total number of animals<br>slaughtered or<br>destroyed      |                           | Quantity of the<br>eggs destroyed<br>(number or kg)                  |                           | Quantity of<br>eggs<br>channelled to<br>egg products<br>(number or kg)                 |                           |
|---------------------------------|----------------------|-----------------------------------|----------------------------------|--|---|-----------------------------------|---|---------------------------|--|---------------------------|---|---------------------------|--|---------------------------|--|---------------------------|
|                                 |                      |                                   |                                  |  |   |                                   | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>serotypes(<br>g) | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotypes(<br>g) | Serotypes<br>targeted in<br>the control<br>programme<br>(f) | Other<br>serotyp<br>es(g) | Serotyp<br>es<br>targete<br>d in the<br>control<br>progra<br>mme (f) | Other<br>serotyp<br>es(g) | Sero<br>type<br>s<br>targe<br>ted<br>in the<br>cont<br>rol<br>prog<br>ram<br>me<br>(f) | Other<br>serotyp<br>es(g) |
| RO 122 BV                       | Turkey               | 96                                | 389982                           | 86   | 329982  | 39                                | 0   | 10                        | 0  | 10                        | 0   | 68983                     | 0  | 0                         | 0  | 0                         |
| RO411-DJ                        | Turkey               | 12                                | 39000                            | 12   | 39000   | 0                                 | 0   | 0                         | 0  | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         |
| RO125-MS                        | Turkey               | 3                                 | 14850                            | 1  | 4950  | 1                                 | 0   | 0                         | 0  | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         |
| RO412-GJ                        | Turkey               | 0                                 | 0                                | 0  | 0   | 0                                 | 0   | 0                         | 0  | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         |
| IDAH                            | Turkey               | 0                                 | 0                                | 0  | 0   | 0                                 | 0   | 0                         | 0  | 0                         | 0   | 0                         | 0  | 0                         | 0  | 0                         |
| <b>Total</b>                    | <b>Turkey</b>        | <b>111</b>                        | <b>443832</b>                    | <b>99</b>  | <b>373932</b>   | <b>40</b>                         | <b>0</b>  | <b>10</b>                 | <b>0</b>   | <b>10</b>                 | <b>0</b>  | <b>68983</b>              | <b>0</b>   | <b>0</b>                  | <b>0</b>   | <b>0</b>                  |

## 6.2 Data on vaccination programme

Not applicable



## 7. Targets

### 7.1. Targets related to testing :to investigate the presence of Salmonella in turkey flocks

**Disease (a): zoonotic Salmonella**

**Animal species:**

| Region (b)   | Type of the test (c) | Target population (d) | Type of sample (e)                 | Objective (f) | Number of planned tests |
|--------------|----------------------|-----------------------|------------------------------------|---------------|-------------------------|
| All regions  | Bacteriological      | turkey                | Faeces<br>boot/sock<br>swabs, dust | Surveillance  | 100                     |
|              | Bacteriological      | turkey                | Isolate of<br>Salmonella           | Confirmation  | 50                      |
| <b>Total</b> | -                    | -                     | -                                  | -             | 150                     |

(a) Disease and species if necessary

(b) Region as defined in the approved eradication programme of the Member State

(c) Description of the test

(d) Specification of the targeted species and the categories of targeted animals

(e) Description of the sample (faeces)

(f) Description of the objective ( e.g. qualification, surveillance, confirmation of suspected cases, monitoring of campaigns, seroconversion, control on deleted vaccines, testing of vaccine , control of vaccination

## 7.1.2. Targets on testing of turkey flocks

Year: 2012

Animal species: Gallus gallus

Disease/infection(a): Zoonotic Salmonella

| Region      | Type of flock <sup>(b)</sup> | Total number of flocks <sup>(c)</sup> | Total number of animals | Total n° of flocks under the programme | Total n° of animals under the programme | Expected n° of flocks to be checked <sup>(d)</sup> | Number of flocks <sup>(e)</sup> expected to be positive <sup>(a)</sup> |      |      | Number of flocks expected to be depopulated |      | Total n° of animals expected to be slaughtered or destroyed <sup>(f)</sup> |      | Expected quantity of eggs to be destroyed (number or kg) <sup>(d)</sup> | Expected quantity of eggs channelled to egg products (number or kg.) <sup>(d)</sup> |    |      |
|-------------|------------------------------|---------------------------------------|-------------------------|--|---|--|--|------|------|---|------|--|------|---|---|----|------|
|             |                              |                                       |                         |  |   |  | (a1)   | (a2) | (a3) | a4  | (a3) | (a4)   | (a3) |   | (a4)  | a3 | (a4) |
| All regions | turkey                       | 200                                   | 750000                  | 200                                    | 750000                                  | 75   | 2  | 2    | 20   | 4   | 20   | 1000   | 0    | 10000   | 0   | 0  | 0    |
| Total       | turkey                       | 200                                   | 750000                  | 200                                    | 750000                                  | 75   | 2  | 2    | 20   | 4   | 20   | 1000   | 0    | 10000   | 0   | 0  | 0    |

(a) For zoonotic salmonella indicate the serotypes covered by the control programmes: (a1) for Salmonella Enteritidis, (a2) for Salmonella Typhimurium, (a3) for other serotypes-specify as appropriate, (a4) for Salmonella Enteritidis or Salmonella Typhimurium.

(b) For example, breeding flocks (rearing, adult flocks), production flocks, laying hen flocks, etc. Flocks equals herds or as appropriate.

(c) Total number of flocks existing in the region including eligible flocks and non-eligible flocks for the programme.

(d) Check means to perform a flock level test under the programme for the presence of Salmonella. In this column a flock should not be counted twice even if it has been checked more than once.

(e) If a flock has been checked, in accordance with footnote (d), more than once, a positive sample should be taken into account only once.

## 7.2. Targets on vaccination

### 7.2.1. Targets on vaccination of turkeys flocks- not applicable

Disease<sup>(a)</sup>: Zoonotic Salmonella

Animal species: gallus gallus

| Region | Total number of herds in | Total number of animals in | Targets on vaccination programme |             |               |                |
|--------|--------------------------|----------------------------|----------------------------------|-------------|---------------|----------------|
|        |                          |                            | No of herds in                   | No of herds | No of animals | No of doses of |

|              | vaccination or programme | vaccination programme | vaccination programme | expected to be vaccinated | expected to be vaccinated | vaccine expected to be administered |
|--------------|--------------------------|-----------------------|-----------------------|---------------------------|---------------------------|-------------------------------------|
| All regions  | -                        | -                     | -                     | -                         | -                         | -                                   |
| <b>Total</b> | -                        | -                     | -                     | -                         | -                         | -                                   |

**The vaccination is not mandatory and the costs regarding purchase of vaccine doses and the vaccination are incurred by the business operators.**

## 8. Detailed analysis of the cost of the programme<sup>1</sup>

Costs mentioned below are estimated for a one-year period (1 of January 2013-31 of December 2013 ) for samples harvest on official control

| <i>Costs related to</i>          | <i>Specification</i>   | <i>Number of units</i> | <i>Unitary cost in €</i> | <i>Total amount in €</i> | <i>Community funding requested (yes/no)</i> |
|----------------------------------|--|------------------------|--------------------------|--------------------------|---|
| <b>1. Testing</b>                |  |                        |                          |                          |   |
| <b>1.1. Cost of the analysis</b> | Test: bacteriological exam for Salmonella spp.detection  | 100                    | 16,70                    | 1670                     | YES   |
|                                  | Test : Salmonella spp. serotyping  | 50                     | 64.97                    | 3248.50                  | YES   |
|                                  | Test : the detection of antimicrobials or bacterial growth inhibitory effect in tissues from birds from flocks tested for Salmonella | 100                    | 47.44                    | 4744                     | YES   |
| <b>1.2. Cost of sampling</b>     | Official sample harvest  | 100                    | 0,5                      | 50                       | YES   |
|                                  | Disposable sterile containers for sampling of faeces   | 100                    | 0,5                      | 50                       | NO  |
|                                  | One use gloves-pairs   | 100                    | 0,1                      | 10                       | NO  |
|                                  | Boot swabs   | 200                    | 2                        | 400                      | NO  |
| <b>1.3. Other costs</b>          | Overcoats  | 100                    | 1                        | 100                      | NO  |
| <b>2. Vaccination</b>            |  |                        |                          |                          |   |
| <b>2.1. Purchase of vaccine</b>  |  |                        |                          |                          |   |
| <b>2.2. Distribution costs</b>   |  |                        |                          |                          |   |
| <b>2.3. Administering costs</b>  |  |                        |                          |                          |   |

|  |   |             |            |                |            |
|--|---|-------------|------------|----------------|------------|
| <b>2.4. Control costs</b>  |   |             |            |                |            |
| <b>3. Slaughter and destruction</b>  |   |             |            |                |            |
| <b>3.1. Compensation of animals</b>  | Compensation for the value of a parent breeding turkey bird of <i>Meleagris gallopavo</i> culled  | <b>500</b>  | <b>10</b>  | <b>5000</b>    | <b>YES</b> |
|  | Compensation for hatching eggs of parent breeding <i>Meleagris gallopavo</i> per hatching egg destroyed;                                | <b>1000</b> | <b>0,5</b> | <b>500</b>     | <b>YES</b> |
| <b>3.2. Transport costs</b>  |   |             |            |                |            |
| <b>3.3. Destruction costs</b>  |   |             |            |                |            |
| <b>3.4. Loss in case of slaughtering</b>   |   |             |            |                |            |
| <b>3.5 Costs from treatment of products (eggs, hatching eggs, heat treat of broilers, etc)</b> |   |             |            |                |            |
| <b>4. Cleaning and disinfection</b>  |   |             |            |                |            |
|  | Test: Bacteriological test to verify the efficiency of disinfection of poultry houses after depopulation of a Salmonella-positive flock | <b>100</b>  | <b>10</b>  | <b>1000</b>    | <b>YES</b> |
| <b>5. Salaries (staff contracted for the programme only)</b>                                   |   |             |            |                |            |
|  |   |             |            |                |            |
| <b>6. Consumables and specific equipment</b>   |   |             |            |                |            |
| <b>7. Other costs</b>  |   |             |            |                |            |
| <b>TOTAL</b>   |   |             |            | <b>16212.5</b> | <b>YES</b> |



**EUROPEAN COMMISSION**  
 HEALTH AND CONSUMERS DIRECTORATE-GENERAL  
 Veterinary and International affairs  
**Veterinary control programmes**

Brussels, 10 August 2012

**Subject: 2013 Salmonella control programmes, Romania**

Following the assessment of all of your programmes there is a need for additional information/clarifications/amendment of the programmes in the following points:

**SCP laying hens**

**Programme element and relevant criteria**

The prevalence trend is not positive, apparently it showed an increase from 0.2% in 2009 to 1.9% in 2011. A better trend analysis of laying hens should be provided.  
 Could you please explain this?

Because of an error of translation in the EU legislation from English into romanian of the word “flock” in 2009 were tested all the holdings , but not also all flocks in he same holding. After 2010 FVO mission, we start to implement correct the words :”Flock” and we start to tested all the flocks of laying hens. Increasing the number of tests, in 2010 and 2011,determine to increase the number of positive flock. Because of this increase of the number of flocks we take into consideration to vaccinate mandatory fos Salmonella enteritidis and Salmonella typhimurium.

Point 1.1 in Part A shall be completed.

We completed point 1.1from part A

Specific references to the Salmonella results obtained in the framework of the programmes, in animal populations, products and humans should be included.

Information from 2011: E.g. Table 6.4 shall be completed.

Table 6.4 is regarding the vaccinatın. In Romania the vaccination against Salmonella is not mandatory.

**Programme element and relevant criteria**

***Proposed changes and/or additions that may be required:***

Monophasic ST should be included in the objectives.

Historical data on the epidemiological evolution of the disease (sampling, vaccination, analyses etc.) should be completed.

Improve information in some part of the programme (structure of production) 2.1 , 2.2.

A better trend analysis of laying hens should be provided.

**SCP breeding flocks**

**Programme element and relevant criteria**

***The evolution of the disease over the past 5 years as demonstrated by the data provided showing a sufficiently positive trend***

Monophasic ST should be included in the objectives.

The monophasic ST was also included in the objectives.

Historical data on the epidemiological evolution of the disease (sampling, vaccination, analyses etc.) should be completed.- We also completed the epidemiological evolution with historical data

Improve information in some part of the programme (structure of production) 2.1 , 2.2, complete all Tables.

Specific references to the Salmonella results obtained in the framework of the programmes, in animal populations, products and humans should be included.

Information from 2011: E.g. Table 6.4 shall be completed.- In Romania the vaccination is not mandatory in breeding flocks.

**SCP broilers**

### **Programme element and relevant criteria**

#### ***Proposed changes and/or additions that may be required:***

Monophasic ST should be included in the objectives.

The monophasic ST was also included in the objectives.

Historical data on the epidemiological evolution of the disease (sampling, vaccination, analyses etc.) should be completed.

We also completed the epidemiological evolution with historical data

All flocks shall be sampled in 2013.

All broiler flocks will be tested in 2013. The National Control Programme will be implemented all throughout Romania and will apply to all operators with just two exceptions:

- Holdings with capacity of less than 500 chickens present at any one time, where the operator supplies small quantities direct to the consumer (i.e. farm gate sales) or via local 1 retailers which only supply the final consumer (essentially householders).
- Where all production is for private domestic use only (i.e. the meat is not being sold on the market).

Improve information in some part of the programme (structure of production) 2.1 , 2.2, complete all Tables.-We improve the information

### **Programme element and relevant criteria**

The prevalence of SE/ST is low and below Community target, but prevalence is increasing. Reported prevalences are: 2009: 0.01%, 2010: 0.2%, and 2011: 0.7%.

In 2011, only 1,535 of 4,910 flocks were checked. Could you please explain this?

*A total no of 4910 flocks were checked in total (both FBO initiative and official control). A no of 1535 broiler flock were tested at official control*

Point 1.1 in Part A shall be completed.

Specific references to the Salmonella results obtained in the framework of the programmes, in animal populations, products and humans should be included.

Information from 2011: E.g. Table 6.4 shall be completed.

Table 6.4 is regarding the vaccination programme. In Romania the broiler flock are not vaccinated against Salmonella



## SCP breeding and fattening turkeys

### **Programme element and relevant criteria**

Target is clearly set up, but the target is only for holdings with more than 500 birds. This shall be in line with Regulation. 2160/2003/EC.

The National Control Programme will be implemented all throughout Romania and will apply to all operators with just two exceptions:

- Holdings with capacity of less than 500 fattening turkeys present at any one time, where the operator supplies small quantities direct to the consumer (i.e. farm gate sales) or via local 1 retailers which only supply the final consumer (essentially householders).
- Where all production is for private domestic use only (i.e. the meat is not being sold on the market)

From the information given, all flocks shall be sampled in 2013 (75/200)?

**All broiler flocks will be tested in 2013.**

### **Programme element and relevant criteria**

Data shall be presented for breeding turkey flocks. Data are presented for 2010 and 2011 for fattening flocks (no positive flocks), but the coverage shall be improved.

In Romania there are not at this time any breeding turkey holding with breeding turkey flocks. In 2010 54 of 71 flocks were checked, and in 2011 less than half of the flocks (40/99).

In 2010 there were tested all 71 flocks (17 of them by the FBO, 54 in official control) in official control the 54. In 2011 all the fattening turkey flocks were tested, from 99 flocks 40 were tested by the official control and the rest by the FBO.

Point 1.1 in Part A shall be described.

Specific references to the Salmonella results obtained in the framework of the programmes, in animal populations, products and humans should be included.

For 2011: E.g. Table 6.4 shall be completed.

In Romania the fattening turkeys are not vaccinated against Salmonella.

### **Programme element and relevant criteria**

Indicator is expected number of positive flocks (programme relevant serotypes)

There were no positive fattening turkey flock since the beginning of the programme.

Indicators for FBO and OC should be given separately.

**Programme element and relevant criteria**

*Proposed changes and/or additions that may be required:*

Monophasic ST should be included in the objectives.

There were included also

Historical data on the epidemiological evolution of the disease (sampling, vaccination, analyses etc.) should be completed.

All flocks under the programme shall be sampled in 2013

All flocks of fattening turkeys will be tested in 2013.

Improve information in some part of the programme (structure of production) 2.1 , 2.2, complete all Tables.

Please, insert the answers to the previous questions in this document and please submit the revised programmes.

Thank you very much for your cooperation.



## **AUTORITATEA NAȚIONALĂ SANITARĂ VETERINARĂ ȘI PENTRU SIGURANȚA ALIMENTELOR**

**DIRECȚIA GENERALĂ SANITARĂ VETERINARĂ ȘI PENTRU SIGURANȚA  
LANȚULUI ALIMENTAR**

### DECLARATION

General Sanitary Veterinary and Food Chain Safety Directorate confirm that provision of below mentioned legislation will be followed /implemented in 2013 in the context of Salmonella National Control Programmes:

1. In the frame of Salmonella NCP in laying flocks of Gallus gallus the provisions of paragraph 1 and 2 (frequency of sampling) and 3,4 (results and reporting) of Annex of Commission Regulation (EC) no 517/2011 (particular provision on exceptional cases ) are implemented;
2. In the frame of Salmonella NCP in breeding flocks of Gallus gallus the provisions of paragraph 1 and 2 (frequency of sampling) and 4 (results and reporting) of Annex of Commission Regulation (EC) no 200/2010 and 213/2009 (particular provision on exceptional cases ) are implemented;
3. In the frame of Salmonella NCP in broilers the provisions of Commission Regulation 646/2007 paragraph 1/2/4 are implemented.
4. In the frame of Salmonella NCP in turkeys the provisions of Commission Regulation 584/2007 paragraph 1/2/4 are implemented.

Sincerely yours,

**GENERAL DIRECTOR**

**Dr Ioan BOGOLIN**

Director : Dr. Nicolae LAZĂR  
Șef Serviciu Ioana NEGHIRLĂ